

# Census of Vascular Plants, up-dated to 29 Nov 2004

Symbols in left margin:

e endemic in Tasmania

t within Australia, occurs only in Tasmania

x considered to be extinct in Tasmania

i introduced and naturalised in Tasmania

# sparingly naturalised or known from only one or two populations or collections

no symbol indicates that the taxon is native in Tasmania and the Australian mainland.

n a name newly added to the Census in 2004

In the right margin there is a part and page reference to The Student's Flora of Tasmania.

## Dicotyledonae

### ACANTHACEAE

- i **Acanthus mollis** L., Sp. Pl. 2: 639 (1753) 3:539

### AIZOACEAE (FICOIDEAE)

- i# **Aptenia cordifolia** (L.f.) Schwantes, Gartenflora 77: 69 (1928)
- i **Carpobrotus aequilaterus** (Haw.) N.E.Br., J. Bot. 66: 324 (1928)
- i **Carpobrotus edulis** (L.) N.E.Br. in E.Phillips, Gen. S.Afr. Flow. Pl. 249 (1926) 2:238
- Carpobrotus rossii** (Haw.) Schwantes, Gartenflora 77: 68 (1928) 2:238
- Disphyma australe sensu W.M.Curtis (1963) = **Disphyma crassifolium** 2:239
- Disphyma crassifolium** (L.) L.Bolus, Fl. Pl. S.Afr. 7: t.276 (1927) 2:239
- subsp. **clavellatum** (Haw.) Chinnock, Fl. S.Aust. ed.4, 1: 194 (1986)
- i **Galenia pubescens** (Ecklon & Zeyher) Druce, Bot. Soc. Exch. Club Brit. Isles 1916: 624 (1917) 2:241
- Galenia secunda sensu W.M.Curtis (1963) = **Galenia pubescens** 2:241
- i **Lampranthus glaucus** (L.) N.E.Br., Gard. Chron. 87: 252 (1930) 2:240
- Mesembryanthemum aequilaterale sensu L.Rodway (1903) = **Carpobrotus rossii** 2:238
- Mesembryanthemum australe sensu L.Rodway (1903) = **Disphyma crassifolium** 2:239
- i **Mesembryanthemum crystallinum** L., Sp. Pl. 480 (1753)
- Tetragonia expansa Murr. = **Tetragonia tetragonoides** 2:240
- Tetragonia implexicoma** (Miq.) Hook.f., Fl. Tasm. 1: 148 (1856) 2:241
- Tetragonia tetragonoides** (Pallas) Kuntze, Revis. Gen. Pl. 1: 264 (1891) 2:240

### AMARANTHACEAE

- Alternanthera denticulata** R.Br., Prodr. 417 (1810) 3:568
- i **Amaranthus albus** L., Syst. Nat. ed.10: 1268 (1759)
- i **Amaranthus deflexus** L., Mant. Pl. 2: 295 (1771) 3:567
- i# **Amaranthus graecizans** L., Sp. Pl. 990 (1753)
- Amaranthus hybridus sensu W.M.Curtis (1967) = **Amaranthus powellii** 3:566
- i# **Amaranthus powellii** S.Watson, Proc. Amer. Acad. 10: 347 (1875)
- Amaranthus retroflexus sensu W.M.Curtis (1967) = **Amaranthus powellii** 3:566

|                                    |  |       |
|------------------------------------|--|-------|
| i#                                 | <b>Amaranthus spinosus</b> L., Sp. Pl. 1405 (1753)   |       |
|                                    | <b>Hemichroa pentandra</b> R.Br., Prodr. 409 (1810)  | 3:576 |
|                                    | <b>Ptilotus spathulatus</b> (R.Br.) Poiret ex F.Muell., Pap. Proc. R. Soc. Tasm. 77 (1874)           | 3:567 |
| <br><b>APIACEAE (UMBELLIFERAE)</b> |  |       |
|                                    | Aciphylla procumbens (F.Muell.) Benth. = <b>Anisotome procumbens</b>                                 | 2:261 |
|                                    | <b>Actinotus bellidioides</b> (Hook.f.) Benth., Fl. Austral. 3: 369 (1867)                           | 2:253 |
| e                                  | <b>Actinotus moorei</b> F.Muell. & Rodway, Pap. Proc. R. Soc. Tasm. for 1894-95: 65 (1896)           | 2:254 |
| t                                  | <b>Actinotus suffocatus</b> (Hook.f.) Rodway, Pap. Proc. R. Soc. Tasm. 1893: 180 (1894)              | 2:254 |
| i                                  | <b>Aegopodium podagraria</b> L., Sp. Pl. 265 (1753)  |       |
| i                                  | <b>Ammi majus</b> L., Sp. Pl. 243 (1753)   |       |
| e                                  | <b>Anisotome procumbens</b> (F.Muell.) C.J.Webb in Barlow, Fl. & Fauna Alpine Australasia 395 (1986) | 2:261 |
| it                                 | <b>Anthriscus caucalis</b> Bieb., Fl. Taur-Cauc. 1: 230 (1808)                                       |       |
|                                    | <b>Apium annuum</b> P.S.Short, J. Adelaide Bot. Gard. 1: 230 (1979)                                  |       |
|                                    | <b>Apium insulare</b> P.S.Short, J. Adelaide Bot. Gard. 1: 228 (1979)                                |       |
|                                    | <b>Apium prostratum</b> Labill. ex Vent., Jard. Malmaison 2: t.81 (1805)                             | 2:255 |
|                                    | var. <b>filiforme</b> (A.Rich.) Kirk, Stud. Fl. N.Z. 196 (1899)                                      |       |
|                                    | var. <b>prostratum</b>   |       |
|                                    | <b>Centella cordifolia</b> (Hook.f.) Nannf., Svensk Bot. Tidskr. 18: 418 (1924)                      | 2:247 |
| i                                  | <b>Conium maculatum</b> L., Sp. Pl. 243 (1753)   | 2:254 |
| i                                  | <b>Cyclospermum leptophyllum</b> (Pers.) Sprague, J. Bot. 61: 131 (1923)                             |       |
| i                                  | <b>Daucus carota</b> L., Sp. Pl. 242 (1753)  | 2:262 |
|                                    | <b>Daucus glochidiatus</b> (Labill.) Fischer et al., Ind. Sem. Petrop. 9, Suppl. 11 (1844)           | 2:262 |
|                                    | <b>Dichosciadium ranunculaceum</b> (F.Muell. ex Hook.) Domin, Feddes Repert. 5: 104 (1908)           | 2:250 |
| e                                  | var. <b>tasmanicum</b> (Hook.f.) Domin, Feddes Repert. 5: 105 (1908)                                 |       |
| e                                  | <b>Diplaspis cordifolia</b> (Hook.) Hook.f., Fl. Tasm. 1: 157 (1856)                                 | 2:252 |
| e                                  | <b>Diplaspis hydrocotyle</b> Hook.f., London J. Bot. 6: 469 bis (1847)                               | 2:251 |
|                                    | <b>Eryngium ovinum</b> A.Cunn. in Field, Geogr. Mem. N.S.W. 358 (1825)                               | 2:252 |
|                                    | Eryngium rostratum sensu W.M.Curtis (1963) = <b>Eryngium ovinum</b>                                  | 2:252 |
|                                    | <b>Eryngium vesiculosum</b> Labill., Nov. Holl. Pl. 1: 73 t.98 (1805)                                | 2:252 |
| i                                  | <b>Foeniculum vulgare</b> Miller, Gard. Dict. ed.8 no.1 (1768)                                       | 2:259 |
|                                    | <b>Hydrocotyle callicarpa</b> Bunge in Lehm., Pl. Preiss. 1: 283 (1845)                              | 2:246 |
|                                    | <b>Hydrocotyle capillaris</b> F.Muell. ex Klatt, Linnaea 29: 707 (1859)                              | 2:247 |
|                                    | <b>Hydrocotyle comocarpa</b> F.Muell., Vict. Nat. 3: 127 (1887)                                      |       |
|                                    | <b>Hydrocotyle foveolata</b> H.Eichler, Suppl. J.Black's Fl. S.Austral. ed.2: 248 (1965)             |       |
|                                    | <b>Hydrocotyle hirta</b> R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 204 (1820)                        | 2:245 |
|                                    | Hydrocotyle javanica sensu W.M.Curtis (1963) = <b>Hydrocotyle hirta</b>                              | 2:245 |
|                                    | <b>Hydrocotyle laxiflora</b> DC., Prodr. 4: 61 (1830)  | 2:246 |
|                                    | <b>Hydrocotyle muscosa</b> R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 208, t.61 fig.27 (1820)         | 2:246 |
|                                    | <b>Hydrocotyle pterocarpa</b> F.Muell., Trans. Proc. Vict. Inst. 126 (1855)                          | 2:246 |
|                                    | <b>Hydrocotyle sibthorpioides</b> Lamk., Encycl. 3: 153 (1789)                                       | 2:245 |
|                                    | <b>Hydrocotyle tripartita</b> R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 209, t.61 fig.25 (1820)      |       |
|                                    | Lilaeopsis brownii A.W.Hill = <b>Lilaeopsis polyantha</b>  | 2:260 |
|                                    | <b>Lilaeopsis polyantha</b> (Gand.) H.Eichler, Taxon 12: 296 (1963)                                  | 2:260 |
|                                    | <b>Oreomyrrhis argentea</b> (Hook.f.) Hook.f., Fl. Tasm. 1: 162 (1856)                               | 2:257 |

|                                    |  |       |
|------------------------------------|--|-------|
|                                    | <b>Oreomyrrhis ciliata</b> Hook.f., London J. Bot. 6: 471 bis (1847)                             | 2:258 |
|                                    | <b>Oreomyrrhis eriopoda</b> (DC.) Hook.f., Fl. Tasm. 1: 162 (1856)                               | 2:257 |
| e                                  | <b>Oreomyrrhis gunnii</b> Mathias & Constance, Univ. Calif. Publ. Bot. 27: 395 (1955)            | 2:258 |
| e                                  | <b>Oreomyrrhis sessiliflora</b> Hook.f., London J. Bot. 6: 471 bis (1847)                        | 2:258 |
| e                                  | <b>Oschatzia saxifraga</b> (Hook.f.) Walp., Ann. Bot. Syst. 1: 340 (1849)                        | 2:251 |
| i                                  | <b>Pastinaca sativa</b> L., Sp. Pl. 1: 262 (1753)  | 2:261 |
| i                                  | <b>Petroselinum crispum</b> (Miller) Nyman ex A.W.Hill, Hand-List Herb. Pl. Kew ed.3: 122 (1925) | 2:256 |
| i                                  | <b>Scandix pecten-veneris</b> L., Sp. Pl. 1: 256 (1753)  | 2:259 |
| i                                  | <b>Torilis nodosa</b> (L.) Gaertner, Fruet. Sem. Pl. 1: 82, t.20 (1788)                          | 2:262 |
|                                    | <b>Trachymene anisocarpa</b> (Turcz.) Burt. J. Bot. 79: 44 (1941)                                | 2:248 |
|                                    | <b>Trachymene humilis</b> (Hook.f.) Benth., Fl. Austral. 3: 351 (1867)                           | 2:248 |
|                                    | subsp. <b>humilis</b>  |       |
|                                    | subsp. <b>breviscapa</b> (Domin) P.S.Short, Muelleria 6: 166 (1986)                              |       |
|                                    | <b>Xanthosia dissecta</b> Hook.f., Icon. Pl. 4: t.302 (1840)                                     | 2:250 |
|                                    | <b>Xanthosia pilosa</b> Rudge, Trans. Linn. Soc. London 10: 301, t.22 fig.1 (1811)               | 2:249 |
|                                    | Xanthosia pusilla sensu W.M.Curtis (1963) = <b>Xanthosia tasmanica</b>                           | 2:250 |
|                                    | <b>Xanthosia tasmanica</b> Domin, Repert. Spec. Nov. Regni Veg. 4: 298 (1907)                    | 2:250 |
|                                    | <b>Xanthosia ternifolia</b> J.M.Hart & Henwood, Aust. Syst. Bot. 13(2): 256 (2000)               |       |
|                                    | <b>Xanthosia tridentata</b> DC., Prodr. 4: 75 (1830)   | 2:249 |
| <br><b>APOCYNACEAE</b>             |  |       |
|                                    | <b>Alyxia buxifolia</b> R.Br., Prodr. 470 (1810)   | 3:472 |
|                                    | <b>Parsonsia brownii</b> (Britten) Pichon, Notul. Syst. (Paris) 14: 10 (1950)                    | 3:473 |
|                                    | Parsonsia straminea sensu W.M.Curtis (1967) = <b>Parsonsia brownii</b>                           | 3:473 |
| i                                  | <b>Vinca major</b> L., Sp. Pl. 1: 209 (1753)   | 3:472 |
| <br><b>AQUIFOLIACEAE</b>           |  |       |
| i                                  | <b>Ilex aquifolium</b> L., Sp. Pl. 1: 125 (1753)   |       |
| <br><b>ARALIACEAE</b>              |  |       |
| i                                  | <b>Hedera helix</b> L., Sp. Pl. 1: 202 (1753)  |       |
|                                    | Nothopanax gunnii (Hook.f.) Seem. = <b>Pseudopanax gunnii</b>                                    | 2:264 |
| i                                  | <b>Polyscias sambucifolia</b> (Sieber ex DC.) Harms, Nat. Pflanzenfam. III 8: 47 (1898)          |       |
|                                    | <b>Polyscias</b> aff. <b>sambucifolia</b>  |       |
| e                                  | <b>Pseudopanax gunnii</b> (Hook.f.) K.Koch, Woch. Gärt. Pflanz. 2: 366 (1859)                    | 2:264 |
| <br><b>ASCLEPIADACEAE</b>          |  |       |
| i                                  | <b>Gomphocarpus fruticosus</b> (L.) W.T.Aiton, Hort. Kew edn.2, 2: 8 (1811)                      |       |
|                                    | subsp. <b>fruticosus</b>   |       |
| <br><b>ASTERACEAE (COMPOSITAE)</b> |  |       |
| e                                  | <b>Abrotanella forsteroides</b> (Hook.f.) Benth., Fl. Austral. 3: 554 (1867)                     | 2:359 |
| e                                  | <b>Abrotanella scapigera</b> (F.Muell.) Benth., Fl. Austral. 3: 554 (1867)                       | 2:359 |
| i                                  | <b>Achillea distans</b> Waldst. & Kit. ex Willd., Sp. Pl. 3: 2207 (1803)                         | 2:351 |

|    |  |       |
|----|--|-------|
| i  | <b>Achillea millefolium</b> L., Sp. Pl. 2: 899 (1753)  | 2:351 |
|    | <i>Achillea tanacetifolia</i> All. = <b>Achillea distans</b>   | 2:351 |
|    | <b>Actites megalocarpa</b> (Hook.f.) Lander, <i>Telopea</i> 1: 130 (1976)                                | 2:390 |
| i  | <b>Ammobium alatum</b> R.Br. in Sims, <i>Bot. Mag.</i> 51: t.2459 (1824)                                 | 2:343 |
| e  | <b>Ammobium calyceroides</b> (Cass.) A.Anderb., <i>Telopea</i> 4: 134 (1990)                             | 2:349 |
|    | <i>Angianthus eriocephalus</i> (A.Gray) Benth. = <b>Angianthus preissianus</b>                           | 2:344 |
|    | <b>Angianthus preissianus</b> (Steetz) Benth., <i>Fl. Austral.</i> 3: 566 (1867)                         | 2:344 |
| i  | <b>Anthemis arvensis</b> L., Sp. Pl. 2: 894 (1753)   | 2:352 |
| i  | <b>Anthemis cotula</b> L., Sp. Pl. 2: 894 (1753)   | 2:352 |
|    | <i>Anthemis nobilis</i> = <b>Chamaemelum nobile</b>  | 2:352 |
| i  | <b>Anthemis tinctoria</b> L., Sp. Pl. 2: 896 (1753)  | 2:352 |
|    | <b>Apalochlamys spectabilis</b> (Labill.) Steudel, <i>Nomen. Bot.</i> ed.2 1: 255 (1840)                 | 2:340 |
| i  | <b>Arctium minus</b> Bernh., <i>Syst. Verz. Pflanz.</i> 154 (1800)                                       | 2:375 |
| i  | <b>Arctotheca calendula</b> (L.) M.Levyns, <i>J. S.African Bot.</i> 8: 284 (1942)                        | 2:374 |
| i  | <b>Arctotheca populifolia</b> (P.Berguis) Norlindh in <i>Aquilo</i> , <i>Ser. Bot.</i> 6: 84 (1967)      |       |
| i  | <b>Arctotis stoechadifolia</b> P.Berguis, <i>Desc. Pl. Cap.</i> 324 (1767)                               | 2:374 |
|    | <b>Argentipallium dealbatum</b> (Labill.) Paul G.Wilson, <i>Nuytsia</i> 8: 458 (1992)                    | 2:330 |
|    | <b>Argentipallium obtusifolium</b> (Sonder) Paul G.Wilson, <i>Nuytsia</i> 8: 458 (1992)                  | 2:329 |
| e  | <b>Argentipallium spiceri</b> (F.Muell.) Paul G.Wilson, <i>Nuytsia</i> 8: 458 (1992)                     | 2:329 |
| n  | <b>Argyrotegium fordianum</b> (M.Gray) J.M.Ward & Breitw., <i>N.Z. J. Bot.</i> 41: 609 (2003)            |       |
| n  | <b>Argyrotegium nitidulum</b> (Hook.f.) J.M.Ward & Breitw., <i>N.Z. J. Bot.</i> 41: 609 (2003)           |       |
| n  | <b>Argyrotegium poliochlorum</b> (N.G.Walsh) J.M.Ward & Breitw., <i>N.Z. J. Bot.</i> 41: 609 (2003)      | 2:320 |
| i  | <b>Aster subulatus</b> Michaux, <i>Fl. Bor.-Amer.</i> 2: 111 (1803)                                      | 2:299 |
|    | <b>Bedfordia arborescens</b> Hochr., <i>Candollea</i> 5: 332 (1934)                                      |       |
| e  | <b>Bedfordia linearis</b> (Labill.) DC., <i>Prodr.</i> 6: 441 (1838)                                     | 2:371 |
| ne | subsp. <b>linearis</b>   |       |
| ne | subsp. <b>oblongifolia</b> Orch., <i>Muelleria</i> 19: 90 (2004)   |       |
| ne | subsp. <b>oblongifolia</b> var. <b>curvifolia</b> Orch., <i>Muelleria</i> 19: 93 (2004)                  |       |
| ne | subsp. <b>oblongifolia</b> var. <b>oblongifolia</b>  |       |
| e  | <b>Bedfordia salicina</b> (Labill.) DC., <i>Prodr.</i> 6: 441 (1838)                                     | 2:371 |
| i  | <b>Bellis perennis</b> L., Sp. Pl. 2: 886 (1753)   | 2:299 |
| i  | <b>Berkheya rigida</b> (Thunb.) Ewart, J.White & B.Rees, <i>Proc. R. Soc. Vict.</i> n.s.22: 20 (1909)    |       |
| e  | <b>Brachyglottis brunonis</b> (Hook.f.) B.Nord., <i>Opera Bot.</i> 44: 30 (1978)                         | 2:365 |
|    | <b>Brachyscome aculeata</b> (Labill.) Less., <i>Syn. Gen. Compos.</i> 192 (1832)                         | 2:296 |
|    | <b>Brachyscome angustifolia</b> A.Cunn. ex DC., <i>Prodr.</i> 5: 306 (1836)                              | 2:295 |
|    | var. <b>angustifolia</b>   | 2:295 |
|    | <b>Brachyscome cardiocarpa</b> F.Muell. ex Benth., <i>Fl. Austral.</i> 3: 517 (1867)                     | 2:296 |
|    | <i>Brachyscome ciliaris</i> (Labill.) Less., not present in Tasmania                                     | 2:298 |
|    | <b>Brachyscome decipiens</b> Hook.f., <i>London J. Bot.</i> 6: 114 (1847)                                | 2:295 |
|    | <b>Brachyscome diversifolia</b> (Graham ex Hook.) Fisch. & Mey., <i>Ind. Semin. Petrop.</i> 2: 31 (1835) | 2:297 |
|    | var. <b>diversifolia</b>   | 2:297 |
| e  | var. <b>maritima</b> Benth., <i>Fl. Austral.</i> 3: 511 (1867)   | 2:297 |
|    | <i>Brachyscome graminea</i> (Labill.) F.Muell. = <b>Paquerina graminea</b>                               | 2:295 |
|    | <i>Brachyscome nivalis</i> sensu W.M.Curtis (1963) = <b>B. radicans</b> sens. lat.                       | 2:297 |

|   |   |       |
|---|---|-------|
|   | <b>Brachyscome parvula</b> Hook.f., Fl. Tasm. 1: 185 (1856)                             | 2:298 |
|   | var. <b>parvula</b>   | 2:298 |
|   | <b>Brachyscome perpusilla</b> (Steetz) J.M.Black, Fl. S.Austral. 4: 587 (1929)          |       |
|   | <b>Brachyscome radicans</b> Steetz in Lehm., Pl. Preiss. 1: 429 (1845)                  | 2:295 |
| t | <b>Brachyscome radicata</b> Hook.f., Fl. Nov.-Zel. 1: 127 (1852)                        | 2:298 |
|   | <b>Brachyscome rigidula</b> (DC.) G.Davis, Proc. Linn. Soc. N.S.W. 73: 219 (1948)       | 2:298 |
|   | <i>Brachyscome scapiformis</i> DC. = <b>Brachyscome spathulata</b> subsp. <b>glabra</b> | 2:296 |
|   | <b>Brachyscome sieberi</b> DC., Prodr. 5: 306 (1836)                                    |       |
| e | var. <b>gunnii</b> DC., Prodr. 5: 306 (1836)  |       |
|   | <b>Brachyscome spathulata</b> Gaud. in Freycinet, Voy. Uranie 468 (1830)                |       |
|   | subsp. <b>glabra</b> (DC.) Stace, Austral. J. Bot. 29: 436 (1981)                       | 2:296 |
|   | subsp. <b>spathulata</b>  |       |
|   | <i>Brachyscome stricta</i> DC. = <b>Brachyscome aculeata</b>                            | 2:296 |
|   | <i>Brachyscome tadgellii</i> Tovey & P.Morris, not present in Tasmania                  |       |
|   | <b>Brachyscome tenuiscapa</b> Hook.f., London J. Bot. 6: 114 (1847)                     | 2:294 |
|   | var. <i>pubescens</i> (Benth.) G.Davis, not present in Tasmania                         | 2:294 |
|   | var. <b>tenuiscapa</b>  | 2:294 |
|   | <i>Bracteantha bicolor</i> = <b>Xerochrysum bicolor</b>                                 | 2:331 |
|   | <i>Bracteantha bracteata</i> = <b>Xerochrysum bracteatum</b>                            |       |
|   | <i>Bracteantha palustris</i> = <b>Xerochrysum palustre</b>                              |       |
|   | <i>Bracteantha papillosa</i> = <b>Xerochrysum papillosum</b>                            | 2:331 |
|   | <i>Bracteantha subundulata</i> = <b>Xerochrysum subundulatum</b>                        | 2:330 |
| i | <b>Calendula arvensis</b> L., Sp. Pl. ed.2, 2: 1303 (1763)                              | 2:372 |
| i | <b>Calendula officinalis</b> L., Sp. Pl. 2: 921 (1753)                                  | 2:373 |
|   | <i>Calocephalus brownii</i> (Cass.) F.Muell. = <b>Leucophyta brownii</b>                | 2:345 |
|   | <b>Calocephalus citreus</b> Less., Syn. Gen. Compos. 271 (1832)                         | 2:346 |
|   | <b>Calocephalus lacteus</b> Less., Syn. Gen. Compos. 271 (1832)                         | 2:345 |
| i | <b>Carduus nutans</b> L., Sp. Pl. 2: 821 (1753)   |       |
| i | <b>Carduus pycnocephalus</b> L., Sp. Pl. ed.2, 2: 1151 (1763)                           |       |
| i | <b>Carduus tenuiflorus</b> Curtis, Fl. Lond. 2(6): t.55 (1789)                          | 2:376 |
| i | <b>Carthamus lanatus</b> L., Sp. Pl. 2: 830 (1753)                                      | 2:380 |
| i | <b>Carthamus tinctorius</b> L., Sp. Pl. 2: 830 (1753)                                   |       |
|   | <b>Cassinia aculeata</b> (Labill.) R.Br., Trans. Linn. Soc. London 12: 127 (1817)       | 2:339 |
|   | <i>Cassinia longifolia</i> sensu W.M.Curtis (1963) = <b>Cassinia trinerva</b>           | 2:340 |
|   | <i>Cassinia spectabilis</i> (Labill.) R.Br. = <b>Apalochlamys spectabilis</b>           | 2:340 |
|   | <b>Cassinia trinerva</b> N.A.Wakef., Vict. Nat. 68: 69 (1951)                           | 2:340 |
| e | <b>Celmisia asteliifolia</b> Hook.f., Fl. Antarct. 1: 35 (1844)                         | 2:310 |
|   | <i>Celmisia longifolia</i> Cass. = <b>Celmisia asteliifolia</b>                         | 2:310 |
| e | <b>Celmisia saxifraga</b> (Benth.) W.M.Curtis, Taxon 17: 467 (1968)                     | 2:310 |
| i | <b>Centaurea calcitrapa</b> L., Sp. Pl. 2: 917 (1753)                                   | 2:379 |
| i | <b>Centaurea jacea</b> L., Sp. Pl. 2: 914 (1753)  | 2:378 |
| i | <b>Centaurea melitensis</b> L., Sp. Pl. 2: 917 (1753)                                   | 2:380 |
| i | <b>Centaurea nigra</b> L., Sp. Pl. 2: 911 (1753)  | 2:379 |
| i | <b>Centaurea solstitialis</b> L., Sp. Pl. 2: 917 (1753)                                 | 2:379 |

|    |  |       |
|----|--|-------|
|    | <b>Centipeda cunninghamii</b> (DC.) A.Braun & Asch., Ind. Sem. Hort. Berol. App.6 (1867)               |       |
|    | <b>Centipeda elatinoides</b> (Less.) Benth. & Hook.f. ex O.Hoffm., Nat. Pflanzenfam. 4(5): 280 (1892)  | 2:358 |
|    | Centipeda minima sensu W.M.Curtis (1963) = <b>Centipeda elatinoides</b>                                | 2:358 |
| i  | <b>Chamaemelum nobile</b> (L.) All., Fl. Ped. 1: 185 (1785)  | 2:352 |
| i  | <b>Chondrilla juncea</b> L., Sp. Pl. 2: 796 (1753)   | 2:388 |
| i  | <b>Chrysanthemoides monilifera</b> (L.) Norlindh, Stud. Calend. 1: 374 (1943) subsp. <b>monilifera</b> | 2:373 |
|    | Chrysanthemum leucanthemum L. = <b>Leucanthemum vulgare</b>  | 2:353 |
|    | Chrysanthemum parthenium (L.) Bernh. = <b>Tanacetum parthenium</b>                                     | 2:353 |
|    | <b>Chrysocephalum apiculatum</b> (Labill.) Steetz in Lehm., Pl. Preiss. 1: 474 (1845)                  | 2:332 |
|    | <b>Chrysocephalum baxteri</b> (A.Cunn. ex DC.) A.Anderb., Comp. Newsl. 19: 22 (1991)                   | 2:328 |
|    | <b>Chrysocephalum semipapposum</b> (Labill.) Steetz in Lehm., Pl. Preiss. 1: 474 (1845)                | 2:332 |
| i  | <b>Cichorium intybus</b> L., Sp. Pl. 2: 813 (1753)   | 2:380 |
| i  | <b>Cirsium arvense</b> (L.) Scop., Fl. Carniol. ed.2, 2: 126 (1772)                                    | 2:377 |
| i  | <b>Cirsium vulgare</b> (Savi) Ten., Fl. Napol. 5: 209 (1835-6)   | 2:376 |
|    | Cnicus arvensis (L.) Hoffm. = <b>Cirsium arvense</b>   | 2:377 |
| i  | <b>Conyza albida</b> Willd. ex Sprengel, Syst. Veg. 3: 514 (1826)                                      | 2:315 |
| i  | <b>Conyza bonariensis</b> (L.) Cronq., Bull. Torrey Bot. Club 70: 632 (1943)                           | 2:315 |
| i  | <b>Conyza canadensis</b> (L.) Cronq., Bull. Torrey Bot. Club 70: 632 (1943)                            | 2:314 |
| ix | <b>Conyza chilensis</b> Spreng., Nov. Prov. 14 (1818)  | 2:314 |
|    | Conyza floribunda sensu W.M.Curtis (1963) = <b>Conyza albida</b>                                       | 2:315 |
|    | Conyza scabiosifolia E.J.Remy = <b>Conyza chilensis</b>  | 2:314 |
|    | <b>Cotula alpina</b> (Hook.f.) Hook.f., Fl. Tasm. 1: 192 t.51A (1856)                                  | 2:356 |
|    | <b>Cotula australis</b> (Sieber ex Sprengel) Hook.f., Fl. Nov.-Zel. 1: 128 (1853)                      | 2:356 |
| i  | <b>Cotula coronopifolia</b> L., Sp. Pl. 2: 892 (1753)  | 2:356 |
|    | Cotula filicula (Hook.f.) Benth. = <b>Leptinella filicula</b>  | 2:357 |
|    | Cotula longipes (Hook.f.) W.M.Curtis = <b>Leptinella longipes</b>                                      | 2:357 |
|    | Cotula reptans (Benth.) Benth. = <b>Leptinella reptans</b>   | 2:357 |
|    | <b>Cotula vulgaris</b> M.R.Levyns, J. S.Afr. Bot. 7: 133 (1941)  | 2:355 |
|    | var. <b>australasica</b> J.H.Willis, Vict. Nat. 73: 201 (1957)   | 2:355 |
|    | <b>Craspedia alpina</b> Backh. ex Hook.f., London J. Bot. 6: 119 (1847)                                | 2:347 |
|    | <b>Craspedia coolaminica</b> J.Everett & Joy Thomps., Telopea 5: 45 (1992)                             |       |
| e  | <b>Craspedia glabrata</b> (Hook.f.) Rozefelds, Telopea 9(4): 814 (2002)                                | 2:347 |
|    | <b>Craspedia glauca</b> (Labill.) Sprengel, Syst. Veg. 3: 441 (1826)                                   | 2:347 |
|    | var. <b>glabrata</b> (Hook.f.) W.M.Curtis (comb. inval.) = <b>Craspedia glabrata</b>                   | 2:347 |
|    | var. <b>gracilis</b> nom. inval.   | 2:347 |
|    | var. <b>macrocephala</b> nom. inval.   | 2:347 |
|    | <b>Craspedia paludicola</b> J.Everett & Doust, Telopea 5: 35 (1992)                                    |       |
| e  | <b>Craspedia preminghana</b> Rozefelds, Telopea 9(4): 816 (2002)                                       |       |
| i  | <b>Crepis capillaris</b> (L.) Wallr., Erst. Beitr. Fl. Hercyn. 287 (1840)                              | 2:384 |
|    | Crepis nicaeensis Balb., not present in Tasmania, formerly recorded in error                           | 2:384 |
| i  | <b>Crepis setosa</b> Haller.f. in Roemer, Arch. Bot. 1: 1 (1796-97)                                    | 2:384 |
|    | Cymbonotus lawsonianus sensu W.M.Curtis (1963) = <b>Cymbonotus preissianus</b>                         | 2:373 |
|    | <b>Cymbonotus preissianus</b> Steetz in Lehm., Pl. Preiss 1: 486 (1845)                                | 2:373 |
| i  | <b>Delairea odorata</b> Lemaire, Ann. Sci. Nat. Bot. ser.3, 1: 380 (1844)                              | 2:367 |

|   |   |       |
|---|---|-------|
| i | <b>Dittrichia graveolens</b> (L.) Greuter, Exsicc. Genev. 4: 71 (1973)                      | 2:348 |
|   | <b>Erigeron bellidioides</b> (Hook.f.) S.J.Forbes & D.I.Morris, Muelleria 9: 183 (1996)     |       |
| e | <b>Erigeron gunnii</b> (Hook.f.) F.Muell. in Hook.f., Fl. Tasm. 1: 183, t.46b (1856)        | 2:312 |
|   | var. bellidioides (Hook.f.) Hook.f. = <b>Erigeron bellidioides</b>                          |       |
| i | <b>Erigeron karvinskianus</b> DC., Prodr. 5: 285 (1836)                                     | 2:312 |
|   | <i>Erigeron mucronatus</i> DC. = <b>Erigeron karvinskianus</b>                              | 2:312 |
| e | <b>Erigeron pappocromus</b> Labill., Nov. Holl. Pl. 2: 47 t.193 (1806)                      | 2:311 |
|   | var. billardieri Benth. = <b>Erigeron pappocromus</b>                                       | 2:311 |
|   | var. gunnii Benth. = <b>Erigeron gunnii</b>   | 2:312 |
|   | var. oblongatus Benth. = <b>Erigeron tasmanicus</b>   |       |
| e | <b>Erigeron stellatus</b> (Hook.f.) W.M.Curtis, Stud. Fl. Tasm. 2: 463 (1963)               | 2:312 |
|   | <b>Erigeron tasmanicus</b> (Hook.f.) Hook.f., Fl. Tasm. 1: 183, t.46A (1856)                |       |
| e | <b>Erigeron trigonus</b> S.J.Forbes & D.I.Morris, Muelleria 9: 187 (1996)                   |       |
|   | <b>Euchiton argentifolius</b> (N.A.Wakef.) A.Anderb., Opera Bot. 104: 167 (1991)            | 2:320 |
|   | <b>Euchiton collinus</b> Cass., Dict. Sci. Nat. 56: 216 (1828)                              | 2:319 |
|   | <i>Euchiton fordianus</i> (M.Gray) P.S.Short = <b>Argyrotegium fordianum</b>                |       |
|   | <b>Euchiton involucratus</b> (Forst.f.) Holub, Fol. Geobot. Phytotax. 9: 271 (1974)         | 2:318 |
| e | <b>Euchiton litticola</b> A.M.Buchanan, Pap. Proc. Roy. Soc. Tasm. 133(1): 115 (1999)       |       |
|   | <i>Euchiton poliochlorus</i> N.G.Walsh = <b>Argyrotegium poliochlorum</b>                   | 2:320 |
|   | <b>Euchiton sphaericus</b> (Willd.) Holub, Fol. Geobot. Phytotax. 9: 271 (1974)             |       |
|   | <b>Euchiton traversii</b> (Hook.f.) Holub, Fol. Geobot. Phytotax. 9: 271 (1974)             | 2:320 |
|   | <b>Euchiton umbricola</b> (J.H.Willis) A.Anderb., Opera Bot. 104: 167 (1991)                | 2:319 |
| i | <b>Euryops abrotanifolius</b> (L.) DC., Prodr. 6: 443 (1838)                                | 2:372 |
| e | <b>Ewartia catipes</b> (DC.) P.Beauv., Bull. Soc. Bot. Gèneve ser.2, 2: 238 (1910)          | 2:322 |
| e | <b>Ewartia meredithiae</b> (F.Muell.) P.Beauv., Bull. Soc. Bot. Gèneve ser.2, 2: 240 (1910) | 2:322 |
| e | <b>Ewartia planchonii</b> (Hook.f.) P.Beauv., Bull. Soc. Bot. Gèneve ser.2, 3: 253 (1911)   | 2:321 |
|   | <i>Felicia erigeroides</i> sensu T.D.Raphael (1955) = <b>Erigeron karvinskianus</b>         |       |
|   | <i>Filago gallica</i> L. = <b>Logfia gallica</b>  | 2:315 |
| i | <b>Galinsoga parviflora</b> Cav., Icon. 3: 41 t.281 (1795)                                  | 2:349 |
| i | <b>Gamochaeta calviceps</b> (Fernald) Cabrera, Bol. Soc. Argent. Bot. 9: 368 (1961)         |       |
| i | <b>Gamochaeta purpurea</b> (L.) Cabrera, Bol. Soc. Argent. Bot. 9: 377 (1961)               | 2:321 |
|   | <i>Gnaphalium argentifolium</i> N.A.Wakef. = <b>Euchiton argentifolius</b>                  | 2:320 |
|   | <i>Gnaphalium candidissimum</i> Lam. = <b>Vellereophyton dealbatum</b>                      | 2:318 |
|   | <i>Gnaphalium collinum</i> Labill.  |       |
|   | var. collinum = <b>Euchiton collinus</b>  | 2:319 |
|   | var. monocephalum Hook.f. = <b>Euchiton argentifolius</b>                                   | 2:319 |
|   | <i>Gnaphalium fordianum</i> M.Gray = <b>Argyrotegium fordianum</b>                          |       |
|   | <i>Gnaphalium indicum</i> auct. non L. = <b>Gnaphalium polycaulon</b>                       |       |
|   | <b>Gnaphalium indutum</b> Hook.f., London J. Bot. 6: 121 (1847)                             | 2:320 |
|   | <i>Gnaphalium involucratum</i> Forst.f. = <b>Euchiton involucratus</b>                      | 2:318 |
|   | <i>Gnaphalium luteo-album</i> L. = <b>Pseudognaphalium luteoalbum</b>                       | 2:318 |
|   | <b>Gnaphalium polycaulon</b> Pers., Syn. Pl. 2: 421 (1807)                                  |       |
|   | <i>Gnaphalium purpureum</i> L. = <b>Gamochaeta purpurea</b>                                 | 2:321 |
|   | <i>Gnaphalium supinum</i> sensu W.M.Curtis (1963) = <b>Argyrotegium poliochlorum</b>        | 2:320 |

|   |  |       |
|---|--|-------|
|   | Gnaphalium traversii Hook.f. = <b>Euchiton traversii</b>   | 2:320 |
|   | Gnaphalium umbricola J.H.Willis = <b>Euchiton umbricola</b>                                      | 2:319 |
|   | Hedypnois cretica (L.) Dum.-Cours. = <b>Hedypnois rhagodiolooides</b>                            | 2:382 |
| i | <b>Hedypnois rhagodiolooides</b> (L.) F.W.Schmidt, Samml. Phys.-Oekon. Aufsätze 1: 279 (1795)    | 2:382 |
|   | Helichrysum acuminatum DC. = <b>Xerochrysum subundulatum</b>                                     | 2:330 |
|   | Helichrysum adenophorum F.Muell., not present in Tasmania, formerly recorded in error            |       |
|   | Helichrysum antennarium (DC.) F.Muell. ex Benth. = <b>Ozothamnus antennaria</b>                  | 2:336 |
|   | Helichrysum apiculatum (Labill.) D.Don = <b>Chrysocephalum apiculatum</b>                        | 2:332 |
|   | Helichrysum argophyllum (A.Cunn. ex DC.) N.A.Wakef. = <b>Ozothamnus argophyllum</b>              | 2:335 |
|   | Helichrysum backhousii (Hook.f.) F.Muell. ex Benth. = <b>Ozothamnus rodwayi</b>                  | 2:337 |
|   | var. backhousii = <b>Ozothamnus rodwayi</b> var. <b>rodwayi</b>                                  |       |
|   | var. kingii W.M.Curtis in Morris & Curtis = <b>Ozothamnus rodwayi</b> var. <b>kingii</b>         |       |
|   | var. oreophilum W.M.Curtis in Morris & Curtis = <b>Ozothamnus rodwayi</b> var. <b>oreophilus</b> |       |
|   | Helichrysum baxteri A.Cunn. ex DC. = <b>Chrysocephalum baxteri</b>                               | 2:328 |
|   | Helichrysum bicolor Lindley = <b>Xerochrysum bicolor</b>   | 2:331 |
|   | Helichrysum bracteatum (Vent.) Andrews = <b>Xerochrysum bracteatum</b>                           |       |
|   | Helichrysum costatifractum R.V.Smith = <b>Ozothamnus costatifractus</b>                          | 2:332 |
|   | Helichrysum dealbatum Labill. = <b>Argentipallium dealbatum</b>                                  | 2:330 |
|   | Helichrysum dendroideum N.A.Wakef. = <b>Ozothamnus ferrugineus</b>                               | 2:335 |
|   | Helichrysum ericeteum W.M.Curtis = <b>Ozothamnus ericifolius</b>                                 | 2:334 |
|   | Helichrysum expansifolium (P.Morris & J.H.Willis) N.Burb. = <b>Ozothamnus expansifolius</b>      | 2:337 |
|   | Helichrysum gunnii (Hook.f.) Benth. = <b>Ozothamnus bracteolatus</b>                             | 2:333 |
|   | Helichrysum hookeri (Sonder) Druce = <b>Ozothamnus hookeri</b>                                   | 2:337 |
|   | Helichrysum ledifolium (DC.) Benth. = <b>Ozothamnus ledifolium</b>                               | 2:334 |
|   | <b>Helichrysum leucopsidium</b> DC., Prodr. 6: 193 (1838)  | 2:331 |
|   | Helichrysum lycopodioides (Hook.f.) Benth. = <b>Ozothamnus lycopodioides</b>                     | 2:338 |
| e | <b>Helichrysum milliganii</b> Hook.f., Fl. Tasm. 1: 214, t.60 (1856)                             | 2:330 |
|   | Helichrysum obcordatum (DC.) Benth. = <b>Ozothamnus obcordatus</b>                               | 2:336 |
|   | Helichrysum obtusifolium F.Muell. & Sonder ex Sonder = <b>Argentipallium obtusifolium</b>        | 2:329 |
|   | Helichrysum papillosum Labill. = <b>Xerochrysum papillosum</b>                                   | 2:331 |
|   | Helichrysum parali (N.Burb.) W.M.Curtis = <b>Ozothamnus turbinatus</b>                           | 2:333 |
|   | Helichrysum pleurandroides sensu Census Vasc. Pl. Tasm. edn.1 = <b>Ozothamnus ericifolius</b>    | 2:334 |
| e | <b>Helichrysum pumilum</b> Hook.f., Fl. Tasm. 1: 213 (1856)                                      | 2:330 |
| e | var. <b>pumilum</b>  |       |
| e | var. <b>spathulatum</b> A.M.Buchanan, Pap. Proc. R. Soc. Tasm. 121: 54 (1987)                    |       |
|   | Helichrysum purpurascens (DC.) W.M.Curtis = <b>Ozothamnus purpurascens</b>                       | 2:334 |
|   | Helichrysum reticulatum (Labill.) Less. = <b>Ozothamnus reticulatus</b>                          | 2:332 |
|   | Helichrysum rosmarinifolium (Labill.) Benth. = <b>Ozothamnus rosmarinifolium</b>                 | 2:335 |
|   | <b>Helichrysum rutidolepis</b> DC., Prodr. 6: 194 (1838)   |       |
|   | <b>Helichrysum scorpioides</b> Labill., Nov. Holl. Pl. 2: 45 t.191 (1806)                        | 2:328 |
|   | Helichrysum scutellifolium (Hook.f.) F.Muell. = <b>Ozothamnus scutellifolius</b>                 | 2:338 |
|   | Helichrysum selaginoides (Sonder & F.Muell.) F.Muell. ex Benth. = <b>Ozothamnus selaginoides</b> | 2:339 |
|   | Helichrysum semipapposum (Labill.) DC. = <b>Chrysocephalum semipapposum</b>                      | 2:332 |
|   | Helichrysum spiceri F.Muell. = <b>Argentipallium spiceri</b>                                     | 2:329 |



|    |  |       |
|----|--|-------|
|    | Helichrysum thyrsoideum (DC.) P.Morris & J.H.Willis = <b>Ozothamnus thyrsoideus</b>              | 2:335 |
|    | Helipterum albicans (A.Cunn.) DC. = <b>Leucochrysum albicans</b>                                 | 2:324 |
|    | var. <i>incanum</i> (Hook.) Paul G.Wilson = <b>Leucochrysum albicans</b> var. <b>tricolor</b>    | 2:325 |
|    | Helipterum anthemoides (Sieber ex Sprengel) DC. = <b>Rhodanthe anthemoides</b>                   | 2:324 |
|    | Helipterum australe (A.Gray) Druce = <b>Triptilodiscus pygmaeus</b>                              |       |
|    | Helipterum demissum (A.Gray) Druce = <b>Hyalosperma demissum</b>                                 | 2:325 |
| i  | <b>Helminthotheca echioides</b> (L.) Holub, Fol. Geobot. Phytotax. 8: 176 (1973)                 | 2:383 |
| it | <b>Hieracium aurantiacum</b> L., Sp. Pl. 2: 801 (1753)   | 2:385 |
|    | Hieracium brunneocroceum Pugls. = <b>Hieracium aurantiacum</b>                                   | 2:385 |
| it | <b>Hieracium pilosella</b> L., Sp. Pl. 2: 800 (1753)   |       |
|    | <b>Hyalosperma demissum</b> (A.Gray) Paul G.Wilson, Nuytsia 7: 85 (1989)                         | 2:325 |
| i  | <b>Hypochoeris glabra</b> L., Sp. Pl. 2: 811 (1753)  | 2:386 |
| i  | <b>Hypochoeris radicata</b> L., Sp. Pl. 2: 811 (1753)  | 2:385 |
|    | Inula graveolens (L.) Desf. = <b>Dittrichia graveolens</b>                                       | 2:348 |
|    | <b>Isoetopsis graminifolia</b> Turcz., Bull. Soc. Nat. Musc. 24: 175 t.3 (1853)                  | 2:360 |
|    | Ixiolaena supina F.Muell. = <b>Leiocarpa supina</b>  | 2:341 |
|    | Ixodia achlaena D.I.Morris = <b>Odixia achlaena</b>  |       |
|    | Ixodia angusta (N.A.Wakef.) N.Burb. = <b>Odixia angusta</b>                                      | 2:343 |
| i  | <b>Lactuca saligna</b> L., Sp. Pl. 2: 796 (1753)   | 2:388 |
| i  | <b>Lactuca serriola</b> L., Cent. Pl. 2: 29 (1756)   | 2:389 |
|    | <b>Lagenophora gracilis</b> Steetz in Lehm., Pl. Preiss. 1: 430 (1845)                           |       |
|    | <b>Lagenophora huegelii</b> Benth. in Endl., Enum. Pl. Hügel 59 (1837)                           | 2:290 |
|    | <b>Lagenophora montana</b> Hook.f., Hooker's Lond. J. Bot. 6: 113 (1847)                         |       |
|    | <b>Lagenophora stipitata</b> (Labill.) Druce, Bot. Exch. Club Soc. Brit. Isles 1916: 630 (1917)  | 2:290 |
|    | Lagenifera, see <b>Lagenophora</b> (nom. cons.)  |       |
| i  | <b>Lapsana communis</b> L., Sp. Pl. 2: 811 (1753)  | 2:382 |
| it | <b>Lasiospermum bipinnatum</b> (Thunb.) Druce, Bot. Exch. Club Soc. Brit. Isles 1916: 631 (1917) | 2:350 |
|    | <b>Leiocarpa supina</b> (F.Muell.) Paul G.Wilson, Nuytsia 13(3): 604 (2001)                      | 2:341 |
|    | Leontodon leysseri (Wallr.) Beck = <b>Leontodon taraxacoides</b>                                 | 2:386 |
| i  | <b>Leontodon taraxacoides</b> (Vill.) Mérat, Ann. Sci. Nat. 22: 108 (1831)                       | 2:386 |
|    | <b>Leptinella filicula</b> (Hook.f.) Hook.f., Fl. Tasm. 1: 194 (1856)                            | 2:357 |
|    | <b>Leptinella longipes</b> Hook.f. in Hook., Lond. J. Bot. 6: 117 (1847)                         | 2:357 |
|    | <b>Leptinella reptans</b> (Benth.) D.Lloyd & C.Webb, N.Z. J. Bot. 25: 103 (1987)                 | 2:357 |
|    | <b>Leptorhynchos elongatus</b> DC., Prodr. 6: 160 (1838)   | 2:323 |
|    | Leptorhynchos linearis sensu W.M.Curtis (1963) = <b>Leptorhynchos nitidulus</b>                  | 2:323 |
|    | <b>Leptorhynchos nitidulus</b> DC., Prodr. 6: 160 (1838)   | 2:323 |
|    | <b>Leptorhynchos squamatus</b> (Labill.) Less., Syn. Gen. Compos. 273 (1832)                     | 2:323 |
|    | subsp. <i>alpinus</i> Flann, Aust. Syst. Bot. 15(2): 217 (2002)                                  |       |
|    | subsp. <i>squamatus</i>  |       |
| i  | <b>Leucanthemum vulgare</b> Lam., Fl. Franc. 2: 137 (1779)                                       | 2:353 |
|    | <b>Leucochrysum albicans</b> (A.Cunn.) Paul G.Wilson, Nuytsia 8: 442 (1992)                      | 2:324 |
|    | subsp. <i>albicans</i> var. <b>tricolor</b> (DC.) Paul G.Wilson, Nuytsia 8: 443 (1992)           | 2:325 |
|    | <b>Leucophyta brownii</b> Cass. in Cuvier, Dict. Sci. Nat. 26: 159 (1823)                        | 2:345 |
| i  | <b>Logfia gallica</b> (L.) Cosson & Germ., Ann. Sci. Nat. ser.2, 20: 291 (1843)                  | 2:315 |

|    |   |       |
|----|---|-------|
|    | Matricaria discoidea DC. = <b>Matricaria matricarioides</b>   | 2:354 |
| i  | <b>Matricaria matricarioides</b> (Less.) Porter, Mem. Torrey Bot. Club 5: 341 (1894)                | 2:354 |
|    | Matricaria perforata sensu Census (1995) = <b>Tripleurospermum maritimum</b> subsp. <b>inodorum</b> |       |
|    | <b>Microseris lanceolata</b> (Walp.) Schultz-Bip., Jahresber. Pollichia 22-24: 310 (1866)           | 2:381 |
|    | Microseris scapigera sensu W.M.Curtis (1963) = <b>Microseris lanceolata</b>                         | 2:381 |
|    | <b>Millotia muelleri</b> (Sonder) P.S.Short, Muelleria 7: 246 (1990)                                |       |
|    | <b>Millotia tenuifolia</b> Cass., Ann. Sci. Nat. 17: 31, 417 (1829)                                 | 2:342 |
|    | var. <b>tenuifolia</b>  |       |
|    | Nablonium calyceroides Cass. = <b>Ammobium calyceroides</b>   | 2:349 |
| e  | <b>Odixia achlaena</b> (D.I.Morris) Orch., Brunonia 4: 194 (1981)                                   |       |
| e  | <b>Odixia angusta</b> (N.A.Wakef.) Orch., Brunonia 4: 194 (1981)                                    | 2:343 |
|    | <b>Olearia algida</b> N.A.Wakef., Vict. Nat. 73: 97 (1956)  | 2:308 |
|    | Olearia alpina (Hook.f.) W.M.Curtis = <b>Olearia tasmanica</b>                                      | 2:304 |
| e  | <b>Olearia archeri</b> Lander, Muelleria 7: 117 (1989)  |       |
|    | <b>Olearia argophylla</b> (Labill.) Benth., Fl. Austral. 3: 470 (1867)                              | 2:303 |
|    | <b>Olearia axillaris</b> (DC.) F.Muell. ex Benth., Fl. Austral. 3: 475 (1867)                       | 2:307 |
|    | <b>Olearia ciliata</b> (Benth.) F.Muell. ex Benth., Fl. Austral. 3: 488 (1867)                      | 2:309 |
| e  | <b>Olearia ericoides</b> (Steetz) N.A.Wakef., Vict. Nat. 73: 97 (1956)                              | 2:307 |
|    | <b>Olearia erubescens</b> (DC.) Dippel, Handb. Laubholz 1: 290 (1889)                               | 2:303 |
|    | <b>Olearia floribunda</b> (Hook.f.) Benth., Fl. Austral. 3: 477 (1867)                              | 2:308 |
|    | <b>Olearia glandulosa</b> (Labill.) Benth., Fl. Austral. 3: 483 (1867)                              | 2:309 |
|    | <b>Olearia glutinosa</b> (Lindley) Benth., Fl. Austral. 3: 482 (1867)                               | 2:309 |
| e  | <b>Olearia hookeri</b> (Sonder) Benth., Fl. Austral. 3: 483 (1867)                                  | 2:309 |
|    | Olearia lanceolata (Benth.) D.I.Morris = <b>Olearia archeri</b>                                     |       |
| e  | <b>Olearia ledifolia</b> (DC.) Benth., Fl. Austral. 3: 477 (1867)                                   | 2:305 |
|    | <b>Olearia lepidophylla</b> (Pers.) Benth., Fl. Austral. 3: 477 (1867)                              | 2:308 |
|    | <b>Olearia lirata</b> (Sims) Hutch., Gard. Chron. ser.3, 61: 14 fig.4 (1917)                        | 2:306 |
|    | <b>Olearia myrsinoides</b> (Labill.) F.Muell. ex Benth., Fl. Austral. 3: 470 (1867)                 | 2:303 |
| e  | <b>Olearia obcordata</b> (Hook.f.) Benth., Fl. Austral. 3: 471 (1867)                               | 2:304 |
| e  | <b>Olearia peroonioides</b> (DC.) Benth., Fl. Austral. 3: 471 (1867)                                | 2:304 |
|    | <b>Olearia phlogopappa</b> (Labill.) DC., Prodr. 5: 272 (1836)                                      | 2:305 |
| e  | var. <b>angustifolia</b> (Hutch.) W.M.Curtis, Stud. Fl. Tasm. 2: 463 (1963)                         | 2:306 |
| e  | var. <b>brevipes</b> (Hutch.) W.M.Curtis, Stud. Fl. Tasm. 2: 463 (1963)                             | 2:306 |
| e  | var. <b>microcephala</b> (Hutch.) W.M.Curtis, Stud. Fl. Tasm. 2: 463 (1963)                         | 2:306 |
|    | var. <b>phlogopappa</b>   | 2:306 |
| e  | var. <b>salicifolia</b> (Hutch.) W.M.Curtis, Stud. Fl. Tasm. 2: 463 (1963)                          | 2:306 |
|    | var. <b>subrepanda</b> (DC.) J.H.Willis, Muelleria 1: 32 (1956)                                     | 2:306 |
| e  | <b>Olearia pinifolia</b> (Hook.f.) Benth., Fl. Austral. 3: 471 (1867)                               | 2:305 |
|    | <b>Olearia ramulosa</b> (Labill.) Benth., Fl. Austral. 3: 476 (1867)                                | 2:307 |
|    | <b>Olearia stellulata</b> (Labill.) DC., Prodr. 5: 272 (1806)                                       | 2:305 |
| e  | <b>Olearia tasmanica</b> (Hook.f.) W.M.Curtis, Vict. Nat. 87: 270 (1970)                            | 2:304 |
|    | <b>Olearia viscosa</b> (Labill.) Benth., Fl. Austral. 3: 468 (1867)                                 | 2:302 |
| i  | <b>Onopordum acanthium</b> L., Sp. Pl. 2: 827 (1753)  | 2:377 |
| i# | <b>Onopordum acaulon</b> L., Sp. Pl. ed.2, 2: 1159 (1763)   |       |

|    |  |       |
|----|--|-------|
| i  | <b>Osteospermum fruticosum</b> (L.) Norl., Stud. Calend. 1: 250 (1943)                         |       |
| e  | <b>Ozothamnus antennaria</b> (DC.) Hook.f., Fl. Tasm. 1: 203 (1856)                            | 2:336 |
|    | <b>Ozothamnus argophyllus</b> (A.Cunn. ex DC.) A.Anderb., Opera Bot. 104: 89 (1991)            | 2:335 |
| e  | <b>Ozothamnus bracteolatus</b> Hook.f., Fl. Tasm. 1(3): 203 (1856)                             | 2:333 |
|    | <i>Ozothamnus cinereus</i> sensu Rodway (1903) = <b>Ozothamnus turbinatus</b>                  | 2:333 |
| e  | <b>Ozothamnus costatifructus</b> (R.V.Smith) A.Anderb., Opera Bot. 104: 89 (1991)              | 2:332 |
| e  | <b>Ozothamnus ericifolius</b> Hook.f., Lond. J. Bot. 6: 119 (1847)                             | 2:334 |
| e  | <b>Ozothamnus expansifolius</b> (P.Morris & J.H.Willis) A.Anderb., Opera Bot. 104: 89 (1991)   | 2:337 |
|    | <b>Ozothamnus ferrugineus</b> (Labill.) Sweet, Hort. Brit. ed.1, 221 (1826)                    | 2:335 |
|    | <i>Ozothamnus gunnii</i> Hook.f. = <b>Ozothamnus bracteolatus</b>                              | 2:333 |
| e  | <b>Ozothamnus hookeri</b> Sonder, Linnaea 25: 509 (1853)                                       | 2:337 |
| e  | <b>Ozothamnus ledifolius</b> (DC.) Hook.f., Fl. Tasm. 1: 204 (1856)                            | 2:334 |
| e  | <b>Ozothamnus lycopodioides</b> Hook.f., London J. Bot. 6: 119 (1847)                          | 2:338 |
|    | <b>Ozothamnus obcordatus</b> DC., Prodr. 6: 165 (1838)   | 2:336 |
| e  | <b>Ozothamnus purpurascens</b> DC., Prodr. 6: 165 (1838)                                       | 2:334 |
| e  | <b>Ozothamnus reflexifolius</b> K.Leeson & Rozefelds, Aust. Syst. Bot. 16(3): 319 (2003)       |       |
| e  | <b>Ozothamnus reticulatus</b> (Labill.) DC., Prodr. 6: 164 (1838)                              | 2:332 |
| e  | <b>Ozothamnus rodwayi</b> Orch., Muelleria 7(4): 522 (1992)                                    | 2:337 |
| e  | var. <b>kingii</b> (W.M.Curtis) P.S.Short, Muelleria 7(4): 522 (1992)                          |       |
| e  | var. <b>oreophilus</b> (W.M.Curtis) P.S.Short, Muelleria 7(4): 522 (1992)                      |       |
| e  | var. <b>rodwayi</b>  |       |
|    | <b>Ozothamnus rosmarinifolius</b> (Labill.) Sweet, Hort. Brit. ed.1, 221 (1826)                | 2:335 |
| e  | <b>Ozothamnus scutellifolius</b> Hook.f., Fl. Tasm. 1: 202 (1856)                              | 2:338 |
| ex | <b>Ozothamnus selaginoides</b> Sonder & F.Muell. in Sonder, Linnaea 25: 510 (1853)             | 2:339 |
|    | <b>Ozothamnus thyrsoideus</b> DC., Prodr. 6: 165 (1838)  | 2:335 |
|    | <b>Ozothamnus turbinatus</b> DC., Prodr. 6: 164 (1838)   | 2:333 |
|    | <b>Paquerina graminea</b> (Labill.) Cass. ex Less., Syn. Gen. Compos. 193 (1832)               | 2:295 |
| i  | <b>Petasites fragrans</b> (Vill.) C.Presl, Fl. Sic. 1: 28 (1826)                               | 2:360 |
|    | <b>Picris angustifolia</b> DC., Prodr. 7: 130 (1838)   | 2:383 |
|    | subsp. <b>angustifolia</b>   |       |
|    | subsp. <b>merxmuelleri</b> Lack & S.Holzappel, Willdenowia 23: 190 (1993)                      |       |
|    | <i>Picris echioides</i> L. = <b>Helminthotheca echioides</b>                                   | 2:383 |
|    | <i>Picris hieracioides</i> sensu W.M.Curtis (1963) = <b>P. angustifolia</b>                    | 2:383 |
|    | <b>Podolepis jaceoides</b> (Sims) Voss, Vilmorin. Blumengart. 1: 537 (1894)                    | 2:348 |
|    | <i>Podosperma angustifolium</i> Labill. = <b>Podotheca angustifolia</b>                        | 2:341 |
| ?x | <b>Podotheca angustifolia</b> (Labill.) Less., Syn. Gen. Comp. 272 (1832)                      | 2:341 |
|    | <b>Pseudognaphalium luteoalbum</b> (L.) Hilliard & B.L.Burt, Bot. J. Linn. Soc. 82: 206 (1981) | 2:318 |
| e  | <b>Pterygopappus lawrencei</b> Hook.f., Lond. J. Bot. 6: 120 (1847)                            | 2:316 |
|    | <b>Rhodanthe anthemoides</b> (Sprengel) Paul G.Wilson, Nuytsia 8: 386 (1992)                   | 2:324 |
|    | <i>Rutidosis multiflora</i> (Nees) Robinson = <b>Siloxerus multiflorus</b>                     | 2:342 |
| i  | <b>Scorzonera laciniata</b> L., Sp. Pl. 2: 791 (1753)  |       |
| i# | <b>Senecio angulatus</b> L.f., Suppl. 369 (1782)   |       |
|    | <b>Senecio biserratus</b> Belcher, Ann. Miss. Bot. Gard. 43: 43 (1956)                         | 2:368 |
| n  | <b>Senecio campylocarpus</b> I.Thomps., Muelleria 20: 139 (2004)                               |       |

|     |  |       |
|-----|--|-------|
| e   | <b>Senecio capillifolius</b> Hook.f., Lond. J. Bot. 6: 123 (1847)                        | 2:365 |
|     | <i>Senecio centropappus</i> F.Muell. = <b>Brachyglottis brunonis</b>                     | 2:365 |
| i   | <b>Senecio elegans</b> L., Sp. Pl. 2: 869 (1753)   | 2:366 |
| n   | <b>Senecio extensus</b> I.Thomps., Muelleria 19: 150 (2004)                              |       |
| n   | <i>Senecio glandulosus</i> (DC.) Sch.Bip. nom. illeg. = <b>Senecio campylocarpus</b>     |       |
|     | <b>Senecio glomeratus</b> Desf. ex Poirlet, Encycl. Suppl. 5: 130 (1817)                 | 2:370 |
|     | subsp. <b>glomeratus</b>   |       |
| n   | subsp. <b>longifructus</b> I.Thomps., Muelleria 19: 148 (2004)                           |       |
|     | <b>Senecio gunnii</b> (Hook.f.) Belcher, Ann. Miss. Bot. Gard. 43: 60 (1956)             | 2:370 |
| n   | <b>Senecio hispidissimus</b> I.Thomps., Muelleria 19: 138 (2004)                         |       |
|     | <b>Senecio hispidulus</b> A.Rich., Voy. Astrolabe 2: 94 t.34 (1834)                      | 2:370 |
| i   | <b>Senecio jacobaea</b> L., Sp. Pl. 2: 870 (1753)  | 2:366 |
|     | <i>Senecio lautus</i> sensu W.M.Curtis (1963) = <b>Senecio pinnatifolius</b>             | 2:365 |
|     | <b>Senecio leptocarpus</b> DC., Prodr. 6: 372 (1838)                                     | 2:364 |
|     | <b>Senecio linearifolius</b> A.Rich., Voy. Astrolabe 2: 129 (1834)                       | 2:366 |
| n   | <b>Senecio longipilus</b> I.Thomps., Muelleria 19: 193 (2004)                            |       |
| ?x  | <b>Senecio macrocarpus</b> Belcher, Muelleria 5: 119 (1983)                              |       |
| n   | <b>Senecio microbasis</b> I.Thomps., Muelleria 19: 175 (2004)                            |       |
|     | <i>Senecio mikanioides</i> Otto ex Harvey = <b>Delairea odorata</b>                      | 2:367 |
|     | <b>Senecio minimus</b> Poirlet, Encycl. Suppl. 5: 130 (1817)                             | 2:368 |
|     | <b>Senecio odoratus</b> Hornem., Hort. Bot. Hafn. 2: 809 (1815)                          | 2:367 |
|     | <b>Senecio orarius</b> J.Black, Trans. R. Soc. S.Austral. 52: 230 (1928)                 |       |
| e   | <b>Senecio papillosus</b> F.Muell., Trans. Philos. Inst. Vict. 2: 69 (1857)              | 2:363 |
|     | <b>Senecio pectinatus</b> DC., Prodr. 6: 372 (1838)                                      | 2:364 |
|     | var. <i>leptocarpus</i> sensu L.Rodway (1903) = <b>Senecio leptocarpus</b>               |       |
| e   | var. <b>ochroleucus</b> F.Muell., Pap. Proc. R. Soc. Tasm. 1870: 16 (1871)               | 2:364 |
| e   | var. <b>pectinatus</b>   |       |
|     | var. <i>pleiocephalus</i> Rodw. = <b>Senecio pinnatifolius</b> var. <b>pleiocephalus</b> |       |
| n   | <b>Senecio phelleus</b> I.Thomps., Muelleria 19: 171 (2004)                              |       |
|     | <b>Senecio pinnatifolius</b> A.Rich. Voy. Astrolabe 2: 117 (1834)                        | 2:365 |
|     | var. <b>pinnatifolius</b>  |       |
|     | var. <b>pleiocephalus</b> (Rodw.) Belcher, Muelleria 9: 127 (1996)                       |       |
| n   | <b>Senecio prenanthoides</b> A.Rich., Voy. Astrolabe 2: 96 (1834)                        |       |
| e   | <b>Senecio primulaefolius</b> F.Muell., Trans. Philos. Inst. Vict. 2: 69 (1857)          | 2:364 |
| n   | <b>Senecio psilocarpus</b> Belcher & Albr., Muelleria 8: 113 (1994)                      |       |
|     | <b>Senecio quadridentatus</b> Labill., Nov. Holl. Pl. 2: 48 t.194 (1806)                 | 2:369 |
|     | <b>Senecio spathulatus</b> A.Rich., Voy. Astrolabe 2: 125 (1834)                         | 2:364 |
|     | <b>Senecio squarrosus</b> A.Rich., Voy. Astrolabe 2: 107 t.35 (1834)                     | 2:369 |
| nex | <b>Senecio tasmanicus</b> I.Thomps., Muelleria 19: 158 (2004)                            |       |
|     | <b>Senecio vagus</b> F.Muell., Trans. Philos. Soc. Vict. 1: 46 (1855)                    |       |
|     | <b>Senecio velleioides</b> A.Cunn. ex DC., Prodr. 6: 374 (1838)                          | 2:365 |
| i   | <b>Senecio vulgaris</b> L., Sp. Pl. 2: 867 (1753)  | 2:367 |
| i#  | <b>Siegesbeckia orientalis</b> L., Sp. Pl. 2: 900 (1753)                                 |       |
|     | <b>Siloxerus multiflorus</b> Nees in Lehm., Pl. Preiss. 2: 244 (1845)                    | 2:342 |

|    |  |       |
|----|--|-------|
| i  | <b>Silybum marianum</b> (L.) Gaertn., Fruct. Sem. Pl. 2: 378 t.162 (1791)                              | 2:378 |
|    | Solenogyne bellioides var. gunnii (Hook.f.) G.Davis = <b>Solenogyne gunnii</b>                         | 2:292 |
|    | <b>Solenogyne dominii</b> L.Adams, Brunonia 2: 58 (1979)   |       |
|    | <b>Solenogyne gunnii</b> (Hook.f.) Cabrera, Blumea 14: 307 (1966)                                      | 2:292 |
| i  | <b>Soliva sessilis</b> Ruiz Lopez & Pavon, Fl. Peruv. et Chil. Prodr. 113, t.24 (1794)                 | 2:358 |
| i  | <b>Sonchus asper</b> (L.) Hill, Herb. Brit. 1: 47 t.34 (1769)  | 2:390 |
| i  | subsp. <b>asper</b>  |       |
| ni | subsp. <b>glaucescens</b> (Jord.) Ball, J. Linn. Soc. Bot. 16: 548 (1878)                              |       |
| n  | <b>Sonchus hydrophilus</b> Boulos in H.Eichler, Fl. South Aust. edn.2, suppl., 331 (1965)              |       |
|    | Sonchus megalocarpus (Hook.f.) J.Black = <b>Actites megalocarpa</b>                                    | 2:390 |
| i  | <b>Sonchus oleraceus</b> L., Sp. Pl. 2: 794 (1753)   | 2:389 |
| i  | <b>Tanacetum cinerariifolium</b> (Trev.) Schultz-Bip., Tanaceteeen 58 (1844)                           |       |
| i  | <b>Tanacetum parthenium</b> (L.) Schultz-Bip., Tanaceteeen 55 (1844)                                   | 2:353 |
| i  | <b>Tanacetum vulgare</b> L., Sp. Pl. 2: 844 (1753)   | 2:360 |
|    | <b>Taraxacum aristum</b> Hagl. & Markl., Bot. Not. 117: 197 figs.198 & 199 (1964)                      |       |
|    | <b>Taraxacum cygnorum</b> Hand.-Maz., Monogr. Gatt. Taraxacum 55 (1907)                                |       |
| i# | <b>Taraxacum kok-saghyz</b> Rodin, Acta Inst. Bot. Acad. Sci. U.R.S.S. 1: 187 (1933)                   | 2:387 |
| i  | <b>Taraxacum officinale</b> Weber in Wiggers, Prim. Fl. Holsat. 56 (1780)                              | 2:387 |
| i  | <b>Tolpis barbata</b> (L.) Gaertner, Fruct. Sem. Pl. 2: 372 (1791)                                     | 2:381 |
|    | Tolpis umbellata Bertol. = <b>Tolpis barbata</b>   | 2:381 |
| i  | <b>Tragopogon porrifolius</b> L., Sp. Pl. 2: 789 (1753)  | 2:390 |
|    | Tripleurospermum inodorum (L.) Schultz-Bip. = <b>Tripleurospermum maritimum</b> subsp. <b>inodorum</b> |       |
| it | <b>Tripleurospermum maritimum</b> (L.) W.Koch, Syn. Fl. Germ. edn.2 1026 (1845)                        |       |
| it | subsp. <b>inodorum</b> (L.) Applequist, Taxon 51: 760 (2002)   |       |
|    | <b>Triptilodiscus pygmaeus</b> Turcz., Bull. Soc. Nat. Mosc. 24: 66 (1851)                             |       |
| it | <b>Urospermum dalechampii</b> (L.) Scop. ex F.W.Schmidt, Samml. Phys. Aufs. 1: 276 (1795)              | 2:391 |
| i  | <b>Vellereophyton dealbatum</b> (Thunb.) Hilliard & B.L.Burt, Bot. J. Linn. Soc. 82: 210 (1981)        | 2:318 |
|    | <b>Vittadinia australasica</b> (Turcz.) N.Burb., Brunonia 5: 42 (1982)                                 |       |
|    | var. <b>oricola</b> N.Burb., Brunonia 5: 44 (1982)   |       |
|    | <b>Vittadinia cuneata</b> DC., Prodr. 5: 281 (1836)  | 2:313 |
|    | var. <b>cuneata</b>  |       |
|    | <b>Vittadinia gracilis</b> (Hook.f.) N.Burb., Brunonia 5: 54 (1982)                                    | 2:313 |
| ?x | <b>Vittadinia megacephala</b> (F.Muell. ex Benth.) J.Black, Trans. R. Soc. S.Austral. 52: 229 (1928)   | 2:314 |
|    | <b>Vittadinia muelleri</b> N.Burb., Proc. Linn. Soc. N.S.W. 93: 440 fig.1D (1969)                      |       |
|    | Vittadinia triloba sensu W.M.Curtis (1963) = <b>Vittadinia cuneata</b>                                 | 2:313 |
|    | var. lanuginosa J.Black = <b>Vittadinia gracilis</b>   | 2:313 |
|    | var. megacephala (F.Muell. ex Benth.) Ewart = <b>Vittadinia megacephala</b>                            | 2:314 |
| i  | <b>Xanthium spinosum</b> L., Sp. Pl. 2: 987 (1753)   | 2:349 |
|    | <b>Xerochrysum bicolor</b> (Lindley) R.J.Bayer, Kew Bull. 56(4): 1014 (2001)                           | 2:331 |
|    | <b>Xerochrysum bracteatum</b> (Vent.) Tzvelev, Novosti Sist. Vyssh. Rast. 27: 151 (1990)               |       |
|    | <b>Xerochrysum palustre</b> (Flann) R.J.Bayer, Kew Bull. 56(4): 1014 (2001)                            |       |
|    | <b>Xerochrysum papillosum</b> (Labill.) R.J.Bayer, Kew Bull. 56(4): 1015 (2001)                        | 2:331 |
|    | <b>Xerochrysum subundulatum</b> (Schultz-Bip.) R.J.Bayer, Kew Bull. 56(4): 1015 (2001)                 | 2:330 |

## BASELLACEAE

- i **Anredera cordifolia** (Tenore) Steenis, Fl. Malesiana 5: 303 fig.2 a-j (1957) 3:580

## BERBERIDACEAE

- i **Berberis darwinii** Hook., Icon. Pl. 7: t.672 (1844)

## BETULACEAE

- i **Alnus cordata** (Lois.) Desf. in K.Koch, Dendrologie 2(1): 634 (1872) 3:645

## BIGNONIACEAE

- Pandorea pandorana** (Andrews) Steenis, Bull. Jard. Bot. Buitenz. 10: 198 (1928) 3:538

## BORAGINACEAE

- i **Amsinckia calycina** (Moris) Chater in Heywood, Bot. J. Linn. Soc. 64(4): 380 (1971) 3:489  
*Amsinckia hispida* (Ruiz Lopez & Pavon) I.M.Johnston = **Amsinckia calycina** 3:489
- i **Anchusa arvensis** (L.) M.Bieb., Fl. Taur.-Caucas. 1: 123 (1808)  
*Anchusa sempervirens* L. = **Pentaglottis sempervirens** 3:491  
**Austrocynoglossum latifolium** (R.Br.) R.R.Mill, Notes Roy. Bot. Gard. Edinburgh 46: 44 (1989) 3:488
- i **Borago officinalis** L., Sp. Pl. 1: 137 (1753) 3:490
- i **Buglossoides arvensis** (L.) I.M.Johnston, J. Arnold Arbor. 35: 42 (1954) 3:494  
**Cynoglossum australe** R.Br., Prodr. 495 (1810) 3:489  
*Cynoglossum latifolium* R.Br. = **Austrocynoglossum latifolium** 3:488  
**Cynoglossum suaveolens** R.Br., Prodr. 495 (1810) 3:488  
*Echium lycopsis* sensu W.M.Curtis (1967) = **Echium plantagineum** 3:495
- i **Echium plantagineum** L., Mant. Pl. 2: 202 (1771) 3:495
- i **Echium vulgare** L., Sp. Pl. 1: 139 (1753) 3:495  
*Lithospermum arvense* L. = **Buglossoides arvensis** 3:494
- it# **Lithospermum officinale** L., Sp. Pl. 1: 132 (1753) 3:494  
*Lycopsis arvensis* L. = **Anchusa arvensis**
- i **Myosotis arvensis** (L.) Hill, Veg. Syst. 7: 55 (1764)  
**Myosotis australis** R.Br., Prodr. 495 (1810) 3:492  
*Myosotis caespitosa* Schultz = **Myosotis laxa** subsp. **caespitosa** 3:493
- i **Myosotis discolor** Pers., Syst. Veg. 190 (1797) 3:493  
**Myosotis exarrhena** F.Muell., Syst. Census Austral. Pl. Suppl. 4: 7 (1889) 3:492
- i **Myosotis laxa** Lehm., Pl. Asperif. Nucif. 83 (1818)  
subsp. **caespitosa** (Schultz) Hyl. ex Nordh., Norsk. Fl. 529 (1940) 3:493
- i **Myosotis scorpioides** L., Sp. Pl. 1: 131 (1753) 3:492  
*Myosotis suaveolens* (R.Br.) Poiret = **Myosotis exarrhena** 3:492
- i **Myosotis sylvatica** Hoffm., Deutschl. Fl. 61 (1791) 3:493
- i **Nonea lutea** (Desr.) Rchb. ex A.DC. in DC., Prodr. 10: 28 (1846)
- i **Pentaglottis sempervirens** (L.) Tausch ex L.Bailey, Man. Cult. Pl. ed.2, 837 (1949) 3:491  
*Symphytum peregrinum* Ledeb. = **Symphytum Xuplandicum** 3:490
- i# **Symphytum Xuplandicum** Nyman, Syll. 80 (1854) 3:490

**BRASSICACEAE (CRUCIFERAE)**

|     |  |      |
|-----|--|------|
| i   | <b>Arabidopsis thaliana</b> (L.) Heynh. in Holl & Heynh., Fl. Sachs. 1: 538 (1842)                     | 1:50 |
|     | <b>Ballantinia antipoda</b> (F.Muell.) E.Shaw, Contr. Gray Herb. 205: 161 (1974)                       | 1:42 |
| e   | <b>Barbarea australis</b> Hook.f., Fl. Nov.-Zel. 1: 14 (1852)  | 1:47 |
| i   | <b>Barbarea intermedia</b> Bor., Fl. Centr. Fr. 2: 48 (1840)   | 1:48 |
| i   | <b>Barbarea verna</b> (Mill.) Asch., Fl. Brandenburg 1: 36 (1860)                                      | 1:48 |
| i#  | <b>Brassica juncea</b> (L.) Czern., Consp. Pl. Chark. 8 (1859)   |      |
| i   | <b>Brassica napus</b> L., Sp. Pl. 2: 666 (1753)  | 1:34 |
| i   | <b>Brassica nigra</b> (L.) Koch in Roehl., Deutschl. Fl. ed.3, 4: 713 (1833)                           | 1:35 |
| ni# | <b>Brassica oleracea</b> L., Sp. Pl. 2: 667 (1753)   |      |
| i   | <b>Brassica rapa</b> L., Sp. Pl. 2: 666 (1753)   | 1:34 |
| i   | <b>Brassica tournefortii</b> Gouan, Illust. Observ. Bot. 44: t.20A (1773)                              |      |
| i   | <b>Cakile edentula</b> (Bigelow) Hook., Fl. Bot.-Amer. 1: 59 (1830)                                    | 1:38 |
| i   | <b>Cakile maritima</b> Scop., Fl. Carniol. ed.2, 2: 35 (1772)  | 1:38 |
| i   | <b>Camelina alyssum</b> (Mill.) Thell., Verz. Tausch Säm. Früchte Zürich 10 (1906)                     | 1:51 |
| i   | <b>Camelina sativa</b> (L.) Crantz, Stirp. Austriac. fasc.1: 18 (1762)                                 | 1:51 |
| i   | <b>Capsella bursa-pastoris</b> (L.) Medikus, Pflanzengatt. 1: 85 (1792)                                | 1:43 |
|     | <b>Capsella tasmanica</b> (Hook.) F.Muell. ex B.D.Jackson = <b>Hutchinsia tasmanica</b>                | 1:43 |
|     | <b>Cardamine astoniae</b> I.Thomps., Muelleria 9: 156 (1996)   |      |
|     | <b>Cardamine</b> aff. <b>corymbosa</b> Hook.f., sensu H.J.Hewson, Fl. Australia 8: 313 (1982)          |      |
|     | <b>Cardamine depressa</b> Hook.f., var. <b>stellata</b> (Hook.f.) Hook.f. = <b>Cardamine corymbosa</b> |      |
| i   | <b>Cardamine flexuosa</b> With., Bot. Arr. Brit. Pl. ed.3, 3: 578 (1796)                               |      |
|     | <b>Cardamine franklinensis</b> I.Thomps., Muelleria 9: 152 (1996)                                      |      |
|     | <b>Cardamine gunnii</b> Hewson, Fl. Australia 8: 390 (1982)  | 1:46 |
|     | <b>Cardamine heterophylla</b> sensu Curtis & Morris (1975) = <b>Cardamine gunnii</b>                   | 1:46 |
| i   | <b>Cardamine hirsuta</b> L., Sp. Pl. 2: 655 (1753)   | 1:46 |
|     | <b>Cardamine lilacina</b> Hook., Companion Bot. Mag. 1: 273 (1835)                                     |      |
|     | <b>Cardamine papillata</b> I.Thomps., Muelleria 9: 161 (1996)  |      |
|     | <b>Cardamine paucijuga</b> Turcz., Bull. Soc. Nat. Mosc. 27(2): 295 (1854)                             |      |
|     | <b>Cardamine tenuifolia</b> Hook., J. Bot. (Hooker) 1: 247 (1837)                                      | 1:46 |
| nx  | <b>Cardamine tryssa</b> I.Thomps., Muelleria 18: 27 (2004)   |      |
| i   | <b>Cardaria draba</b> (L.) Desv., J. Bot. (Desvaux) 3: 163 (1815)                                      | 1:42 |
| i#  | <b>Carrichtera annua</b> (L.) DC., Mém. Mus. Hist. Nat. Paris 7: 244, 250 (1821)                       |      |
| e   | <b>Cheesemaniania radicata</b> (Hook.f.) O.E.Schulz, Notizbl. Bot. Gart. Berlin-Dahlem 10: 552 (1929)  | 1:47 |
| i   | <b>Coronopus didymus</b> (L.) Smith, Fl. Brit. 2: 691 (1800)   | 1:41 |
| i   | <b>Coronopus squamatus</b> (Forsk.) Asch., Fl. Brandenburg 1(2): 62 (1864)                             | 1:41 |
|     | <b>Cuphonotus antipodus</b> (F.Muell.) J.Black = <b>Ballantinia antipoda</b>                           | 1:42 |
| i   | <b>Diplotaxis muralis</b> (L.) DC., Syst. Nat. 2: 634 (1821)   | 1:36 |
| i   | <b>Diplotaxis tenuifolia</b> (L.) DC., Syst. Nat. 2: 632 (1821)  | 1:36 |
|     | <b>Draba muralis</b> sensu Curtis & Morris (1975) = <b>Draba nemorosa</b>                              | 1:45 |
| i   | <b>Draba nemorosa</b> L., Sp. Pl. 2: 643 (1753)  | 1:45 |
| i   | <b>Erophila verna</b> (L.) Chevall., Fl. Gén. Env. Paris 2: 898 (1827)                                 | 1:45 |
| i   | <b>Eruca vesicaria</b> (L.) Cav., Descr. Pl. 426 (1801)  |      |
| i   | subsp. <b>sativa</b> (Mill.) Thell. in Hegi, Ill. Fl. Mitt.-Eur. 4(1): 201 (1918)                      |      |

|                         |  |       |
|-------------------------|--|-------|
|                         | <i>Eruca sativa</i> Mill. = <b><i>Eruca vesicaria</i></b> subsp. <b><i>sativa</i></b>                                      |       |
|                         | <b><i>Geococcus pusillus</i></b> J.Drumm. ex Harvey, Hook. J. Bot. Kew Gard. Misc. 7: 52 (1855)                            | 1:50  |
| i                       | <b><i>Hirschfeldia incana</i></b> (L.) Lagrèze-Fossat, Fl. Tarn Garonne 19 (1847)  | 1:35  |
| e                       | <b><i>Hutchinsia tasmanica</i></b> Hook., Icon. Pl. 9: fig.848 (1851)  | 1:43  |
| i                       | <b><i>Hymenolobus procumbens</i></b> (L.) Nutt. in Torr. & A.Gray, Fl. N. Amer. 1: 117 (1838)                              | 1:44  |
| i                       | <b><i>Lepidium africanum</i></b> (N.Burm.) DC., Syst. Nat. 2: 522 (1821)   |       |
| i                       | <b><i>Lepidium campestre</i></b> (L.) R.Br. in Aiton, Hort. Kew. ed.2, 4: 88 (1812)  | 1:40  |
|                         | <b><i>Lepidium desvauxii</i></b> Thell., Neue Denkschr. Schweiz. Naturw. Ges. 41(1): 307 (1906)                            | 1:39  |
| t                       | <b><i>Lepidium flexicaule</i></b> Kirk, Trans. Proc. N.Z. Inst. 14: 380 (1882)   |       |
|                         | <b><i>Lepidium foliosum</i></b> Desv., J. Bot. (Desvaux) 3: 164, 180 (1815)  | 1:39  |
|                         | <i>Lepidium halmaturinum</i> J.Black = <b><i>Lepidium desvauxii</i></b>  | 1:39  |
| i#                      | <b><i>Lepidium heterophyllum</i></b> (DC.) Benth., Cat. Pl. Pyr. Bas-Languedoc 95 (1826)                                   | 1:40  |
|                         | <b><i>Lepidium hyssopifolium</i></b> Desv., J. Bot. (Desvaux) 3: 164, 179 (1815)   |       |
| i                       | <b><i>Lepidium latifolium</i></b> L., Sp. Pl. 2: 644 (1753)  | 1:41  |
|                         | <i>Lepidium praetervisum</i> Domin = <b><i>Lepidium desvauxii</i></b>  | 1:39  |
|                         | <b><i>Lepidium pseudotasmanicum</i></b> Thell., Neue Denkschr. Schweiz. Naturw. Ges. 41(1): 307 (1906)                     | 1:39  |
| i                       | <b><i>Lepidium sativum</i></b> L., Sp. Pl. 2: 644 (1753)   | 1:40  |
| i                       | <b><i>Lobularia maritima</i></b> (L.) Desv., J. Bot. (Desvaux) 3: 162 (1815)   | 1:45  |
| i#                      | <b><i>Lunaria annua</i></b> L., Sp. Pl. 2: 652 (1753)  |       |
|                         | <i>Nasturtium microphyllum</i> Boenn. ex Rchb. = <b><i>Rorippa microphylla</i></b>   | 1:49  |
|                         | <i>Nasturtium officinale</i> R.Br. = <b><i>Rorippa nasturtium-aquaticum</i></b>  | 1:49  |
| i                       | <b><i>Raphanus maritimus</i></b> Smith, Engl. Bot. 23: 1643 (1806)   | 1:37  |
| i                       | <b><i>Raphanus raphanistrum</i></b> L., Sp. Pl. 2: 669 (1753)  | 1:37  |
| i                       | <b><i>Rapistrum rugosum</i></b> (L.) All., Fl. Pedem. 1: 257 t.78 (1755)   | 1:37  |
|                         | <b><i>Rorippa dictyosperma</i></b> (Hook.) L.A.S.Johnson, Contr. N.S.W. Nat. Herb. 3: 97 (1962)                            | 1:49  |
|                         | <b><i>Rorippa gigantea</i></b> (Hook.f.) Garnock-Jones, N.Z. J. Bot. 16: 119 (1978)  | 1:48  |
|                         | <i>Rorippa islandica</i> sensu Curtis & Morris (1975) = <b><i>Rorippa palustris</i></b>                                    | 1:48  |
| i                       | <b><i>Rorippa microphylla</i></b> (Boenn. ex Rchb.) Hyl., Rep. Univ. Inst. Appl. Sci. Reykj. Dept. Ag. Bull. 3: 109 (1948) | 1:49  |
| i                       | <b><i>Rorippa nasturtium-aquaticum</i></b> (L.) Hayek, Sched. Fl. Stiriac. 22 (1905)                                       | 1:49  |
| i                       | <b><i>Rorippa palustris</i></b> (L.) Besser, Enum. Pl. 27 (1822)   | 1:48  |
|                         | <i>Rorippa stylosa</i> sensu Curtis & Morris (1975) = <b><i>Rorippa gigantea</i></b>                                       | 1:48  |
| it#                     | <b><i>Rorippa sylvestris</i></b> (L.) Besser, Enum. Pl. 27 (1822)  |       |
| i                       | <b><i>Sinapis alba</i></b> L., Sp. Pl. 2: 668 (1753)   |       |
| i                       | <b><i>Sinapis arvensis</i></b> L., Sp. Pl. 2: 668 (1753)   | 1:35  |
| i                       | <b><i>Sisymbrium irio</i></b> L., Sp. Pl. 2: 659 (1753)  |       |
| i                       | <b><i>Sisymbrium officinale</i></b> (L.) Scop., Fl. Carniole ed.2, 2: 26 (1772)  | 1:50  |
| i                       | <b><i>Sisymbrium orientale</i></b> L., Cent. Pl. 2: 24 (1756)  | 1:50  |
|                         | <b><i>Stenopetalum lineare</i></b> R.Br. ex DC., Syst. Nat. 2: 513 (1821)  | 1:44  |
| it                      | <b><i>Teesdalia nudicaulis</i></b> (L.) W.T.Aiton, Hortus Kew. ed.2, 4: 83 (1812)  | 1:43  |
| i#                      | <b><i>Thlaspi arvense</i></b> L., Sp. Pl. 2: 646 (1753)  |       |
| <br><b>BRUNONIACEAE</b> |  |       |
|                         | <b><i>Brunonia australis</i></b> Smith ex R.Br., Prodr. 590 (1810)   | 2:405 |



## BUDDLEJACEAE

- i **Buddleja davidii** Franchet, Nouv. Arch. Mus. Hist. Nat. Paris ser.2, 10: 103 (1888)

## CALLITRICHACEAE

- Callitriche brachycarpa** Hegelm., Verh. Bot. Vereins Prov. Brandenburg 10: 115 (1868) 1:195
- i# **Callitriche brutia** Petagna, Inst. Bot. 2: 10 (1787)
- Callitriche sonderi** Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 18 (1867)
- i **Callitriche stagnalis** Scop., Fl. Carniol. ed.2, 2: 251 (1772) 1:195
- Callitriche umbonata** Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 19 (1867) 1:196

## CAMPANULACEAE

- i **Campanula rapunculoides** L., Sp. Pl. 1: 165 (1753) 2:410
- Isotoma fluviatilis** (R.Br.) F.Muell. ex Benth., Fl. Austral. 4: 136 (1868) 2:411
- subsp. **australis** McComb, Contr. N.S.W. Natl. Herb. 4: 109 (1970)
- Lobelia alata** Labill. = **Lobelia anceps** 2:414
- Lobelia anceps** L.f., Suppl. Pl. 395 (1782) 2:414
- Lobelia gibbosa** Labill., Nov. Holl. Pl. 1: 50, t.71 (1805) 2:414
- var. **browniana** (Roem. & Schult.) F.M.Bailey, Queensland Fl. 916 (1900)
- var. **gibbosa**
- Lobelia pratioides** Benth., Fl. Austral. 4: 131 (1868) 2:415
- Lobelia rhombifolia** Vriese in Lehm., Pl. Preiss. 1: 397 (1845) 2:414
- Pratia irrigua** (R.Br.) Benth., Fl. Austral. 4: 132 (1868) 2:412
- Pratia pedunculata** (R.Br.) Benth., Fl. Austral. 4: 133 (1868) 2:413
- Pratia platycalyx** (F.Muell.) Benth. = **Pratia irrigua** 2:412
- Pratia surrepens** (Hook.f.) F.E.Wimmer, Pflanzenr. 106: 108 (1943) 2:412
- Wahlenbergia ceracea** Loth., Vict. Nat. 72: 166 (1956) 2:407
- Wahlenbergia consimilis** Loth. = **Wahlenbergia stricta** 2:408
- Wahlenbergia gracilentata** Loth., Proc. Linn. Soc. N.S.W. 71: 217 (1947) 2:407
- Wahlenbergia gracilis** (Forst.f.) A.DC., Monogr. Camp. 142 (1830) 2:409
- Wahlenbergia gymnoclada** Loth., Proc. Linn. Soc. N.S.W. 71: 227 (1947) 2:408
- Wahlenbergia littoricola** P.J.Smith = **Wahlenbergia multicaulis** pro parte maj. 2:409
- Wahlenbergia multicaulis** Benth. in Endl., Enum. Pl. Hügel. 75 (1837) 2:409
- Wahlenbergia quadrifida** (R.Br.) A.DC. = **Wahlenbergia gracilis** 2:409
- e **Wahlenbergia saxicola** A.DC., Monogr. Camp. 144 (1830) 2:410
- Wahlenbergia stricta** (R.Br.) Sweet, Hort. Brit. ed.2, 593 (1830) 2:408
- Wahlenbergia tadgellii** Loth. = **Wahlenbergia multicaulis** 2:409

## CAPRIFOLIACEAE

- i **Leycesteria formosa** Wallich in Roxb., Fl. Ind. 2: 182 (1824) 2:266
- ni **Lonicera japonica** Thunb. ex Murray, Syst. Veg. edn.14, 216 (1784)
- i# **Lonicera periclymenum** L., Sp. Pl. 1: 173 (1753) 2:266
- Sambucus gaudichaudiana** DC., Prodr. 4: 322 (1830) 2:265
- i **Sambucus nigra** L., Sp. Pl. 1: 269 (1753) 2:265

## CARYOPHYLLACEAE

- i **Agrostemma githago** L., Sp. Pl. 1: 435 (1753) 1:69
- i **Arenaria leptoclados** (Rchb.) Guss., Fl. Sicul. Syn. 2: 824 (1845)
- i **Arenaria serpyllifolia** L., Sp. Pl. 1: 423 (1753) 1:72
- i **Cerastium balearicum** Herm., Verh. Bot. Vereins Prov. Brandenburg 54: 247 (1913)
- i **Cerastium fontanum** Baumg., Enum. Stirp. Transs. 1: 425 (1816) 1:69
- i subsp. **vulgare** (Hartm.) Greuter & Burdet, Willdenowia 12: 37 (1982) 1:69
- i **Cerastium glomeratum** Thuill., Fl. Paris ed.2: 226 (1799) 1:69
- i **Cerastium semidecandrum** L., Sp. Pl. 1: 438 (1753)
- Cerastium vulgare** Hartm. = **Cerastium fontanum** subsp. **vulgare** 1:69
- Colobanthus affinis** (Hook.) Hook.f., Fl. Tasm. 1: 45 (1855) 1:74
- Colobanthus apetalus** (Labill.) Druce, Bot. Exch. Club Soc. Brit. Isles 1916: 616 (1917) 1:74
- var. **apetalus**
- e **Colobanthus curtisiae** J.West in Banks et al., Aspects Tasm. Bot. 75 (1991)
- Colobanthus pulvinatus** F.Muell., Trans. Phil. Soc. Vict. 1: 101 (1855)
- i **Dianthus armeria** L. Sp. Pl. 1: 410 (1753)
- Gypsophila australis** (Schldl.) A.Gray = **Gypsophila tubulosa** 1:66
- i **Gypsophila tubulosa** (Jaub. & Spach) Boiss., Diagn. Pl. Orient. Nov. 11 (1842) 1:66
- Minuartia hybrida** sensu Cens. Vasc. Pl. Tasm. edn.1 = **Minuartia mediterranea**
- i **Minuartia mediterranea** (Link) K.Maly, Glasn. Zemaljsk. Muz. Bosni Hercegovini 20: 363 (1908)
- i **Melandrium album** (Mill.) Garcke = **Silene latifolia**
- i **Moenchia erecta** (L.) Gaertn, Mey. & Scherb., Oekon.-Tech. Fl. Wetterau 1: 219 (1799) 1:70
- i **Paronychia brasiliiana** DC. in Lam. & Poiret, Encycl. 5: 23 (1804)
- i **Petrorhagia nanteuillii** (Burnat) Ball & Heywood, Bull. Brit. Mus. Bot. 3: 164 (1964) 1:65
- Petrorhagia prolifera** sensu Curtis & Morris (1975) = **Petrorhagia nanteuillii** 1:65
- i **Petrorhagia velutina** (Guss.) Ball & Heywood, Bull. Brit. Mus. Bot. 3: 166 (1964) 1:66
- i **Polycarpon tetraphyllum** (L.) L., Syst. Nat. ed.10, 2: 881 (1759) 1:76
- i **Sagina apetala** Ard., Animad. Bot. Spec. Alt. 2: 22 (1764) 1:73
- e **Sagina diemensis** L.G.Adams, Muellera 9: 64 (1996)
- i **Sagina maritima** G.Don, Herb. Brit. fasc.7: 155 (1806) 1:73
- Sagina namadgi** L.G.Adams, Muellera 9: 63 (1996)
- i **Sagina procumbens** L., Sp. Pl. 1: 128 (1753) 1:73
- i **Saponaria officinalis** L., Sp. Pl. 1: 408 (1753) 1:67
- i **Scleranthus annuus** L., Sp. Pl. 1: 406 (1753) 1:77
- Scleranthus biflorus** (Forst. & Forst.f.) Hook.f., Fl. Nov.-Zel. 1: 74 (1852) 1:76
- Scleranthus brockiei** P.A.Williamson, Rec. Dom. Mus. Wellington 3: 16 (1956)
- Scleranthus diander** R.Br., Prodr. 412 (1810) 1:77
- Scleranthus fasciculatus** (R.Br.) Hook.f., Fl. Tasm. 1: 42 (1855)
- i **Silene alba** (Mill.) E.H.Krause = **Silene latifolia** 1:67
- i **Silene conica** L., Sp. Pl. 1: 418 (1753)
- i **Silene dichotoma** Ehrh., Beitr. Natur. 7: 144 (1792)
- i **Silene gallica** L., Sp. Pl. 1: 417 (1753) 1:68
- i **Silene latifolia** Poiret, Voy. Barb. 2: 165 (1789) 1:67

|   |   |      |
|---|---|------|
| i | <b>Silene longicaulis</b> Pourret ex Lagasca, Gen. Sp. Pl. 15 (1816)                            |      |
| i | <b>Silene nocturna</b> L., Sp. Pl. 1: 416 (1753)  | 1:68 |
| i | <i>Silene pratensis</i> (Rafn.) Godron & Gren. = <b>Silene latifolia</b>                        | 1:67 |
| i | <b>Silene vulgaris</b> (Moench) Garcke, Fl. N.-Mitt.-Deutschland ed.9, 64 (1869)                | 1:68 |
| i | <b>Spergula arvensis</b> L., Sp. Pl. 1: 440 (1753)  | 1:74 |
| i | <b>Spergularia bocconii</b> (Scheele) Graebn., Syn. Mitteleur. Fl. 5: 849 (1919)                |      |
| i | <b>Spergularia marina</b> (L.) Griseb., Spic. Fl. Rumel. 1: 213 (1843)                          | 1:75 |
|   | <b>Spergularia aff. media</b> (L.) C.Presl, Fl. Sicula 161 (1826)                               | 1:75 |
| i | <b>Spergularia rubra</b> (L.) J. & C.Presl, Fl. Czech. 94 (1819)                                | 1:75 |
|   | <b>Stellaria angustifolia</b> Hook., J. Bot. (Hooker) 1: 250 (1834)                             |      |
|   | <b>Stellaria caespitosa</b> Hook.f., J. Bot. (Hooker) 2: 411 (1840)                             |      |
|   | <b>Stellaria flaccida</b> Hook., Comp. Bot. Mag. 1: 275 (1836)                                  | 1:71 |
| i | <b>Stellaria graminea</b> L., Sp. Pl. 1: 422 (1753)   | 1:72 |
| i | <b>Stellaria media</b> (L.) Cirillo, Essent. Pl. Char. Comment. 36 (1784)                       | 1:71 |
|   | <b>Stellaria multiflora</b> Hook., Comp. Bot. Mag. 1: 275 (1836)                                | 1:71 |
| i | <b>Stellaria pallida</b> (Dumort.) Crep., Man. Fl. Belgique edn.2, 19 (1866)                    | 1:72 |
|   | <i>Stellaria palustris</i> sensu auct. Aust. non Ehrh. ex Retz. = <b>Stellaria angustifolia</b> | 1:71 |
|   | <b>Stellaria pungens</b> Brongn. in Duperrey, Voy. Coquille 16: t.78 (1834)                     | 1:71 |
| i | <b>Vaccaria hispanica</b> (Miller) Rauschert, Feddes Repert. 73: 52 (1966)                      | 1:66 |
|   | <i>Vaccaria pyramidata</i> Medik. = <b>Vaccaria hispanica</b>                                   | 1:66 |

#### CASUARINACEAE

|   |   |       |
|---|---|-------|
| e | <b>Allocasuarina crassa</b> L.A.S.Johnson, Fl. Australia 3: 194 (1989)                                |       |
| e | <b>Allocasuarina duncanii</b> L.A.S.Johnson & D.I.Morris, Telopea 5(4): 793 (1994)                    |       |
|   | <b>Allocasuarina littoralis</b> (Salisb.) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 76 (1982)          | 3:644 |
| e | <b>Allocasuarina monilifera</b> (L.A.S.Johnson) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 76 (1982)    | 3:644 |
|   | <b>Allocasuarina paludosa</b> (Sieber ex Sprengel) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 77 (1982) | 3:644 |
|   | <b>Allocasuarina verticillata</b> (Lam.) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 79 (1982)           | 3:644 |
| e | <b>Allocasuarina zephyrea</b> L.A.S.Johnson, Fl. Australia 3: 199 (1989)                              |       |
|   | <i>Casuarina bicuspidata</i> sensu L.Rodway (1903) = <b>Allocasuarina paludosa</b>                    | 3:644 |
|   | <i>Casuarina distyla</i> sensu L.Rodway (1903) = <b>Allocasuarina monilifera</b>                      | 3:644 |
|   | <i>Casuarina littoralis</i> Salisb. = <b>Allocasuarina littoralis</b>                                 | 3:644 |
|   | <i>Casuarina monilifera</i> L.A.S.Johnson = <b>Allocasuarina monilifera</b>                           | 3:644 |
|   | <i>Casuarina paludosa</i> Sieber ex Sprengel = <b>Allocasuarina paludosa</b>                          | 3:644 |
|   | <i>Casuarina quadrivalvis</i> Labill. = <b>Allocasuarina verticillata</b>                             | 3:644 |
|   | <i>Casuarina stricta</i> Dryander = <b>Allocasuarina verticillata</b>                                 | 3:644 |
|   | <i>Casuarina suberosa</i> Otto & A.Dietr. = <b>Allocasuarina littoralis</b>                           | 3:644 |

#### CHENOPODIACEAE

|    |   |       |
|----|---|-------|
|    | <i>Arthrocnemum arbusculum</i> (R.Br.) Moq. = <b>Sclerostegia arbuscula</b> | 3:577 |
| ?i | <b>Atriplex australasica</b> Moq., Chenop. Monogr. Enum. 59 (1840)          |       |
|    | <b>Atriplex billardierei</b> (Moq.) Hook.f., Fl. Nov.-Zel. 1: 215 (1853)    | 3:575 |
|    | <b>Atriplex cinerea</b> Poiret in Lam., Encycl. Suppl. 1: 471 (1811)        | 3:574 |
|    | <i>Atriplex hastata</i> sensu W.M.Curtis (1967) = <b>Atriplex prostrata</b> | 3:575 |

|  |  |       |
|--|--|-------|
| i  | <b>Atriplex hortensis</b> L., Sp. Pl. 2: 1053 (1753)   | 3:576 |
|  | <b>Atriplex paludosa</b> R.Br., Prodr. 406 (1810)  | 3:574 |
|  | subsp. <b>paludosa</b>   |       |
| i  | <b>Atriplex patula</b> L., Sp. Pl. 2: 1053 (1753)  | 3:575 |
| i  | <b>Atriplex prostrata</b> Boucher ex DC. in Lam. & DC., Fl. Franc. ed.3, 3: 387 (1805)               | 3:575 |
| i#   | <b>Atriplex semibaccata</b> R.Br., Prodr. 406 (1810)   |       |
|  | <b>Atriplex suberecta</b> I.Verd., Bothalia 6: 418 (1954)  |       |
| i#   | <b>Bassia scoparia</b> (L.) A.J.Scott, Feddes Repert. 89: 108 (1978)                                 |       |
| i  | <b>Beta vulgaris</b> L., Sp. Pl. 1: 222 (1753)   | 3:573 |
| i  | subsp. <b>maritima</b> (L.) Arcangeli, Comp. Fl. Ital. 593 (1882)                                    | 3:573 |
| i  | <b>Chenopodium album</b> L., Sp. Pl. 1: 219 (1753)   | 3:571 |
| x?   | <b>Chenopodium erosum</b> R.Br., Prodr. 407 (1810)   |       |
| i?   | <b>Chenopodium glaucum</b> L., Sp. Pl. 1: 220 (1753)   | 3:572 |
|  | subsp. <b>ambiguum</b> sensu W.M.Curtis (1967) = <b>Chenopodium glaucum</b>                          | 3:572 |
| i  | <b>Chenopodium murale</b> L., Sp. Pl. 1: 219 (1753)  | 3:572 |
| i  | <b>Chenopodium pumilio</b> R.Br., Prodr. 407 (1810)  | 3:573 |
| i  | <b>Chenopodium vulvaria</b> L., Sp. Pl. 1: 220 (1753)  |       |
|  | <b>Einadia nutans</b> (R.Br.) A.J.Scott, Feddes Repert. 89: 3 (1978)                                 | 3:570 |
|  | subsp. <b>nutans</b>   |       |
|  | <b>Rhagodia baccata</b> sensu W.M.Curtis (1967) = <b>Rhagodia candolleana</b>                        | 3:570 |
|  | <b>Rhagodia candolleana</b> Moq., Chemsp. Monogr. Enum. 10 (1840)                                    | 3:570 |
|  | subsp. <b>candolleana</b>  |       |
|  | <b>Rhagodia nutans</b> R.Br. = <b>Einadia nutans</b>   | 3:570 |
|  | <b>Salicornia blackiana</b> Ulbrich = <b>Sarcocornia blackiana</b>                                   | 3:579 |
|  | <b>Salicornia quinqueflora</b> Bunge ex Ung.-Sternb. = <b>Sarcocornia quinqueflora</b>               | 3:578 |
| i  | <b>Salsola kali</b> L., Sp. Pl. 1: 222 (1753)  |       |
|  | <b>Sarcocornia blackiana</b> (Ulbrich) A.J.Scott, Bot. J. Linn. Soc. 75: 369 (1978)                  | 3:579 |
|  | <b>Sarcocornia quinqueflora</b> (Bunge ex Ung.-Sternb.) A.J.Scott, Bot. J. Linn. Soc. 75: 368 (1978) | 3:578 |
|  | subsp. <b>quinqueflora</b>   |       |
|  | subsp. <b>tasmanica</b> Paul G.Wilson, Nuytsia 3: 74 (1980)  |       |
|  | <b>Sclerostegia arbuscula</b> (R.Br.) Paul G.Wilson, Nuytsia 3: 20 (1980)                            | 3:577 |
|  | <b>Suaeda australis</b> (R.Br.) Moq., Ann. Sci. Nat. 23: 318 (1831)                                  | 3:580 |
| i  | <b>Suaeda maritima</b> (L.) Dumort., Fl. Belg. 22 (1827)   |       |
|  | <b>Threlkeldia diffusa</b> R.Br., Prodr. 410 (1810)  | 3:577 |
| <br><b>CISTACEAE</b>                             |  |       |
| i#   | <b>Cistus psilosepalus</b> Sweet, Cistineae t.33 (1826)  |       |
| <br><b>CLUSIACEAE (GUTTIFERAE, HYPERICACEAE)</b> |  |       |
| i  | <b>Hypericum androsaemum</b> L., Sp. Pl. 2: 784 (1753)   | 1:82  |
| i  | <b>Hypericum calycinum</b> L., Mant. Pl. 1: 106 (1767)   | 1:82  |
|  | <b>Hypericum gramineum</b> Forst.f., Fl. Ins. Austral. 53 (1786)                                     | 1:81  |
| ni   | <b>Hypericum humifusum</b> L., Sp. Pl. 1: 785 (1753)   |       |
|  | <b>Hypericum japonicum</b> Thunb., Fl. Jap. 295 (1784)   | 1:81  |

|                       |   |       |
|-----------------------|---|-------|
| i                     | <b>Hypericum perforatum</b> L., Sp. Pl. 2: 785 (1753)   | 1:81  |
| i                     | <b>Hypericum tetrapterum</b> Fries, Novit. Fl. Suec. Alt. 236 (1828)                                    |       |
| <b>CONVOLVULACEAE</b> |   |       |
|                       | <b>Calystegia sepium</b> (L.) R.Br., Prodr. 483 (1810)  | 3:498 |
| i                     | <b>Calystegia silvatica</b> (Kit.) Griseb., Spic. Fl. Rumel. 1: 74 (1843)                               | 3:498 |
|                       | <b>Calystegia soldanella</b> (L.) R.Br., Prodr. 484 (1810)  | 3:499 |
|                       | <b>Convolvulus angustissimus</b> R.Br., Prodr. 482 (1810)   | 3:497 |
|                       | var. <b>angustissimus</b>   |       |
| i                     | <b>Convolvulus arvensis</b> L., Sp. Pl. 1: 153 (1753)   | 3:497 |
|                       | <i>Convolvulus erubescens</i> sensu W.M.Curtis (1967) = <b>Convolvulus angustissimus</b>                | 3:497 |
|                       | <b>Dichondra repens</b> Forst. & Forst.f., Char. Gen. Pl. 40: t.20 (1776)                               | 3:499 |
|                       | <b>Wilsonia backhousei</b> Hook.f., Lond. J. Bot. 6: 275 (1847)   | 3:500 |
|                       | <b>Wilsonia humilis</b> R.Br., Prodr. 490 (1810)  | 3:500 |
|                       | <b>Wilsonia rotundifolia</b> Hook., Icon. Pl. 5: t.410 (1842)   |       |
| <b>CRASSULACEAE</b>   |   |       |
|                       | <b>Crassula decumbens</b> Thunb., Prodr. Pl. Cap. 54 (1794)   | 1:185 |
|                       | var. <b>decumbens</b>   |       |
|                       | <b>Crassula exserta</b> (Reader) Ostenf., Dansk. Bot. Ark. 2: 45 (1918)                                 | 1:184 |
|                       | <b>Crassula helmsii</b> (Kirk) Cockayne, Trans. N.Z. Inst. 39: 349 (1907)                               | 1:185 |
|                       | <i>Crassula macrantha</i> (Hook.f.) Diels & Pritzel = <b>Crassula decumbens</b>                         | 1:185 |
| t                     | <b>Crassula moschata</b> Forst.f., Comm. Gotting. 9: 26 (1789)  |       |
| i                     | <b>Crassula natans</b> Thunb., Prodr. Pl. Cap. 54 (1794)  |       |
|                       | var. <b>minus</b> (Eckl. & Zeyher) Rowley, Cactus Succ. J. Gt. Brit. 40: 53 (1978)                      |       |
|                       | <b>Crassula pedicellosa</b> (F.Muell.) Ostenf., Dansk. Bot. Ark. 2: 42 (1918)                           |       |
|                       | <b>Crassula peduncularis</b> (Smith) Meigen, Bot. Jahrb. 17: 239 (1893)                                 | 1:184 |
|                       | <b>Crassula sieberiana</b> (Schultes & Schultes f.) Druce, Bot. Exch. Club Brit. Isles 1916: 618 (1917) | 1:184 |
|                       | subsp. <b>sieberiana</b>  |       |
|                       | subsp. <b>tetramera</b> Toelken, J. Adelaide Bot. Gard. 3: 77 (1981)                                    |       |
| i                     | <b>Sedum acre</b> L., Sp. Pl. 1: 432 (1753)   | 1:186 |
| i                     | <b>Sedum forsterianum</b> Smith in Sowerby, Engl. Bot. 26: t.1802 (1808)                                |       |
| i                     | <b>Sedum reflexum</b> L., Sp. Pl. ed.2, 618 (1762)  |       |
| i                     | <b>Sedum sexangulare</b> L., Sp. Pl. 1: 432 (1753)  | 1:186 |
| <b>CUCURBITACEAE</b>  |   |       |
| i                     | <b>Cucumis myriocarpus</b> Naudin, Ann. Sci. Nat. Bot. ser.4, 11: 22 (1859)                             |       |
| i#                    | <b>Ecballium elaterium</b> (L.) A.Rich., Dict. Class. Hist. Nat. 6: 19 (1824)                           |       |
|                       | <i>Sicyos angulata</i> L. = <b>Sicyos australis</b>   | 2:237 |
|                       | <b>Sicyos australis</b> Endl., Prodr. Fl. Norfolk 67 (1833)   | 2:237 |
| <b>CUNONIACEAE</b>    |   |       |
| e                     | <b>Anodopetalum biglandulosum</b> (A.Cunn. ex Hook.) Hook.f., Fl. Tasm. 1: 148 (1856)                   | 1:180 |
|                       | <b>Bauera rubioides</b> Andrews, Bot. Repos. 3: t.198 (1801)  | 1:180 |

## CUSCUTACEAE

- i **Cuscuta epithymum** (L.) Murray, Syst. Veg. ed.13, 140 (1774) 3:501  
i **Cuscuta suaveolens** Ser., Ann. Sci. Phys. Nat. Lyon 3: 519 (1840)  
**Cuscuta tasmanica** Engelm., Trans. Acad. Sci. St. Louis 1: 512 (1859) 3:501

## DILLENACEAE

- Hibbertia acicularis** (Labill.) F.Muell., Pl. Indig. Col. Vict. 1: 17 (1862) 1:23  
**Hibbertia appressa** Toelken, J. Adelaide Bot. Gard. 19: 54 (2000)  
*Hibbertia aspera* sensu Curtis & Morris (1975) = **Hibbertia hirticalyx** 1:23  
**Hibbertia calycina** (DC.) N.A.Wakef., Vict. Nat. 72: 122 (1955)  
**Hibbertia empetrifolia** (DC.) Hoogl., Kew Bull. 29: 155 (1974) 1:23  
    subsp. **empetrifolia**  
*Hibbertia fasciculata* sensu Curtis & Morris (1975) = **Hibbertia prostrata** 1:24  
e **Hibbertia hirsuta** (Hook.) Benth., Fl. Austral. 1: 26 (1863) 1:22  
**Hibbertia hirticalyx** Toelken, J. Adelaide Bot. Gard. 18(2): 146 (1998) 1:23  
**Hibbertia obtusifolia** DC., Syst. Nat. 1: 429 (1817) 1:24  
**Hibbertia procumbens** (Labill.) DC., Syst. Nat. 1: 427 (1817) 1:24  
**Hibbertia prostrata** Hook., J. Bot. (Hooker) 1: 246 (1834) 1:24  
**Hibbertia riparia** (R.Br. ex DC.) Hoogl., Kew Bull. 29: 155 (1974) 1:22  
**Hibbertia rufa** N.A.Wakef., Vict. Nat. 72: 119 (1955) 1:23  
**Hibbertia sericea** (R.Br. ex DC.) Benth., Fl. Austral. 1: 26 (1863) 1:22  
    var. **sericea**  
**Hibbertia serpyllifolia** R.Br. ex DC., Syst. Nat. 1: 430 (1817) 1:24  
**Hibbertia virgata** R.Br. ex DC., Syst. Nat. 1: 428 (1817) 1:24

## DIPSACACEAE

- i **Dipsacus fullonum** L., Sp. Pl. 1: 97 (1753) 2:280  
*Dipsacus sylvestris* Huds. = **Dipsacus fullonum** 2:280  
i **Scabiosa atropurpurea** L., Sp. Pl. 1: 100 (1753) 2:280

## DONATIACEAE

- t **Donatia novae-zelandiae** Hook.f., Fl. Nov.-Zel. 1: 81 (1852) 2:397

## DROSERACEAE

- Drosera arcturi** Hook., Lond. J. Bot. 1: 247 (1843) 1:187  
*Drosera auriculata* Backh. ex Planchon = **Drosera peltata** subsp. **auriculata** 1:188  
**Drosera binata** Labill., Nov. Holl. Pl. 1: 78 t.105 (1805) 1:188  
**Drosera glanduligera** Lehm., Nov. Strip. Pugill. 8: 37 (1844) 1:187  
*Drosera gracilis* Hook.f. ex Planchon = **Drosera peltata** 1:188  
**Drosera macrantha** Endl., Enum. Pl. Hügel. 6 (1837) 1:189  
    subsp. **planchonii** (Hook.f. ex Planchon) N.Marchant = **Drosera macrantha** 1:189  
**Drosera peltata** Thunb., Diss. Bot. Drosera 7 (1797) 1:188  
    subsp. **auriculata** (Backh. ex Planchon) Conn, J. Adelaide Bot. Gard. 3: 98 (1981) 1:188

|                       |  |       |
|-----------------------|--|-------|
|                       | subsp. <b>peltata</b>  |       |
|                       | <i>Drosera planchonii</i> Hook.f. = <b>Drosera macrantha</b>                                 | 1:189 |
|                       | <b>Drosera pygmaea</b> DC., Prodr. 1: 317 (1824)   | 1:187 |
|                       | <b>Drosera spatulata</b> Labill., Nov. Holl. Pl. 1: 79 t.106 fig.1 (1805)                    | 1:187 |
| <br>                  |  |       |
| <b>ELAEOCARPACEAE</b> |  |       |
| e                     | <b>Aristotelia peduncularis</b> (Labill.) Hook.f., Fl. Tasm. 1: 52 (1855)                    | 1:91  |
|                       | <b>Elaeocarpus reticulatus</b> Smith in Rees, Cycl. 12, Elaeocarpus no.6 (1809)              | 1:91  |
| <br>                  |  |       |
| <b>ELATINACEAE</b>    |  |       |
|                       | <b>Elatine gratioides</b> A.Cunn., Ann. Nat. Hist. 4: 26 (1840)                              | 1:80  |
| <br>                  |  |       |
| <b>EPACRIDACEAE</b>   |  |       |
| n                     | <b>Acrotriche affinis</b> DC., Prodr. 7(2): 757 (1839)                                       |       |
|                       | <b>Acrotriche cordata</b> (Labill.) R.Br., Prodr. 547 (1810)                                 |       |
|                       | <b>Acrotriche serrulata</b> R.Br., Prodr. 547 (1810)   | 2:440 |
| e                     | <b>Archeria comberi</b> Melville, Kew Bull. 3: 389 (1957)                                    | 2:453 |
| e                     | <b>Archeria eriocarpa</b> Hook.f., Fl. Tasm. 1: 263 t.80B (1857)                             | 2:452 |
| e                     | <b>Archeria hirtella</b> (Hook.f.) Hook.f., Fl. Tasm. 1: 263 t.81 (1857)                     | 2:452 |
| e                     | <b>Archeria serpyllifolia</b> Hook.f., Fl. Tasm. 1: 263 t.80B (1857)                         | 2:452 |
| e                     | var. <b>minor</b> Benth., Fl. Austral. 4: 246 (1868)   | 2:452 |
| e                     | var. <b>serpyllifolia</b>  |       |
|                       | <b>Astroloma humifusum</b> (Cav.) R.Br., Prodr. 538 (1810)                                   | 2:424 |
|                       | <b>Astroloma pinifolium</b> (R.Br.) Benth., Fl. Austral. 4: 159 (1868)                       | 2:424 |
|                       | <b>Brachyloma ciliatum</b> (R.Br.) Benth., Fl. Austral. 4: 173 (1868)                        | 2:432 |
|                       | <b>Brachyloma depressum</b> (F.Muell.) Benth., Fl. Austral. 4: 173 (1868)                    | 2:432 |
|                       | <i>Cyathodes abietina</i> (Labill.) R.Br. = <b>Leptecophylla abietina</b>                    | 2:427 |
| e                     | <b>Cyathodes dealbata</b> R.Br., Prodr. 539 (1810)   | 2:427 |
|                       | <i>Cyathodes divaricata</i> Hook.f. = <b>Leptecophylla divaricata</b>                        | 2:428 |
| e                     | <b>Cyathodes glauca</b> Labill., Nov. Holl. Pl. 1: 57 t.81 (1805)                            | 2:426 |
|                       | <i>Cyathodes juniperina</i> (Forst. & Forst.f.) Druce = <b>Leptecophylla juniperina</b>      | 2:427 |
|                       | <i>Cyathodes nitida</i> Jarman = <b>Planocarpa nitida</b>                                    |       |
|                       | <i>Cyathodes parvifolia</i> R.Br. = <b>Leptecophylla juniperina</b> subsp. <b>parvifolia</b> | 2:428 |
|                       | <i>Cyathodes pendulosa</i> Jarman = <b>Leptecophylla pendulosa</b>                           |       |
|                       | <i>Cyathodes petiolaris</i> (DC.) Druce = <b>Planocarpa petiolaris</b>                       | 2:426 |
| e                     | <b>Cyathodes platystoma</b> C.M.Weiller, Aust. Syst. Bot. 9(4): 502 (1996)                   |       |
| e                     | <b>Cyathodes straminea</b> R.Br., Prodr. 539 (1810)  | 2:426 |
|                       | <i>Cyathodes sulcata</i> Mihaich in Banks et al. = <b>Planocarpa sulcata</b>                 |       |
| e                     | <b>Dracophyllum milliganii</b> Hook.f., Icon. Pl. 9: t.845 (1852)                            | 2:461 |
| e                     | <b>Dracophyllum minimum</b> F.Muell., Fragm. 1: 39 (1859)                                    | 2:461 |
| e                     | <b>Epacris acuminata</b> Benth., Fl. Austral. 4: 240 (1868)                                  | 2:451 |
| e                     | <b>Epacris apsleyensis</b> Crowden, Pap. Proc. R. Soc. Tasm. 120: 17 (1986)                  |       |
| e                     | <b>Epacris barbata</b> Melville, Kew Bull. 176 (1952)  | 2:450 |
| e                     | <b>Epacris corymbiflora</b> Hook.f., Fl. Tasm. 1: 261 (1857)                                 | 2:447 |

|    |  |       |
|----|--|-------|
| e  | <b>Epacris curtisiae</b> Jarman, Pap. Proc. R. Soc. Tasm. 122: 115 (1988)                    |       |
| e  | <b>Epacris exserta</b> R.Br., Prodr. 551 (1810)  | 2:448 |
| ne | <b>Epacris franklinii</b> Hook.f., Fl. Tasm. 1: 261 (1857)                                   | 2:448 |
| e  | <b>Epacris glabella</b> Jarman in Banks et al., Aspects Tasm. Bot. 100 (1991)                |       |
| e  | <b>Epacris grandis</b> Crowden, Pap. Proc. R. Soc. Tasm. 120: 17 (1986)                      |       |
|    | <b>Epacris gunnii</b> Hook.f., Hook. Lond. J. Bot. 6: 272 (1847)                             | 2:451 |
| e  | <b>Epacris heteronema</b> Labill., Nov. Holl. Pl. 1: 42 t.56 (1805)                          | 2:449 |
|    | <b>Epacris impressa</b> Labill., Nov. Holl. Pl. 1: 43 t.58 (1805)                            | 2:446 |
|    | <b>Epacris lanuginosa</b> Labill., Nov. Holl. Pl. 1: 42 t.57 (1805)                          | 2:448 |
| e  | <b>Epacris limbata</b> K.J.Williams & F.Duncan in Banks et al., Aspects Tasm. Bot. 95 (1991) |       |
| e  | <b>Epacris marginata</b> Melville, Kew Bull. 175 (1952)                                      | 2:449 |
| e  | <b>Epacris mucronulata</b> R.Br., Prodr. 552 (1810)  | 2:448 |
| e  | <b>Epacris myrtifolia</b> Labill., Nov. Holl. Pl. 1: 41 t.55 (1805)                          | 2:447 |
| e  | <b>Epacris navicularis</b> Jarman, Pap. Proc. R. Soc. Tasm. 112: 3 (1978)                    |       |
|    | <b>Epacris obtusifolia</b> Smith, Exot. Bot. 1: 77 t.40 (1806)                               | 2:447 |
|    | <b>Epacris paludosa</b> R.Br., Prodr. 551 (1810)   | 2:449 |
|    | <b>Epacris petrophila</b> Hook.f., Fl. Tasm. 1: 261 (1857)                                   | 2:446 |
| e  | <b>Epacris serpyllifolia</b> R.Br., Prodr. 551 (1810)  | 2:450 |
|    | <i>Epacris squarrosa</i> Hook.f. = <b>Epacris tasmanica</b>                                  | 2:450 |
| e  | <b>Epacris stuartii</b> Stapf, Kew Bull. 7: 217 (1910)                                       | 2:449 |
| e  | <b>Epacris tasmanica</b> W.M.Curtis, Taxon 18: 244 (1969)                                    | 2:450 |
| e  | <b>Epacris virgata</b> Hook.f., Lond. J. Bot. 6: 271 (1847)                                  | 2:448 |
| e  | <b>Leptecophylla abietina</b> (Labill.) C.M.Weiller, Muelleria 12(2): 211 (1999)             | 2:427 |
| e  | <b>Leptecophylla divaricata</b> (Hook.f.) C.M.Weiller, Muelleria 12(2): 207 (1999)           | 2:428 |
|    | <b>Leptecophylla juniperina</b> (Forst. & Forst.f.) C.M.Weiller, Muelleria 12(2): 200 (1999) | 2:427 |
| t  | subsp. <b>juniperina</b>   | 2:427 |
|    | subsp. <b>oxycedrus</b> (Labill.) C.M.Weiller, Muelleria 12(2): 205 (1999)                   |       |
| e  | subsp. <b>parvifolia</b> (R.Br.) C.M.Weiller, Muelleria 12(2): 204 (1999)                    | 2:428 |
| e  | <b>Leptecophylla pendulosa</b> (Jarman) C.M.Weiller, Muelleria 12(2): 209 (1999)             |       |
| e  | <b>Leptecophylla pogonocalyx</b> C.M.Weiller, Muelleria 12(2): 206 (1999)                    |       |
|    | <b>Leucopogon australis</b> R.Br., Prodr. 541 (1810)   | 2:436 |
|    | <b>Leucopogon collinus</b> (Labill.) R.Br., Prodr. 543 (1810)                                | 2:436 |
|    | <b>Leucopogon ericoides</b> (Smith) R.Br., Prodr. 543 (1810)                                 | 2:438 |
|    | <b>Leucopogon esquamatus</b> R.Br., Prodr. 546 (1810)  | 2:439 |
|    | <b>Leucopogon hookeri</b> Sonder, Linnaea 26: 248 (1854)                                     | 2:438 |
|    | <b>Leucopogon lanceolatus</b> (Smith) R.Br., Prodr. 541 (1810)                               | 2:435 |
|    | var. <b>lanceolatus</b>  |       |
| e  | <b>Leucopogon milliganii</b> (F.Muell.) Rodway, Tasm. Fl. 118 (1903)                         | 2:439 |
|    | <b>Leucopogon montanus</b> (R.Br.) J.H.Willis, Vict. Nat. 73: 56 (1956)                      | 2:433 |
| e  | <b>Leucopogon oreophilus</b> J.M.Powell in Banks et al., Aspects Tasm. Bot. 108 (1991)       |       |
|    | <b>Leucopogon parviflorus</b> (Andrews) Lindley, Edwards's Bot. Reg. 18: t.1560 (1833)       | 2:435 |
|    | <b>Leucopogon pilifer</b> N.A.Wakef., Vict. Nat. 73: 58 (1956)                               |       |
|    | <b>Leucopogon stuartii</b> F.Muell. ex Sonder, Linnaea 26: 249 (1854)                        | 2:439 |
|    | <b>Leucopogon virgatus</b> (Labill.) R.Br., Prodr. 543 (1810)                                | 2:438 |



|   |   |       |
|---|---|-------|
|   | var. <b>brevifolius</b> Benth., Fl. Austral. 4: 202 (1869)                                |       |
|   | var. <b>virgatus</b>  |       |
|   | Lissanthe montana R.Br. = <b>Leucopogon montanus</b>                                      | 2:433 |
|   | <b>Lissanthe strigosa</b> (Smith) R.Br., Prodr. 540 (1810)                                | 2:433 |
|   | subsp. <b>subulata</b> (R.Br.) J.M.Powell, Telopea 5: 741 (1994)                          |       |
|   | <b>Monotoca elliptica</b> (Smith) R.Br., Prodr. 546 (1810)                                | 2:441 |
| e | <b>Monotoca empetrifolia</b> R.Br., Prodr. 547 (1810)                                     | 2:443 |
|   | <b>Monotoca glauca</b> (Labill.) Druce, Bot. Exch. Club Soc. Brit. Isles 1916: 636 (1917) | 2:442 |
| e | <b>Monotoca linifolia</b> (Rodway) W.M.Curtis, Stud. Fl. Tasm. 2: 464 (1963)              | 2:442 |
| e | subsp. <b>algida</b> Jarman in Banks et al., Aspects Tasm. Bot. 102 (1991)                |       |
| e | subsp. <b>linifolia</b>   |       |
|   | <b>Monotoca scoparia</b> (Smith) R.Br., Prodr. 547 (1810)                                 | 2:442 |
| e | <b>Monotoca submutica</b> (Benth.) Jarman, Pap. Proc. R. Soc. Tasm. 112: 1 (1978)         |       |
| e | var. <b>autumnalis</b> Jarman, Pap. Proc. R. Soc. Tasm. 112: 1 (1978)                     |       |
| e | var. <b>submutica</b>   |       |
| e | <b>Pentachondra ericifolia</b> Hook.f., Lond. J. Bot. 6: 271 (1847)                       | 2:430 |
| e | <b>Pentachondra involucrata</b> R.Br., Prodr. 549 (1810)                                  | 2:429 |
|   | <b>Pentachondra pumila</b> (Forst. & Forst.f.) R.Br., Prodr. 549 (1810)                   | 2:429 |
| e | <b>Planocarpa nitida</b> (Jarman) Weiller, Aust. Syst. Bot. 9(4): 517 (1996)              |       |
| e | <b>Planocarpa petiolaris</b> (DC.) Weiller, Aust. Syst. Bot. 9(4): 514 (1996)             | 2:426 |
| e | <b>Planocarpa sulcata</b> (Mihaich) Weiller, Aust. Syst. Bot. 9(4): 516 (1996)            |       |
| e | <b>Prionotes cerinthoides</b> (Labill.) R.Br., Prodr. 553 (1810)                          | 2:453 |
| e | <b>Richea acerosa</b> (Lindley) F.Muell., Fragm. 6: 69 (1868)                             | 2:458 |
| e | <b>Richea alpina</b> Y.Menadue, Aust. Syst. Bot. 13(5): 798 (2000)                        |       |
|   | Richea angustifolia B.L.Burtt = <b>Richea scoparia</b>                                    | 2:460 |
| e | <b>Richea Xcurtisiae</b> A.M.Gray, Muelleria 2: 143 (1971)                                |       |
| e | <b>Richea dracophylla</b> R.Br., Prodr. 555 (1810)  | 2:460 |
| e | <b>Richea gunnii</b> Hook.f., Lond. J. Bot. 6: 273 (1847)                                 | 2:459 |
| e | <b>Richea milliganii</b> (Hook.f.) F.Muell., Fragm. 6: 69 (1868)                          | 2:458 |
| e | <b>Richea pandanifolia</b> Hook.f., Fl. Antarct. 1: 50 (1844)                             | 2:460 |
| e | subsp. <b>pandanifolia</b>  | 2:460 |
| e | subsp. <b>ramulosa</b> Y.Menadue, Aust. Syst. Bot. 13(5): 795 (2000)                      |       |
| e | <b>Richea procera</b> (F.Muell.) F.Muell., Fragm. 6: 68 (1868)                            | 2:458 |
| e | <b>Richea scoparia</b> Hook.f., Lond. J. Bot. 6: 273 (1847)                               | 2:459 |
| e | <b>Richea sprengeloides</b> (R.Br.) F.Muell., Fragm. 6: 68 (1868)                         | 2:457 |
| e | <b>Sprengelia distichophylla</b> (Rodway) W.M.Curtis, Stud. Fl. Tasm. 2: 464 (1963)       | 2:456 |
|   | <b>Sprengelia incarnata</b> Smith, Tracts Nat. Hist. 272 t.2 (1798)                       | 2:454 |
|   | <b>Sprengelia montana</b> R.Br., Prodr. 555 (1810)  | 2:455 |
|   | <b>Styphelia adscendens</b> R.Br., Prodr. 537 (1810)                                      | 2:423 |
| e | <b>Trochocarpa cunninghamii</b> (DC.) W.M.Curtis, Stud. Fl. Tasm. 2: 463 (1963)           | 2:431 |
| e | <b>Trochocarpa disticha</b> (Labill.) Sprengel, Syst. 1: 660 (1824)                       | 2:430 |
| e | <b>Trochocarpa gunnii</b> (Hook.f.) Benth., Fl. Austral. 4: 167 (1868)                    | 2:431 |
| e | <b>Trochocarpa thymifolia</b> (R.Br.) Sprengel, Syst. 1: 660 (1824)                       | 2:431 |

## ERICACEAE

|   |  |       |
|---|--|-------|
| i | <b>Calluna vulgaris</b> (L.) Hull, Brit. Fl. ed.2, 1: 114 (1808)                         | 2:418 |
| i | <b>Erica andromedaeflora</b> Andrews, Coll. Heaths 3: t.146 (1805)                       | 2:419 |
| i | <b>Erica arborea</b> L., Sp. Pl. 1: 353 (1753)   | 2:420 |
| i | <b>Erica baccans</b> L., Mant. Pl. 2: 233 (1767)   | 2:419 |
| i | <b>Erica caffra</b> L., Sp. Pl. 1: 353 (1753)  | 2:421 |
| i | <b>Erica lusitanica</b> Rudolphi, J. Bot. (Schrader) 2: 286 (1800)                       | 2:420 |
| i | <b>Erica scoparia</b> L., Sp. Pl. 1: 353 (1753)  |       |
| i | <b>Erica Xwillmorei</b> Knowles & Westcott, Fl. Cab. 2: 115 (1838)                       | 2:420 |
| t | <b>Gaultheria depressa</b> Hook.f., Lond. J. Bot. 6: 267 (1847)                          | 2:417 |
|   | <b>Gaultheria hispida</b> R.Br., Prodr. 558 (1810)                                       | 2:417 |
| e | <b>Gaultheria lanceolata</b> Hook.f., Lond. J. Bot. 6: 267 (1847)                        | 2:416 |
| e | <b>Gaultheria tasmanica</b> (Hook.f.) Middleton, Edinb. J. Bot. 47: 299 (1990)           | 2:416 |
|   | <i>Pernettya lanceolata</i> (Hook.f.) B.L.Burt & A.W.Hill = <b>Gaultheria lanceolata</b> | 2:416 |
|   | <i>Pernettya tasmanica</i> Hook.f. = <b>Gaultheria tasmanica</b>                         | 2:416 |

## ESCALLONIACEAE (GROSSULARIACEAE)

|   |   |       |
|---|---|-------|
| e | <b>Anopterus glandulosus</b> Labill., Nov. Holl. Pl. 1: 86 t.112 (1805) | 1:181 |
| e | <b>Tetracarpaea tasmanica</b> Hook.f., Icon. Pl. 3: t.264 (1840)        | 1:182 |

## EUCRYPHIACEAE

|   |   |       |
|---|---|-------|
| e | <b>Eucryphia lucida</b> (Labill.) Baill., Hist. Pl. 1: 401 (1869)                                   | 1:183 |
| e | <b>Eucryphia milliganii</b> Hook.f., Fl. Tasm. 1: 54 (1855)   | 1:183 |
| e | subsp. <b>milliganii</b>  | 1:183 |
| e | subsp. <b>pubescens</b> R.W.Barnes, G.J.Jord., R.S.Hill & C.J.McCoull, Aust. J. Bot. 48: 488 (2000) |       |

## EUPHORBIACEAE

|   |  |       |
|---|--|-------|
|   | <b>Amperea xiphoclada</b> (Sieber ex Sprengel) Druce, Bot. Exch. Club Brit. Isles 1916: 604 (1917)     | 3:637 |
|   | <i>Bertya rosmarinifolia</i> sensu W.M.Curtis (1967) = <b>Bertya tasmanica</b> subsp. <b>tasmanica</b> | 3:637 |
|   | <b>Bertya tasmanica</b> (Sond. & F.Muell.) Müll.Arg., Linnaea 34: 63 (1865)                            |       |
| e | subsp. <b>tasmanica</b>  | 3:637 |
|   | <b>Beyeria lechenaultii</b> (DC.) Baill., Adansonia 6: 307 (1866)                                      | 3:635 |
|   | var. <b>latifolia</b> Grüning, Pflanzenr. IV 147 (Heft 58): 71 (1913)                                  | 3:635 |
|   | <b>Beyeria viscosa</b> (Labill.) Miq., Ann. Sci. Nat. Bot. ser.3 1: 350 t.15 (1844)                    | 3:635 |
|   | <b>Chamaesyce drummondii</b> (Boiss.) D.Hassall, Austral. J. Bot. 24: 640 (1976)                       | 3:632 |
| i | <b>Euphorbia cyparissias</b> L., Sp. Pl. 1: 461 (1753)   | 3:633 |
|   | <i>Euphorbia drummondii</i> Boiss. = <b>Chamaesyce drummondii</b>                                      | 3:632 |
| i | <b>Euphorbia exigua</b> L., Sp. Pl. 1: 456 (1753)  |       |
| i | <b>Euphorbia helioscopia</b> L., Sp. Pl. 1: 459 (1753)   | 3:632 |
| i | <b>Euphorbia lathyris</b> L., Sp. Pl. 1: 457 (1753)  | 3:632 |
| i | <b>Euphorbia paralias</b> L., Sp. Pl. 1: 458 (1753)  | 3:633 |
| i | <b>Euphorbia peplus</b> L., Sp. Pl. 1: 456 (1753)  | 3:633 |
|   | <b>Micrantheum hexandrum</b> Hook.f., Lond. J. Bot. 6: 283 (1847)                                      | 3:634 |
| e | <b>Micrantheum serpentinum</b> Orch. in Banks et al., Aspects Tasm. Bot. 60 (1991)                     |       |

|                               |   |       |
|-------------------------------|---|-------|
| e                             | <b>Oreopranthera petalifera</b> Orch. & J.Davies, Pap. Proc. R. Soc. Tasm. 119: 62 (1985)                     |       |
|                               | <b>Phyllanthus australis</b> Hook.f., Lond. J. Bot. 6: 284 (1847)   | 3:638 |
|                               | <b>Phyllanthus gunnii</b> Hook.f., Lond. J. Bot. 6: 284 (1847)  | 3:638 |
|                               | <b>Poranthera microphylla</b> Brongn., Ann. Sci. Nat. 29: 385 (1833)  | 3:634 |
|                               | <i>Pseudanthus divaricatissimus</i> (Muell. Arg.) Benth., not present in Tasmania, formerly recorded in error |       |
|                               | <b>Pseudanthus ovalifolius</b> F.Muell., Trans. Phil. Inst. Vict. 2: 66 (1858)                                | 3:634 |
|                               | <i>Pseudanthus tasmanicus</i> Rodway = <b>Muehlenbeckia axillaris</b> (Polygonaceae)                          | 3:593 |
|                               | <b>Ricinocarpos pinifolius</b> Desf., Mem. Mus. Hist. Nat. Paris 3: 459 t.22 (1817)                           | 3:636 |
| <br>                          |   |       |
| <b>FABACEAE (LEGUMINOSAE)</b> |   |       |
|                               | <b>Almaleea subumbellata</b> (Hook.) Crisp & P.Weston, Telopea 4(2): 310 (1991)                               | 1:141 |
|                               | <b>Aotus ericoides</b> (Vent.) G.Don, Gen. Syst. 2: 120 (1832)  | 1:137 |
|                               | <i>Aotus villosa</i> (Andrews) Smith = <b>Aotus ericoides</b>   | 1:137 |
|                               | <b>Bossiaea cinerea</b> R.Br. in W.T.Aiton, Hortus Kew. ed.2, 4: 268 (1812)                                   | 1:147 |
|                               | <b>Bossiaea cordigera</b> Benth. ex Hook.f., Fl. Tasm. 1: t.16 (1855), 95 (1856)                              | 1:147 |
|                               | <i>Bossiaea ensata</i> Sieber ex DC., not present in Tasmania, formerly recorded in error                     |       |
|                               | <b>Bossiaea obcordata</b> (Vent.) Druce, Bot. Exch. Club Soc. Brit. Isles 1916: 610 (1917)                    | 1:148 |
|                               | <b>Bossiaea prostrata</b> R.Br., Bot. Mag. 36: t.1493 (1812)  | 1:147 |
|                               | <b>Bossiaea riparia</b> A.Cunn. ex Benth., Fl. Austral. 2: 166 (1864)   | 1:148 |
| i                             | <b>Chamaecytisus palmensis</b> (Christ) F.A.Bisby & K.W.Nicholls, J. Linn. Soc. Bot. 74: 114 (1977)           | 1:151 |
|                               | <i>Chamaecytisus proliferus</i> sensu Curtis & Morris (1975) = <b>Chamaecytisus palmensis</b>                 | 1:151 |
|                               | <i>Coronilla varia</i> L. = <b>Securigera varia</b>   | 1:165 |
|                               | <b>Cullen microcephalum</b> (Rechb. ex Kunze) J.W.Grimes, Muellera 9: 195 (1996)                              | 1:163 |
|                               | <i>Cytisus palmensis</i> (Christ) Hutch. = <b>Chamaecytisus palmensis</b>                                     | 1:151 |
| i                             | <b>Cytisus scoparius</b> (L.) Link, Enum. Hort. Berol. Alt. 2: 241 (1822)                                     | 1:151 |
|                               | <b>Daviesia latifolia</b> R.Br. in W.T.Aiton, Hortus Kew. ed.2, 3: 20 (1811)                                  | 1:136 |
|                               | <b>Daviesia sejugata</b> G.Chandler & Crisp, Aust. Syst. Bot. 10(1): 33 (1997)                                |       |
|                               | <b>Daviesia ulicifolia</b> Andrews, Bot. Repos. 5: t.304 (1803)   | 1:136 |
|                               | subsp. <b>ruscifolia</b> (A.Cunn. ex Benth.) G.Chandler & Crisp, Aust. Syst. Bot. 10(1): 44 (1997)            |       |
|                               | subsp. <b>ulicifolia</b>  |       |
|                               | <b>Desmodium gunnii</b> Benth. ex Hook.f., Fl. Tasm. 1: 101 (1856)  | 1:166 |
| n                             | <b>Desmodium varians</b> (Labill.) G.Don, Gen. Hist. 2: 298 (1832)  | 1:166 |
|                               | <b>Dillwynia cinerascens</b> R.Br. ex Sims, Bot. Mag. 48: t.2247 (1821)                                       | 1:145 |
|                               | <b>Dillwynia glaberrima</b> Smith, Ann. Bot. (Koenig & Sims) 1: 510 (1805)                                    | 1:144 |
|                               | <b>Dillwynia sericea</b> A.Cunn. in Field, Geogr. Mem. N.S.W. 347 (1825)                                      | 1:144 |
| i                             | <b>Dipogon lignosus</b> (L.) Verdc., Taxon 17: 537 (1968)   |       |
|                               | <i>Dolichos lignosus</i> L. = <b>Dipogon lignosus</b>   |       |
|                               | <b>Eutaxia microphylla</b> (R.Br.) C.H.Wright & Dewar, Johnson's Gard. Dict. 1052 (1894)                      | 1:143 |
|                               | <i>Genista maderensis</i> sensu Curtis & Morris (1975) = <b>Genista stenopetala</b>                           | 1:152 |
| i                             | <b>Genista monspessulana</b> (L.) L.A.S.Johnson, Contr. N.S.W. Nat. Herb. 3: 98 (1962)                        | 1:151 |
| i                             | <b>Genista stenopetala</b> Webb & Berthel., Phyt. Canar. 3(2): t.45 (1836)                                    | 1:152 |
|                               | <b>Glycine clandestina</b> J.Wendl., Bot. Beobacht. 54 (1798)   | 1:168 |
|                               | <b>Glycine latrobeana</b> (Meisn.) Benth., Fl. Austral. 2: 244 (1864)   | 1:168 |
|                               | <b>Glycine microphylla</b> (Benth.) Tind., Brunonia 9: 181 (1987)   |       |

|    |   |       |
|----|---|-------|
|    | <b>Gompholobium ecostatum</b> Kuchel in H.Eichler, Suppl. Black's Fl. S.Austral. 182 (1965)   | 1:135 |
|    | <b>Gompholobium huegelii</b> Benth. in Endl. et al., Enum. Pl. 29 (1837)                      | 1:135 |
|    | <b>Goodia lotifolia</b> Salisb., Parad. Londin. 1: t.41 (1806)                                | 1:149 |
|    | var. <b>lotifolia</b>   | 1:149 |
|    | var. <b>pubescens</b> (Sims) Williamson in Ewart, Fl. Vict. 658 (1931)                        | 1:149 |
|    | Goodia pubescens Sims = <b>Goodia lotifolia</b> var. <b>pubescens</b>                         | 1:149 |
|    | <b>Hardenbergia violacea</b> (Schneev.) Stearn, J. Bot. 78: 70 (1940)                         | 1:169 |
| i# | <b>Hedysarum coronarium</b> L., Sp. Pl. 2: 750 (1753)   | 1:165 |
|    | <b>Hovea corrickiae</b> J.H.Ross, Muelleria 7: 203 (1990)                                     |       |
|    | <b>Hovea heterophylla</b> A.Cunn. ex Hook.f., Fl. Tasm. 1: 93 (1856)                          | 1:148 |
|    | Hovea lanceolata sensu Census 1999 = <b>Hovea tasmanica</b>                                   |       |
|    | Hovea linearis sensu Census 1999 = <b>Hovea heterophylla</b>                                  |       |
|    | Hovea longifolia sensu Curtis & Morris (1975) = <b>Hovea montana</b> & <b>Hovea tasmanica</b> | 1:149 |
|    | <b>Hovea magnibractea</b> I.Thomps., Aust. Syst. Bot. 14(1): 74 (2001)                        |       |
|    | <b>Hovea montana</b> (Hook.f.) J.H.Ross, Muelleria 6: 427 (1988)                              | 1:149 |
| e  | <b>Hovea tasmanica</b> I.Thomps. & J.H.Ross, Aust. Syst. Bot. 14(1): 68 (2001)                |       |
|    | <b>Indigofera australis</b> Willd., Sp. Pl. ed.5, 3: 1235 (1802)                              | 1:164 |
| i  | <b>Kennedia nigricans</b> Lindley, Edwards's Botanical Register 20: 1715, pl. 1715 (1835)     |       |
|    | <b>Kennedia prostrata</b> R.Br. in W.T.Aiton, Hortus Kew. ed.2, 4: 299 (1812)                 | 1:169 |
| i  | <b>Kennedia rubicunda</b> (Schneev.) Vent., Jard. Malmaison 2: 104 t.104 (1804)               |       |
| i? | <b>Lathyrus nissolia</b> L., Sp. Pl. 2: 729 (1753)  | 1:168 |
| i  | <b>Lathyrus tingitanus</b> L., Sp. Pl. 2: 732 (1753)  |       |
| i# | <b>Lotus angustissimus</b> L., Sp. Pl. 774 (1753)   |       |
|    | <b>Lotus australis</b> Andrews, Bot. Repos. 10 t.624 (1811)                                   | 1:162 |
| i  | <b>Lotus corniculatus</b> L., Sp. Pl. 2: 775 (1753)   | 1:162 |
| i  | var. <b>corniculatus</b>  |       |
| i  | var. <b>tenuifolius</b> L., Sp. Pl. 2: 776 (1753)   |       |
|    | Lotus hispidus Desf. ex DC. = <b>Lotus suaveolens</b>   | 1:163 |
|    | Lotus pedunculatus sensu Curtis & Morris (1975) = <b>Lotus uliginosus</b>                     | 1:163 |
| i  | <b>Lotus suaveolens</b> Pers., Syn. Pl. 2: 354 (1807)   | 1:163 |
|    | Lotus subbiflorus sensu Census Vasc. Pl. Tasm. edn.1 = <b>Lotus suaveolens</b>                | 1:163 |
|    | Lotus tenuis Waldst. & Kit. = <b>Lotus corniculatus</b> var. <b>tenuifolius</b>               |       |
| i  | <b>Lotus uliginosus</b> Schkuhr, Handb. 2: 412 1796   | 1:163 |
| i  | <b>Lupinus angustifolius</b> L., Sp. Pl. 2: 721 (1753)  | 1:150 |
| i  | <b>Lupinus arboreus</b> Sims, Bot. Mag. 18: t.682 (1803)                                      | 1:150 |
| i  | <b>Medicago arabica</b> (L.) Huds., Fl. Angl. 288 (1762)                                      | 1:154 |
| i# | <b>Medicago arborea</b> L., Sp. Pl. 2: 778 (1753)   |       |
| i# | <b>Medicago falcata</b> L., Sp. Pl. 2: 779 (1753)   | 1:153 |
| i  | <b>Medicago lupulina</b> L., Sp. Pl. 2: 779 (1753)  | 1:153 |
| i  | <b>Medicago minima</b> (L.) Bartal., Cat. Piante Siena 61 (1776)                              | 1:154 |
| i  | <b>Medicago polymorpha</b> L., Sp. Pl. 2: 779 (1753)  | 1:154 |
| i  | <b>Medicago sativa</b> L., Sp. Pl. 2: 778 (1753)  | 1:153 |
| i  | <b>Medicago scutellata</b> (L.) Mill., Gard. Dict. edn.8 no.2 (1768)                          | 1:154 |
| i  | <b>Melilotus albus</b> Medik., Varies. Churpfalz. Phys.-Ocon. Ges. 2: 382 (1787)              | 1:155 |

|   |   |       |
|---|---|-------|
| i | <b>Melilotus indicus</b> (L.) All., Fl. Pedem. 1: 308 (1785)                            | 1:155 |
| i | <b>Melilotus officinalis</b> (L.) Pallas, Reise 3: 537 (1776)                           | 1:155 |
|   | <b>Mirbelia oxylobioides</b> F.Muell., Fragm. 2: 154 (1861)                             |       |
| i | <b>Onobrychis viciifolia</b> Scop., Fl. Carn. edn.2, 2: 76 (1772)                       | 1:165 |
|   | <i>Ononis arvensis</i> sensu L.Rodway (1903) = <b>Ononis spinosa</b>                    | 1:152 |
|   | <i>Ononis repens</i> sensu Curtis & Morris (1975) = <b>Ononis spinosa</b>               | 1:152 |
| i | <b>Ononis spinosa</b> L., Sp. Pl. 2: 716 (1753)   | 1:152 |
| i | <b>Ornithopus compressus</b> L., Sp. Pl. 2: 744 (1753)                                  |       |
| i | <b>Ornithopus pinnatus</b> (Miller) Druce, J. Bot. 45: 420 (1907)                       |       |
|   | <b>Oxylobium arborescens</b> R.Br. in W.T.Aiton, Hortus Kew. edn.2, 3: 10 (1811)        | 1:134 |
|   | <b>Oxylobium ellipticum</b> (Vent.) R.Br. in W.T.Aiton, Hortus Kew. edn.2, 3: 10 (1811) | 1:134 |
| e | <b>Phyllota diffusa</b> (Hook.f.) F.Muell., Fragm. 1: 8 (1858)                          | 1:137 |
|   | <b>Platylobium formosum</b> Smith, Spec. Bot. New Holl. 1: 17, t.6 (1793)               | 1:146 |
|   | subsp. <i>parviflorum</i> (Smith) A.Lee, deleted from the flora (Ross 1983)             |       |
|   | <b>Platylobium obtusangulum</b> Hook., Bot. Mag. 60: t.3258 (1833)                      | 1:146 |
|   | <b>Platylobium triangulare</b> R.Br. in W.T.Aiton, Hortus Kew. edn.2, 4: 266 (1812)     | 1:146 |
| i | <b>Podalyria sericea</b> R.Br. in W.T.Aiton, Hortus Kew. edn.2, 3: 6 (1811)             |       |
|   | <i>Psoralea adscendens</i> F.Muell. = <b>Cullen microcephalum</b>                       | 1:163 |
| i | <b>Psoralea pinnata</b> L., Sp. Pl. 2: 762 (1753)                                       | 1:163 |
|   | <b>Pultenaea daphnoides</b> J.Wendl., Bot. Bosbacht. 49 (1798)                          | 1:139 |
|   | var. <b>obcordata</b> (Andrews) Hook.f., Fl. Tasm. 1(2): 86 (1856)                      | 1:139 |
|   | <b>Pultenaea dentata</b> Labill., Nov. Holl. Pl. 1: 103 t.131 (1806)                    | 1:141 |
|   | <b>Pultenaea fasciculata</b> Benth., Ann. Wien. Mus. Naturg. 2: 82 (1840)               | 1:142 |
|   | <b>Pultenaea gunnii</b> Benth., Comm. Legum. Gen. 18 (1837)                             | 1:140 |
| e | var. <b>baeckeoides</b> (A.Cunn. ex Benth.) Rodway, Tasm. Fl. 33 (1903)                 | 1:140 |
|   | var. <b>gunnii</b>  | 1:140 |
|   | <i>Pultenaea hibbertioides</i> Hook.f. = <b>Pultenaea mollis</b>                        | 1:141 |
|   | <b>Pultenaea humilis</b> Benth. ex Hook.f., Fl. Tasm. 1: 91 (1856)                      | 1:142 |
|   | <b>Pultenaea juniperina</b> Labill., Nov. Holl. Pl. 1: 102 t.130 (1806)                 | 1:142 |
|   | <b>Pultenaea mollis</b> Lindley in T.Mitch., Three Exped. Australia 2: 258 (1838)       | 1:141 |
|   | <i>Pultenaea paleacea</i> var. <i>sericea</i> = <b>Pultenaea sericea</b>                | 1:140 |
|   | <b>Pultenaea pedunculata</b> Hook., Bot. Mag. 55: t.2859 (1828)                         | 1:140 |
|   | <b>Pultenaea prostrata</b> Benth. ex Hook.f., Fl. Tasm. 1: 89 (1856)                    | 1:142 |
|   | <i>Pultenaea selaginoides</i> Hook.f. = <b>Stonesiella selaginoides</b>                 | 1:141 |
|   | <b>Pultenaea sericea</b> (Benth.) Corrick, Muelleria 8(3): 392 (1995)                   | 1:140 |
|   | <b>Pultenaea stricta</b> Sims, Bot. Mag. 38: t.1588 (1813)                              | 1:140 |
|   | <i>Pultenaea subumbellata</i> Hook. = <b>Almaleea subumbellata</b>                      | 1:141 |
|   | <b>Pultenaea tenuifolia</b> R.Br. ex Sims, Bot. Mag. 46: t.2086 (1819)                  | 1:143 |
|   | <i>Sarothamnus scoparius</i> (L.) Wimmer ex Koch = <b>Cytisus scoparius</b>             | 1:151 |
| i | <b>Securigera varia</b> (L.) Lassen, Svensk. Bot. Tidskr. 83: 86 (1989)                 | 1:165 |
| i | <b>Spartium junceum</b> L., Sp. Pl. 2: 708 (1753)                                       |       |
|   | <b>Sphaerolobium minus</b> Labill., Nov. Holl. Pl. 1: 108, t.138 (1805)                 |       |
|   | <b>Sphaerolobium vimineum</b> Sm., Ann. Bot. (Konig & Sims) 1: 509 (1805)               | 1:135 |
| e | <b>Stonesiella selaginoides</b> (Hook.f.) Crisp & P.H.Weston, Taxon 48(4): 711 (1999)   | 1:141 |

|                                     |   |       |
|-------------------------------------|---|-------|
|                                     | <b>Swainsona lessertiifolia</b> DC., Ann. Sci. Nat. Bot. ser.1, 4: 99 (1825)                              | 1:164 |
| i                                   | <b>Trifolium alexandrinum</b> L., Cent. Pl. 1: 25 (1755)  |       |
| i                                   | <b>Trifolium angustifolium</b> L., Sp. Pl. 2: 769 (1753)  | 1:161 |
| i                                   | <b>Trifolium arvense</b> L., Sp. Pl. 2: 769 (1753)  | 1:160 |
| i                                   | <b>Trifolium campestre</b> Schreber in Sturm, Deutschl. Fl. Abt. 1 (Heft 16): t.253 (1804)                | 1:157 |
| i                                   | <b>Trifolium cernuum</b> Brot., Phytogr. Lusit. Select. 1: 150 (1816)                                     | 1:159 |
| i                                   | <b>Trifolium dubium</b> Sibth., Fl. Oxon. 231 (1794)  | 1:157 |
| i                                   | <b>Trifolium fragiferum</b> L., Sp. Pl. 2: 772 (1753)   | 1:158 |
| i                                   | <b>Trifolium glomeratum</b> L., Sp. Pl. 2: 770 (1753)   | 1:159 |
| i                                   | <b>Trifolium hybridum</b> L., Sp. Pl. 2: 766 (1753)   | 1:159 |
| i                                   | <b>Trifolium incarnatum</b> L., Sp. Pl. 2: 769 (1753)   | 1:160 |
| i                                   | <b>Trifolium lappaceum</b> L., Sp. Pl. 2: 768 (1753)  |       |
|                                     | <i>Trifolium medium</i> Huds. Not present in Tas. This name was based on a specimen of <i>T. pratense</i> | 1:159 |
| i                                   | <b>Trifolium micranthum</b> Viv., Fl. Lig. 45 (1824)  | 1:158 |
| i                                   | <b>Trifolium ornithopodioides</b> L., Sp. Pl. 2: 766 (1753)   | 1:152 |
| i                                   | <b>Trifolium pratense</b> L., Sp. Pl. 2: 768 (1753)   | 1:160 |
| i                                   | <b>Trifolium repens</b> L., Sp. Pl. 2: 767 (1753)   | 1:158 |
| i                                   | <b>Trifolium resupinatum</b> L., Sp. Pl. 2: 771 (1753)  | 1:158 |
| i                                   | <b>Trifolium scabrum</b> L., Sp. Pl. 2: 770 (1753)  | 1:160 |
| i                                   | <b>Trifolium squamosum</b> L., Amoen. Acad. 4: 105 (1759)   | 1:161 |
| i                                   | <b>Trifolium stellatum</b> L., Sp. Pl. 2: 769 (1753)  | 1:161 |
| i                                   | <b>Trifolium striatum</b> L., Sp. Pl. 2: 770 (1753)   | 1:160 |
| i                                   | <b>Trifolium subterraneum</b> L., Sp. Pl. 2: 767 (1753)   | 1:159 |
| i                                   | <b>Trifolium suffocatum</b> L., Mant. Pl. 2: 276 (1771)   | 1:159 |
| i                                   | <b>Trifolium tomentosum</b> L., Sp. Pl. 2: 771 (1753)   | 1:158 |
| i                                   | <b>Trifolium uniflorum</b> L., Sp. Pl. 2: 771 (1753)  |       |
|                                     | <i>Trigonella ornithopodioides</i> (L.) DC. = <b>Trifolium ornithopodioides</b>                           | 1:152 |
| i                                   | <b>Ulex europaeus</b> L., Sp. Pl. 2: 741 (1753)   | 1:150 |
|                                     | <i>Vicia angustifolia</i> L. = <b>Vicia sativa</b>  | 1:167 |
|                                     | <i>Vicia cracca</i> sensu Curtis & Morris (1975) = <b>Vicia villosa</b>                                   | 1:167 |
| i                                   | <b>Vicia hirsuta</b> (L.) Gray, Nat. Arr. Brit. Pl. 2: 614 (1821)   | 1:166 |
|                                     | <i>Vicia lathyroides</i> L. Not present in Tasmania, formerly recorded in error                           |       |
| i                                   | <b>Vicia sativa</b> L., Sp. Pl. 2: 736 (1753)   | 1:167 |
| i                                   | subsp. <b>nigra</b> (L.) Ehrh., Hannover. Mag. 1780(15): 229 (1780)                                       |       |
| i                                   | <b>Vicia tetrasperma</b> (L.) Schreber, Spicil. Fl. Lips. 26 (1771)                                       | 1:167 |
| i                                   | <b>Vicia villosa</b> Roth, Tent. Fl. Germ. 2(2): 182 (1793)   | 1:167 |
| i                                   | subsp. <b>eriocarpa</b> (Hauskn.) P.W.Ball, Feddes Repert. 79: 45 (1968)                                  |       |
|                                     | <b>Viminaria juncea</b> (Schrader & J.Wendl.) Hoffsgg., Verz. Pfl.-Kult. 200 (1824)                       | 1:136 |
| <br><b>FAGACEAE (NOTHOFAGACEAE)</b> |   |       |
|                                     | <b>Nothofagus cunninghamii</b> (Hook.) Oersted, Vidensk. Sel. Skr. ser.5, 9: 355 (1871)                   | 3:647 |
| e                                   | <b>Nothofagus gunnii</b> (Hook.f.) Oersted, Vidensk. Sel. Skr. ser.5, 9: 354 (1871)                       | 3:646 |
| <br><b>FRANKENIACEAE</b>            |   |       |

|                     |  |       |
|---------------------|--|-------|
|                     | <b>Frankenia pauciflora</b> DC., Prodr. 1: 350 (1824)  | 1:64  |
|                     | var. <b>gunnii</b> Summerh., J. Linn. Soc. Bot. 48: 366 (1930)   |       |
| <b>FUMARIACEAE</b>  |  |       |
| i                   | <b>Fumaria bastardii</b> Boreau in Duchartre, Rev. Bot. 2: 359 (1847)                                    | 1:30  |
| i                   | <b>Fumaria densiflora</b> DC., Cat. Pl. Horti Monsp. 113 (1813)  | 1:30  |
| i                   | <b>Fumaria muralis</b> Sonder ex Koch, Syn. Fl. Germ. ed.2, 1017 (1845)                                  | 1:30  |
| i                   | <b>Fumaria officinalis</b> L., Sp. Pl. 2: 700 (1753)   |       |
| i#                  | <b>Pseudofumaria alba</b> (Miller) Lidén, Opera Bot. 88: 32 (1986)                                       |       |
|                     | subsp. <b>alba</b>   |       |
| <b>GENTIANACEAE</b> |  |       |
|                     | <i>Centaureum australe</i> (R.Br.) Druce = <b>Centaureum spicatum</b>                                    | 3:479 |
| i                   | <b>Centaureum erythraea</b> Rafn, Danm. Holst. Fl. 2: 75 (1800)  | 3:480 |
|                     | <i>Centaureum pulchellum</i> sensu W.M.Curtis (1967) = <b>Centaureum tenuiflorum</b>                     | 3:480 |
| ?i                  | <b>Centaureum spicatum</b> (L.) Fritsch ex Janch., Mitt. Naturw. Ver., Wien 5: 97 (1907)                 | 3:479 |
| i                   | <b>Centaureum tenuiflorum</b> (Hoffsgg. & Link) Fritsch ex Janch., Mitt. Naturw. Ver., Wien 5: 97 (1907) | 3:480 |
| e                   | <b>Chionogentias brevisepala</b> L.G.Adams, Aust. Syst. Bot. 8(5): 967 (1995)                            | 3:482 |
| x                   | <b>Chionogentias cunninghamii</b> L.G.Adams, Aust. Syst. Bot. 8(5): 962 (1995)                           | 3:482 |
| x                   | subsp. <b>cunninghamii</b>   |       |
| e                   | <b>Chionogentias demissa</b> L.G.Adams, Aust. Syst. Bot. 8(5): 960 (1995)                                | 3:482 |
| e                   | <b>Chionogentias diemensis</b> (Griseb.) L.G.Adams, Aust. Syst. Bot. 8(5): 952 (1995)                    | 3:482 |
| e                   | subsp. <b>diemensis</b>  |       |
| e                   | subsp. <b>plantaginea</b> L.G.Adams, Aust. Syst. Bot. 8(5): 954 (1995)                                   |       |
| e                   | <b>Chionogentias eichleri</b> L.G.Adams, Aust. Syst. Bot. 8(5): 973 (1995)                               | 3:482 |
| e                   | <b>Chionogentias grandis</b> L.G.Adams, Aust. Syst. Bot. 8(5): 960 (1995)                                | 3:482 |
| e                   | <b>Chionogentias gunniana</b> L.G.Adams, Aust. Syst. Bot. 8(5): 969 (1995)                               | 3:482 |
| e                   | <b>Chionogentias pleurogynoides</b> (Griseb.) L.G.Adams, Aust. Syst. Bot. 8(5): 965 (1995)               | 3:482 |
| e                   | subsp. <b>pleurogynoides</b>   |       |
| e                   | subsp. <b>milliganii</b> L.G.Adams, Aust. Syst. Bot. 8(5): 967 (1995)                                    | 3:482 |
|                     | <b>Chionogentias polysperes</b> L.G.Adams, Aust. Syst. Bot. 8(5): 970 (1995)                             | 3:482 |
| i                   | <b>Cicendia filiformis</b> (L.) Delarbre, Fl. Auvergne 1: 20 (1795)                                      | 3:481 |
|                     | <i>Gentianella diemensis</i> (Griseb.) J.H.Willis = <b>Chionogentias</b> spp.                            | 3:482 |
|                     | <b>Sebaea albidiflora</b> F.Muell., Trans. Phil. Soc. Vict. 1: 46 (1855)                                 | 3:479 |
|                     | <b>Sebaea ovata</b> (Labill.) R.Br., Prodr. 452 (1810)   | 3:478 |
| <b>GERANIACEAE</b>  |  |       |
| i                   | <b>Erodium botrys</b> (Cav.) Bertol., Amoen. Ital. 35 (1819)   | 1:98  |
| i                   | <b>Erodium cicutarium</b> (L.) L'Hérit. in Aiton, Hort. Kew. 2: 414 (1789)                               | 1:97  |
| i                   | <b>Erodium malacoides</b> (L.) Willd., Phytograph. 1: 10 (1794)  | 1:98  |
| i                   | <b>Erodium moschatum</b> (L.) L'Hérit. in Aiton, Hort. Kew. 2: 414 (1789)                                | 1:97  |
| i                   | <b>Geranium dissectum</b> L., Cent. Pl. 1: 21 (1755)   | 1:96  |
| i                   | <b>Geranium molle</b> L., Sp. Pl. 2: 682 (1753)  | 1:96  |
|                     | <b>Geranium potentilloides</b> L'Hérit. ex DC., Prodr. 1: 639 (1824)                                     | 1:95  |

|    |  |      |
|----|--|------|
|    | <b>Geranium retrorsum</b> L'Hérit. ex DC., Prodr. 1: 644 (1824)                  |      |
| it | <b>Geranium rotundifolium</b> L., Sp. Pl. 2: 683 (1753)                          | 1:96 |
|    | <b>Geranium sessiliflorum</b> Cav., Diss. 4: 198, t.77 fig.2 (1787)              | 1:95 |
|    | subsp. <b>brevicaule</b> (Hook.) Carolin, Proc. Linn. Soc. N.S.W. 89: 357 (1964) | 1:95 |
|    | <b>Geranium solanderi</b> Carolin, Proc. Linn. Soc. N.S.W. 89: 350 (1964)        | 1:95 |
| i# | <b>Geranium yeoi</b> Aedo & Muñoz Garm., Kew Bull. 52: 727 (1997)                |      |
|    | <b>Pelargonium australe</b> Willd., Sp. Pl. ed.5, 3: 675 (1800)                  | 1:98 |
|    | <b>Pelargonium inodorum</b> Willd., Enum. Hort. Bot. Berol. Alt. 702 (1809)      | 1:99 |
|    | <b>Pelargonium littorale</b> Huegel, Bot. Arch. t.5 (1837)                       |      |
| i  | <b>Pelargonium Xasperum</b> Ehrh. ex Willd., Sp. Pl. ed.4, 3: 678 (1800)         | 1:99 |
| i  | <b>Pelargonium Xdomesticum</b> L.H.Bailey, Stand. Cycl. Hort. 2532 (1916)        | 1:99 |

#### GOODENIACEAE

|   |   |       |
|---|---|-------|
| x | <b>Cooperhooikia barbata</b> (R.Br.) Carolin, Proc. Linn. Soc. New South Wales 92: 213 (1968)     | 2:401 |
|   | <b>Dampiera stricta</b> (Sm.) R.Br., Prodr. 589 (1810)  | 2:404 |
|   | <i>Goodenia amplexans</i> F.Muell., not present in Tasmania, formerly recorded in error           | 2:400 |
|   | <i>Goodenia barbata</i> R.Br. = <b>Cooperhooikia barbata</b>                                      | 2:401 |
|   | <b>Goodenia elongata</b> Labill., Nov. Holl. Pl. 1: 52, t.75 (1805)                               | 2:401 |
|   | <b>Goodenia geniculata</b> R.Br., Prodr. 577 (1810)   | 2:401 |
|   | <b>Goodenia humilis</b> R.Br., Prodr. 575 (1810)  | 2:402 |
|   | <b>Goodenia lanata</b> R.Br., Prodr. 577 (1810)   | 2:401 |
|   | <b>Goodenia ovata</b> Sm., Trans. Linn. Soc. London 2: 347 (1794)                                 | 2:400 |
| x | <b>Goodenia pinnatifida</b> Schldl., Linnaea 21: 450 (1848)                                       |       |
|   | <b>Scaevola aemula</b> R.Br., Prodr. 584 (1810)   | 2:404 |
|   | <b>Scaevola albida</b> (Sm.) Druce, Rep. Bot. Exch. Club Brit. Isles 1916, Suppl. 2: 644 (1917)   | 2:404 |
|   | <i>Scaevola calendulacea</i> (Andrews) Druce, not present in Tasmania, formerly recorded in error | 2:403 |
|   | <b>Scaevola hookeri</b> (Vriese) F.Muell. ex Hook.f., Fl. Tasm. 1: 231 (1856)                     | 2:403 |
|   | <b>Selliera radicans</b> Cav., Ann. Hist. Nat. Madrid 1: 41, t.5 (1799)                           | 2:402 |
|   | <b>Velleia montana</b> Hook.f., Lond. J. Bot. 6: 265 (1847)                                       | 2:399 |
|   | <b>Velleia paradoxa</b> R.Br., Prodr. 580 (1810)  | 2:398 |

#### GROSSULARIACEAE (see also ESCALLONIACEAE)

|   |  |  |
|---|--|--|
| i | <b>Ribes sanguineum</b> Pursh, Fl. Amer. Sept. 1: 164 (1813) |  |
|---|--|--|

#### GUNNERACEAE

|   |  |       |
|---|--|-------|
| e | <b>Gunnera cordifolia</b> (Hook.f.) Hook.f., Fl. Tasm. 1: 125 (1845) | 1:194 |
|---|--|-------|

#### GYROSTEMONACEAE

|  |  |       |
|--|--|-------|
|  | <i>Didymotheca thesioides</i> Hook.f. = <b>Gyrostemon thesioides</b>           | 3:581 |
|  | <b>Gyrostemon thesioides</b> (Hook.f.) A.S.George, Fl. Australia 8: 392 (1982) | 3:581 |

#### HALORAGACEAE

|  |   |       |
|--|---|-------|
|  | <b>Gonocarpus humilis</b> Orch., Bull. Auckland Inst. Mus. 10: 195 (1975) |       |
|  | <b>Gonocarpus micranthus</b> Thunb., Nov. Gen. Pl. 3: 69 (1783)           | 1:190 |



|   |   |       |
|---|---|-------|
|   | subsp. <b>micranthus</b>  |       |
|   | <b>Gonocarpus montanus</b> (Hook.f.) Orch., Bull. Auckland Inst. Mus. 10: 172 (1975)              | 1:192 |
|   | <b>Gonocarpus serpyllifolius</b> Hook.f., Icon. Pl. 3: t.290 (1840)                               | 1:192 |
|   | <b>Gonocarpus tetragynus</b> Labill., Nov. Holl. Pl. 1: 39, t.53 (1805)                           | 1:191 |
|   | <b>Gonocarpus teucroides</b> DC., Prodr. 3: 66 (1828)   | 1:191 |
|   | <b>Haloragis aspera</b> Lindley in T.Mitchell, J. Exped. Trop. Austral. 306 (1848)                | 1:191 |
|   | <b>Haloragis brownii</b> (Hook.f.) Schindler, Pflanzenr. 23: 54 (1905)                            | 1:192 |
|   | <b>Haloragis heterophylla</b> Brongn., Voy. Monde t.68A (1829)                                    | 1:190 |
|   | Haloragis micrantha (Thunb.) R.Br. ex Sieber & Zucc. = <b>Gonocarpus micranthus</b>               | 1:190 |
|   | Haloragis montana Hook.f. = <b>Gonocarpus montanus</b>  | 1:192 |
|   | <b>Haloragis myriocarpa</b> Orch., Bull. Auckland Inst. Mus. 10: 132 (1975)                       |       |
|   | Haloragis serpyllifolia (Hook.f.) Walp. = <b>Gonocarpus serpyllifolius</b>                        | 1:192 |
|   | Haloragis tetragyna (Labill.) Hook.f. = <b>Gonocarpus tetragynus</b>                              | 1:191 |
|   | Haloragis teucroides (DC.) Schldl. = <b>Gonocarpus teucroides</b> & <b>Gonocarpus humilis</b>     | 1:191 |
|   | <b>Myriophyllum amphibium</b> Labill., Nov. Holl. Pl. 2: 70, t.220 (1806)                         | 1:193 |
| i | <b>Myriophyllum aquaticum</b> (Vell.) Verdc., Kew Bull. 23: 36 (1973)                             | 1:193 |
| e | <b>Myriophyllum austropygmaeum</b> Orch., Brunonia 8: 280 (1986)                                  |       |
|   | Myriophyllum brasiliense Cambess. = <b>Myriophyllum aquaticum</b>                                 | 1:193 |
| i | <b>Myriophyllum crispatum</b> Orch., Brunonia 8: 210 (1986)                                       |       |
|   | Myriophyllum elatinoides sensu Curtis & Morris (1975) = <b>Myriophyllum salsugineum</b>           | 1:193 |
| x | <b>Myriophyllum glomeratum</b> Schindler, Pflanzenr. 23: 103 (1905)                               |       |
|   | <b>Myriophyllum integrifolium</b> (Hook.f.) Hook.f., Fl. Tasm. 1: 123 (1856)                      | 1:194 |
|   | <b>Myriophyllum muelleri</b> Sonder, Linnaea 28: 233 (1856)                                       |       |
|   | <b>Myriophyllum pedunculatum</b> Hook.f., London J. Bot. 6: 474 (1847)                            | 1:194 |
|   | subsp. <b>longibracteolatum</b> (Schindler) Orch., Brunonia 8: 273 (1986)                         |       |
|   | subsp. <b>pedunculatum</b>  |       |
|   | Myriophyllum propinquum sensu Curtis & Morris (1975) = <b>M. simulans</b> & <b>M. variifolium</b> | 1:193 |
|   | <b>Myriophyllum salsugineum</b> Orch., Brunonia 4: 48 (1981)                                      | 1:193 |
|   | <b>Myriophyllum simulans</b> Orch., Brunonia 8: 203 (1986)  | 1:193 |
|   | <b>Myriophyllum variifolium</b> Hook.f., Icon. Pl. 3: t.289 (1840)                                | 1:193 |
|   | <b>HYDROPHYLLACEAE</b>  |       |
| i | <b>Phacelia tanacetifolia</b> Benth., Trans. Linn. Soc. 17: 280 (1835)                            | 3:486 |
|   | <b>LAMIACEAE (LABIATAE)</b>   |       |
|   | <b>Ajuga australis</b> R.Br., Prodr. 503 (1810)   | 3:557 |
| i | <b>Ajuga reptans</b> L., Sp. Pl. 561 (1753)   |       |
| i | <b>Cedronella canariensis</b> (L.) Webb & Berthel., Phyt. Canar. 3: 87 (1845)                     | 3:548 |
| i | <b>Lamium amplexicaule</b> L., Sp. Pl. 2: 579 (1753)  | 3:552 |
| i | <b>Lamium purpureum</b> L., Sp. Pl. 2: 579 (1753)   | 3:552 |
|   | <b>Lycopus australis</b> R.Br., Prodr. 500 (1810)   | 3:546 |
| i | <b>Marrubium vulgare</b> L., Sp. Pl. 2: 583 (1753)  | 3:550 |
| i | <b>Melissa officinalis</b> L., Sp. Pl. 2: 592 (1753)  | 3:546 |
|   | <b>Mentha australis</b> R.Br., Prodr. 505 (1810)  | 3:544 |

|                             |  |       |
|-----------------------------|--|-------|
|                             | <b>Mentha diemenica</b> Sprengel, Syst. Veg. 2: 724 (1825)                     | 3:544 |
|                             | var. <b>diemenica</b>  |       |
|                             | var. <b>serpyllifolia</b> (Benth.) J.H.Willis, Muelleria 1: 144 (1967)         |       |
| i                           | <b>Mentha pulegium</b> L., Sp. Pl. 2: 577 (1753)                               | 3:545 |
| i                           | <b>Mentha spicata</b> L., Sp. Pl. 2: 576 (1753)                                | 3:545 |
| i                           | <b>Mentha Xpiperita</b> L., Sp. Pl. 2: 576 (1753)                              | 3:545 |
| i                           | <b>Nepeta cataria</b> L., Sp. Pl. 2: 570 (1753)                                | 3:548 |
| ?x                          | <b>Prostanthera cuneata</b> Benth. in DC., Prodr. 12: 560 (1848)               | 3:554 |
|                             | <b>Prostanthera lasianthos</b> Labill., Nov. Holl. Pl. 2: 18 t.157 (1806)      | 3:553 |
|                             | var. <b>lasianthos</b>   |       |
|                             | <b>Prostanthera rotundifolia</b> R.Br., Prodr. 509 (1810)                      | 3:553 |
| i                           | <b>Prunella laciniata</b> (L.) L., Sp. Pl. ed.2, 2: 837 (1763)                 | 3:550 |
| i                           | <b>Prunella vulgaris</b> L., Sp. Pl. 2: 600 (1753)                             | 3:550 |
|                             | <i>Salvia horminoides</i> Pourr. = <b>Salvia verbenaca</b>                     | 3:547 |
| i                           | <b>Salvia verbenaca</b> L., Sp. Pl. 1: 25 (1753)                               | 3:547 |
|                             | <b>Scutellaria humilis</b> R.Br., Prodr. 507 (1810)                            | 3:549 |
| i                           | <b>Stachys arvensis</b> L., Sp. Pl. ed.2, 2: 814 (1763)                        | 3:551 |
| it                          | <b>Stachys palustris</b> L., Sp. Pl. 2: 580 (1753)                             | 3:551 |
|                             | <b>Teucrium corymbosum</b> R.Br., Prodr. 504 (1810)                            | 3:556 |
| it                          | <b>Teucrium scorodonia</b> L., Sp. Pl. 2: 564 (1753)                           |       |
| e                           | <b>Westringia angustifolia</b> R.Br., Prodr. 501 (1810)                        | 3:555 |
| e                           | <b>Westringia brevifolia</b> Benth., Lab. Gen. & Sp. 459 (1834)                | 3:555 |
| e                           | var. <b>brevifolia</b>   | 3:555 |
| e                           | var. <b>raleighii</b> (B.Boivin) W.M.Curtis, Stud. Fl. Tasm. 3: 650 (1967)     | 3:555 |
|                             | <b>Westringia rigida</b> R.Br., Prodr. 501 (1810)                              | 3:555 |
| e                           | <b>Westringia rubiaefolia</b> R.Br., Prodr. 501 (1810)                         | 3:556 |
| <br><b>LAURACEAE</b>        |  |       |
|                             | <b>Cassytha glabella</b> R.Br., Prodr. 404 (1810)                              | 3:597 |
|                             | <b>Cassytha melantha</b> R.Br., Prodr. 404 (1810)                              | 3:597 |
| e                           | <b>Cassytha pedicellosa</b> J.Z.Weber, J. Adelaide Bot. Gard. 3(3): 214 (1981) |       |
|                             | <b>Cassytha pubescens</b> R.Br., Prodr. 404 (1810)                             | 3:597 |
| <br><b>LENTIBULARIACEAE</b> |  |       |
|                             | <i>Polypompholyx tenella</i> (R.Br.) Lehm. = <b>Utricularia tenella</b>        | 3:537 |
|                             | <b>Utricularia australis</b> R.Br., Prodr. 430 (1810)                          | 3:536 |
|                             | <b>Utricularia dichotoma</b> Labill., Nov. Holl. Pl. 1: 11, t.8 (1804)         | 3:536 |
|                             | <i>Utricularia flexuosa</i> Vahl. = <b>Utricularia australis</b>               | 3:536 |
|                             | <b>Utricularia lateriflora</b> R.Br., Prodr. 431 (1810)                        | 3:536 |
|                             | <b>Utricularia monanthos</b> Hook.f., Fl. Tasm. 1: 299 (1857)                  | 3:537 |
|                             | <b>Utricularia tenella</b> R.Br., Prodr. 432 (1810)                            | 3:537 |
|                             | <b>Utricularia violacea</b> R.Br., Prodr. 431 (1810)                           |       |
| <br><b>LINACEAE</b>         |  |       |

|    |  |      |
|----|--|------|
| e  | <b>Linum albidum</b> Ewart & Jean White, Proc. R. Soc. Vict. ser.2, 23(2): 294, t.54 figs.1,2,4 (1911) |      |
| i  | <b>Linum bienne</b> Miller, Gard. Dict. ed.8, no.8 (1768)  |      |
| it | <b>Linum catharticum</b> L., Sp. Pl. 1: 281 (1753)   | 1:92 |
|    | <b>Linum marginale</b> A.Cunn. in Field, Geogr. Mem. N.S.W. 357 (1825)                                 | 1:92 |
| i  | <b>Linum trigynum</b> L., Sp. Pl. 1: 279 (1753)  | 1:93 |
| i  | <b>Linum usitatissimum</b> L., Sp. Pl. 1: 277 (1753)   | 1:92 |

#### LOGANIACEAE

|   |   |       |
|---|---|-------|
|   | Mitrasacme archeri Hook.f. = <b>Schizacme archeri</b>                             | 3:475 |
|   | Mitrasacme distylis F.Muell. = <b>Phyllangium distylis</b>                        | 3:477 |
|   | Mitrasacme divergens Hook.f. = <b>Phyllangium divergens</b>                       | 3:476 |
|   | Mitrasacme montana Hook.f. ex Benth. = <b>Schizacme montana</b>                   | 3:475 |
|   | Mitrasacme paradoxa sensu W.M.Curtis (1967) = <b>Phyllangium divergens</b>        | 3:476 |
|   | <b>Mitrasacme pilosa</b> Labill., Nov. Holl. Pl. 1: 36, t.49 (1804)               | 3:476 |
|   | var. <b>pilosa</b>  |       |
|   | var. <b>stuartii</b> Hook.f., Fl. Tasm. 1: 274 (1857)                             | 3:476 |
|   | <b>Mitrasacme serpyllifolia</b> R.Br., Prodr. 454 (1810)                          | 3:475 |
|   | <b>Phyllangium distylis</b> (F.Muell.) Dunlop, Fl. Australia 28: 315 (1996)       | 3:477 |
|   | <b>Phyllangium divergens</b> (Hook.f.) Dunlop, Fl. Australia 28: 315 (1996)       | 3:476 |
| e | <b>Schizacme archeri</b> (Hook.f.) Dunlop, Fl. Australia 28: 315 (1996)           | 3:475 |
|   | <b>Schizacme montana</b> (Hook.f. ex Benth.) Dunlop, Fl. Australia 28: 314 (1996) | 3:475 |

#### LYTHRACEAE

|  |   |       |
|--|---|-------|
|  | <b>Lythrum hyssopifolia</b> L., Sp. Pl. 1: 447 (1753) | 2:227 |
|  | <b>Lythrum salicaria</b> L., Sp. Pl. 1: 446 (1753)    | 2:226 |

#### MALVACEAE

|     |  |      |
|-----|--|------|
| e   | <b>Asterotrichion discolor</b> (Hook.) Melville, Kew Bull. 20: 512 (1967)  | 1:86 |
|     | <b>Gynatrix pulchella</b> (Willd.) Alef., Osterr. Bot. Z. 12: 35 (1862)    | 1:87 |
| i#  | <b>Hibiscus trionum</b> L., Sp. Pl. 2: 697 (1753)                          | 1:88 |
|     | Lavatera arborea L. = <b>Malva dendromorpha</b>                            | 1:84 |
|     | Lavatera cretica L. = <b>Malva linnaei</b>                                 | 1:84 |
|     | Lavatera plebeia Sims = <b>Malva australiana</b>                           | 1:83 |
|     | <b>Lawrencia spicata</b> Hook., Icon. Pl. 3: t.261 (1840)                  | 1:88 |
| ?i# | <b>Lawrencia squamata</b> Nees ex Miq. in Lehm., Pl. Preiss. 1: 242 (1845) |      |
|     | <b>Malva australiana</b> M.F.Ray, Novon 8(3): 291 (1998)                   | 1:83 |
| i   | <b>Malva dendromorpha</b> M.F.Ray, Novon 8(3): 292 (1998)                  | 1:84 |
| i   | <b>Malva linnaei</b> M.F.Ray, Novon 8(3): 292 (1998)                       | 1:84 |
| it  | <b>Malva moschata</b> L., Sp. Pl. 2: 690 (1753)                            | 1:85 |
| i   | <b>Malva neglecta</b> Wallr., Syll. Pl. Nov. Ratisbon. 1: 140 (1824)       | 1:85 |
| i   | <b>Malva nicaeensis</b> All., Fl. Pedem. 2: 40 (1785)                      | 1:85 |
| i   | <b>Malva parviflora</b> L., Demonstr. Pl. 18 (1753)                        | 1:85 |
| i   | <b>Malva sylvestris</b> L., Sp. Pl. 2: 689 (1753)                          | 1:85 |
| i   | <b>Modiola caroliniana</b> (L.) G.Don, Gen. Syst. 1: 466 (1831)            | 1:86 |

|  |   |      |
|--|---|------|
|  | Plagianthus pulchellus (Willd.) Hook.f. = <b>Gynatrix pulchella</b> | 1:87 |
|  | Plagianthus sidoides Hook. = <b>Asterotrichion discolor</b>         | 1:86 |
|  | Plagianthus spicatus (Hook.) Benth. = <b>Lawrencia spicata</b>      | 1:88 |

#### MENYANTHACEAE

|   |   |       |
|---|---|-------|
| t | <b>Liparophyllum gunnii</b> Hook.f., London J. Bot. 6: 473 bis (1847)                     | 3:484 |
|   | Nymphoides crenata (F.Muell.) Kuntze, not present in Tasmania, formerly recorded in error | 3:483 |
| e | <b>Nymphoides exigua</b> (F.Muell.) Kuntze, Rev. Gen. Pl. 2: 429 (1891)                   | 3:483 |
|   | <b>Villarsia exaltata</b> (Sol. ex Sims) G.Don, Gen. Hist. 4: 169 (1838)                  | 3:483 |
|   | <b>Villarsia reniformis</b> R.Br., Prodr. 457 (1810)                                      | 3:483 |

#### MIMOSACEAE

|   |   |       |
|---|---|-------|
|   | Acacia armata R.Br. = <b>Acacia paradoxa</b>  | 1:127 |
| e | <b>Acacia axillaris</b> Benth., Lond. J. Bot. 1: 341 (1842)                         | 1:127 |
|   | Acacia botrycephala (Vent.) Desf. = <b>Acacia terminalis</b>                        | 1:129 |
|   | <b>Acacia dealbata</b> Link, Enum. Hort. Berol. Alt. 2: 445 (1822)                  | 1:130 |
|   | subsp. <b>dealbata</b>  |       |
| i | <b>Acacia decurrens</b> (J.Wendl.) Willd., Sp. Pl. ed.4, 4: 1072 (1806)             | 1:130 |
|   | <b>Acacia genistifolia</b> Link, Enum. Hort. Berol. Alt. 2: 442 (1822)              | 1:126 |
|   | <b>Acacia gunnii</b> Benth., Lond. J. Bot. 1: 332 (1842)                            | 1:127 |
|   | <b>Acacia implexa</b> Benth., Lond. J. Bot. 1: 368 (1842)                           |       |
|   | <b>Acacia longifolia</b> (Andrews) Willd., Sp. Pl. ed.4, 4: 1052 (1806)             | 1:127 |
|   | subsp. <b>longifolia</b>  | 1:129 |
|   | subsp. <b>sophorae</b> (Labill.) Court, Fl. Australia 11B: 491 (2001)               | 1:129 |
|   | <b>Acacia mearnsii</b> De Wild., Pl. Bequaert. 3: 61 (1925)                         | 1:130 |
|   | <b>Acacia melanoxylon</b> R.Br. in W.T.Aiton, Hortus Kew. ed.2, 5: 462 (1813)       | 1:128 |
|   | <b>Acacia mucronata</b> Willd. ex Wendl.f., Comment. Acac. 6, 46 t.12 (1820)        | 1:129 |
| e | subsp. <b>dependens</b> (Hook.f.) Court, Fl. Australia 11B: 491 (2001)              | 1:129 |
|   | subsp. <b>longifolia</b> (Benth.) Court, Fl. Australia 11B: 491 (2001)              | 1:129 |
|   | subsp. <b>mucronata</b>   | 1:129 |
|   | <b>Acacia myrtifolia</b> (Smith) Willd., Sp. Pl. ed.4, 4: 1054 (1806)               | 1:128 |
| i | <b>Acacia paradoxa</b> DC., Cat. Pl. Horti Monsp. 74 (1813)                         | 1:127 |
| e | <b>Acacia pataczekii</b> D.I.Morris, Rec. Q. Vic. Mus. 50: 1 (1974)                 | 1:128 |
| i | <b>Acacia pycnantha</b> Benth., Lond. J. Bot. 1: 351 (1842)                         |       |
|   | <b>Acacia retinodes</b> Schldtl., Linnaea 20: 664 (1847)                            | 1:128 |
|   | var. <b>uncifolia</b> J.M.Black, Trans. & Proc. Roy. Soc. S.Australia 56: 42 (1932) | 1:128 |
| e | <b>Acacia riceana</b> Henslow in Maund, Botanist 3: t.135 (1839)                    | 1:127 |
|   | <b>Acacia siculiformis</b> A.Cunn. ex Benth., Lond. J. Bot. 1: 337 (1842)           | 1:126 |
|   | Acacia sophorae (Labill.) R.Br. = <b>Acacia longifolia</b> subsp. <b>sophorae</b>   | 1:129 |
|   | <b>Acacia stricta</b> (Andrews) Willd., Sp. Pl. ed.4, 4: 1052 (1806)                | 1:127 |
|   | <b>Acacia suaveolens</b> (Smith) Willd., Sp. Pl. ed.5, 4: 1050 (1806)               | 1:128 |
|   | <b>Acacia terminalis</b> (Salisb.) Macbr., Contr. Gray Herb. n.ser.59: 7 (1919)     | 1:129 |
|   | <b>Acacia ulicifolia</b> (Salisb.) Court, Vict. Nat. 73: 173 (1957)                 | 1:126 |
|   | <b>Acacia verniciflua</b> A.Cunn. in Field, Geogr. Mem. N.S.W. 344 (1825)           | 1:127 |

|                        |  |       |
|------------------------|--|-------|
|                        | <b>Acacia verticillata</b> (L'Hérit.) Willd., Sp. Pl. ed.4, 4: 1049 (1806)                         | 1:126 |
|                        | subsp. <b>ovoidea</b> (Benth.) Court, Fl. Australia 11B: 492 (2001)                                | 1:126 |
|                        | subsp. <b>ruscifolia</b> (A.Cunn. ex G.Don) Court, Fl. Australia 11B: 492 (2001)                   | 1:126 |
|                        | subsp. <b>verticillata</b>   | 1:126 |
|                        | var. <i>latifolia</i> sensu Curtis & Morris (1975) = subsp. <b>ruscifolia</b>                      | 1:126 |
|                        | <i>Albizia lophantha</i> (Willd.) Benth. = <b>Paraserianthes lophantha</b>                         | 1:130 |
| i                      | <b>Paraserianthes lophantha</b> (Willd.) I.Nielsen, Adansonia ser.4, 5: 326 (1983)                 | 1:130 |
| i                      | subsp. <b>lophantha</b>  |       |
|                        | <i>Racosperma axillare</i> (Benth.) Pedley = <b>Acacia axillaris</b>                               | 1:127 |
|                        | <i>Racosperma dealbatum</i> (Link) Pedley = <b>Acacia dealbata</b>                                 | 1:130 |
|                        | <i>Racosperma decurrens</i> (Willd.) Pedley = <b>Acacia decurrens</b>                              | 1:130 |
|                        | <i>Racosperma gunnii</i> (Benth.) Pedley = <b>Acacia gunnii</b>                                    | 1:127 |
|                        | <i>Racosperma implexum</i> (Benth.) Pedley = <b>Acacia implexa</b>                                 |       |
|                        | <i>Racosperma mearnsii</i> (De Wild.) Pedley = <b>Acacia mearnsii</b>                              | 1:130 |
|                        | <i>Racosperma melanoxyton</i> (R.Br.) Pedley = <b>Acacia melanoxyton</b>                           | 1:128 |
|                        | <i>Racosperma myrtifolium</i> (Smith) Martius = <b>Acacia myrtifolia</b>                           | 1:128 |
|                        | <i>Racosperma paradoxum</i> (DC.) Martius = <b>Acacia paradoxa</b>                                 | 1:127 |
|                        | <i>Racosperma riceanum</i> (Henslow) Pedley = <b>Acacia riceana</b>                                | 1:127 |
|                        | <i>Racosperma sophorae</i> (Labill.) Martius = <b>Acacia longifolia</b> subsp. <b>sophorae</b>     | 1:129 |
|                        | <i>Racosperma strictum</i> (Andrews) Martius = <b>Acacia stricta</b>                               | 1:127 |
|                        | <i>Racosperma suaveolens</i> (Smith) Martius = <b>Acacia suaveolens</b>                            | 1:128 |
|                        | <i>Racosperma ulicifolium</i> (Salisb.) Pedley = <b>Acacia ulicifolia</b>                          | 1:126 |
|                        | <i>Racosperma vernicifluum</i> (A.Cunn.) Pedley = <b>Acacia verniciflua</b>                        | 1:127 |
|                        | <i>Racosperma verticillatum</i> (L'Hérit.) Martius = <b>Acacia verticillata</b>                    | 1:126 |
| <br><b>MONIMIACEAE</b> |  |       |
|                        | <b>Atherosperma moschatum</b> Labill., Nov. Holl. Pl. 2: 74, t.224 (1806)                          | 3:594 |
|                        | <b>Hedycarya angustifolia</b> R.Cunn., Ann. Nat. Hist. ser.1 1: 215 (1838)                         | 3:595 |
| <br><b>MYOPORACEAE</b> |  |       |
|                        | <b>Myoporum insulare</b> R.Br., Prodr. 516 (1810)  | 3:540 |
|                        | <b>Myoporum parvifolium</b> R.Br., Prodr. 516 (1810)   | 3:540 |
| <br><b>MYRTACEAE</b>   |  |       |
|                        | <b>Baeckea gunniana</b> Schauer in Walpers, Repert. Bot. Sys. 2: 920 (1843)                        | 1:198 |
| e                      | <b>Baeckea leptocaulis</b> Hook.f., Icon. Pl. 3: t.298 (1840)                                      | 1:199 |
|                        | <i>Baeckea ramosissima</i> A.Cunn. = <b>Euryomyrtus ramosissima</b>                                | 1:198 |
|                        | <b>Callistemon pallidus</b> (Bonpl.) DC., Prodr. 3: 223 (1828)                                     | 1:203 |
|                        | <i>Callistemon paludosus</i> sensu Curtis & Morris (1975) = <b>Callistemon pallidus</b> sens. lat. | 1:203 |
| e                      | <b>Callistemon viridiflorus</b> (Sims) Sweet, Hort. Brit. 155 (1826)                               | 1:203 |
|                        | <b>Calytrix tetragona</b> Labill., Nov. Holl. Pl. 2: 8, t.146 (1806)                               | 1:197 |
| e                      | <b>Eucalyptus amygdalina</b> Labill., Nov. Holl. Pl. 2: 14, t.154 (1806)                           | 1:224 |
| e                      | <b>Eucalyptus archeri</b> Maiden & Blakely, Crit. Revis. Eucalyptus 8: 58 (1929)                   | 1:215 |
| e                      | <b>Eucalyptus barberi</b> L.A.S.Johnson & Blaxell, Contr. N.S.W. Natl. Herb. 4: 288 (1972)         | 1:211 |

|   |   |       |
|---|---|-------|
|   | <i>Eucalyptus bicostata</i> sensu Curtis & Morris (1975) = <b>E. globulus</b> subsp. <b>pseudoglobulus</b>                | 1:219 |
|   | <b>Eucalyptus brookeriana</b> A.M.Gray, Austral. Forest Res. 9: 111 (1979)  |       |
| e | <b>Eucalyptus coccifera</b> Hook.f., London J. Bot. 6: 477 (1847)   | 1:225 |
| e | <b>Eucalyptus cordata</b> Labill., Nov. Holl. Pl. 2: 13, t.152 (1806)   | 1:218 |
|   | <b>Eucalyptus dalrympleana</b> Maiden, For. Fl. N.S.W. 7: 137, t.241 (1920)   | 1:214 |
|   | subsp. <b>dalrympleana</b>  |       |
|   | <b>Eucalyptus delegatensis</b> R.Baker, Proc. Linn. Soc. N.S.W. 25: 305, t.16 (1900)                                      | 1:220 |
| e | subsp. <b>tasmaniensis</b> Boland, Austral. Forest Res. 15: 177 (1985)  |       |
|   | <i>Eucalyptus gigantea</i> Hook.f. = <b>Eucalyptus delegatensis</b>   | 1:220 |
|   | <b>Eucalyptus globulus</b> Labill., Voy. Rech. Per. 1: 153, t.13 (1800)   | 1:218 |
|   | subsp. <b>globulus</b>  |       |
|   | subsp. <b>pseudoglobulus</b> (Naudin ex Maiden) J.B.Kirkp., Bot. J. Linn. Soc. 69: 101 (1975)                             | 1:219 |
| e | <b>Eucalyptus gunnii</b> Hook.f., London J. Bot. 3: 499 (1844)  | 1:215 |
| e | subsp. <b>divaricata</b> (McAulay & Brett) B.M.Potts, Pap. Proc. R. Soc. Tasm. 135: 57 (2001)                             |       |
| e | subsp. <b>gunnii</b>  |       |
| e | <b>Eucalyptus johnstonii</b> Maiden, Crit. Revis. Eucalyptus 6: 280 (1922)  | 1:219 |
|   | <i>Eucalyptus linearis</i> Dehnh. = <b>Eucalyptus pulchella</b>   | 1:224 |
| e | <b>Eucalyptus morrisbyi</b> Brett, Proc. R. Soc. Tasm. 1938: 129 (1939)   | 1:215 |
| e | <b>Eucalyptus nitida</b> Hook.f., Fl. Tasm. 1: 137, t.29 (1856)   | 1:225 |
|   | <b>Eucalyptus obliqua</b> L'Hérit., Sert. Angl. 18 (1789); t.20 (1792)  | 1:220 |
|   | <b>Eucalyptus ovata</b> Labill., Nov. Holl. Pl. 2: 13, t.153 (1806)   | 1:211 |
|   | <b>Eucalyptus pauciflora</b> Sieber ex Sprengel, Sys. Veg. 4: 195 (1827)  | 1:222 |
|   | subsp. <b>pauciflora</b>  |       |
|   | <b>Eucalyptus perriniana</b> F.Muell. ex Rodway, Pap. Proc. R. Soc. Tasm. 1893: 181 (1894)                                | 1:217 |
| e | <b>Eucalyptus pulchella</b> Desf., Cat. Hort. Paris. ed.3 284: 408 (1829)   | 1:224 |
|   | <b>Eucalyptus radiata</b> Sieber ex DC., Prodr. 3: 218 (1828)   | 1:224 |
|   | subsp. <b>robertsonii</b> (Blakely) L.A.S.Johnson & Blaxell, Cont. N.S.W. Natl. Herb. 4: 380 (1973)                       |       |
|   | <b>Eucalyptus regnans</b> F.Muell., Ann. Rep. Vict. Acclim. Soc. 20 in obs. (1870-71)                                     | 1:220 |
| e | <b>Eucalyptus risdonii</b> Hook.f., London J. Bot. 6: 477 bis (1847)  | 1:226 |
| e | <b>Eucalyptus rodwayi</b> R.Baker & H.G.Smith, Pap. Proc. R. Soc. Tasm. 191 (1912);<br>Res. Eucalyptus Tasmania 53 (1912) | 1:211 |
|   | <b>Eucalyptus rubida</b> Deane & Maiden, Proc. Linn. Soc. N.S.W. 24: 456, t.x1 (1899)                                     | 1:214 |
|   | <i>Eucalyptus salicifolia</i> Cav. = <b>E. saligna</b> Smith (not present in Tasmania (Cameron 1955))                     | 1:224 |
|   | <b>Eucalyptus sieberi</b> L.A.S.Johnson, Contr. N.S.W. Natl. Herb. 3: 125 (1962)  | 1:222 |
|   | <i>Eucalyptus simmondsii</i> Maiden = <b>Eucalyptus nitida</b>  | 1:225 |
| e | <b>Eucalyptus subcrenulata</b> Maiden & Blakely, Crit. Revis. Eucalyptus 8: 59 (1929)                                     | 1:219 |
|   | <i>Eucalyptus tasmanica</i> Blakely = <b>Eucalyptus tenuiramis</b>  | 1:226 |
| e | <b>Eucalyptus tenuiramis</b> Miq., Ned. Kruidk. Arch. 4: 128 (1856)   | 1:226 |
| e | <b>Eucalyptus urnigera</b> Hook.f., London J. Bot. 6: 477 bis (1847)  | 1:217 |
| e | <b>Eucalyptus vernicosa</b> Hook.f., London J. Bot. 6: 478 bis (1847)   | 1:219 |
|   | <b>Eucalyptus viminalis</b> Labill., Nov. Holl. Pl. 2: 12, t.151 (1806)   | 1:214 |
|   | subsp. <b>viminalis</b>   |       |
|   | <b>Euryomyrtus ramosissima</b> (A.Cunn.) Trudgen, Nuytsia 13(3): 560 (2001)   | 1:198 |
|   | subsp. <b>prostrata</b> (Hook.f.) Trudgen, Nuytsia 13(3): 563 (2001)  |       |

|                       |  |       |
|-----------------------|--|-------|
|                       | subsp. <b>ramosissima</b>  |       |
|                       | <b>Kunzea ambigua</b> (Smith) Druce, Bot. Soc. Exch. Club Brit. Isles 1916: 629 (1917)                                   | 1:202 |
| ni                    | <b>Kunzea ericoides</b> (A.Rich.) Joy Thomps., Telopea 2: 379 (1983)   |       |
| e                     | <b>Leptospermum glaucescens</b> S.Schauer, Linnaea 15: 421 (1841)  | 1:201 |
| e                     | <b>Leptospermum grandiflorum</b> Lodd., Bot. Cab. 6: t.514 (1821)  | 1:200 |
|                       | <b>Leptospermum laevigatum</b> (Gaertner) F.Muell., Cat. Pl. Melb. Bot. Gard. 22 (1858)                                  | 1:200 |
|                       | <b>Leptospermum lanigerum</b> (Aiton) Smith, Trans. Linn. Soc. 3: 263 (1797)   | 1:200 |
| e                     | <b>Leptospermum nitidum</b> Hook.f., Fl. Tasm. 1: 139 (1856)   | 1:201 |
| e                     | <b>Leptospermum riparium</b> D.I.Morris, Rec. Q. Vic. Mus. 50: 2 (1974)  | 1:201 |
| e                     | <b>Leptospermum rupestre</b> Hook.f., Icon. Pl. 4: t.308 (1840)  | 1:202 |
|                       | <b>Leptospermum scoparium</b> Forst. & Forst.f., Char. Gen. Pl. 72, t.36, figs.f-l (1776)                                | 1:200 |
|                       | var. <b>eximium</b> B.L.Burtt, Bot. Mag. 162: t.9582 (1939)  | 1:200 |
|                       | var. <b>scoparium</b>  |       |
|                       | <b>Melaleuca armillaris</b> (Sol. ex Gaertner) Smith, Trans. Linn. Soc. 3: 277 (1797)                                    | 1:205 |
|                       | subsp. <b>armillaris</b>   |       |
|                       | <b>Melaleuca ericifolia</b> Smith, Trans. Linn. Soc. 3: 276 (1797)   | 1:204 |
|                       | <b>Melaleuca gibbosa</b> Labill., Nov. Holl. Pl. 2: 30, t.172 (1806)   | 1:204 |
| e                     | <b>Melaleuca pustulata</b> Hook.f., Lond. J. Bot. 6: 476 bis (1847)  | 1:205 |
|                       | <b>Melaleuca squamea</b> Labill., Nov. Holl. Pl. 2: 28, t.168 (1806)   | 1:204 |
|                       | <b>Melaleuca squarrosa</b> Donn ex Smith, Trans. Linn. Soc. 6: 300 (1802)  | 1:204 |
|                       | <b>Thryptomene micrantha</b> Hook.f., J. Bot. Kew Gard. (Hooker) 5: 299, t.8 (1853)                                      | 1:198 |
| <br><b>OLEACEAE</b>   |  |       |
| i                     | <b>Ligustrum vulgare</b> L., Sp. Pl. 1: 7 (1753)   | 3:470 |
|                       | <b>Notelaea ligustrina</b> Vent., Choix Pl. 25 (1804)  | 3:470 |
| <br><b>ONAGRACEAE</b> |  |       |
|                       | <b>Boisduvalia tasmanica</b> (Hook.f.) Munz = <b>Epilobium curtisiae</b>   | 2:234 |
|                       | <b>Epilobium adenocaulon</b> Hausskn. = <b>Epilobium ciliatum</b>  | 2:233 |
|                       | <b>Epilobium billardierianum</b> Sér. ex DC., Prodr. 3: 41 (1828)  | 2:232 |
|                       | subsp. <b>billardierianum</b>  |       |
|                       | subsp. <b>cinereum</b> (A.Rich.) Raven & Engelhorn, N.Z. J. Bot. 9: 349 (1971)   | 2:230 |
|                       | subsp. <b>hydrophilum</b> Raven & Engelhorn (Tasmanian plants) = <b>E. billardierianum</b> subsp. <b>billardierianum</b> |       |
|                       | subsp. <b>intermedium</b> Raven & Engelhorn, N.Z. J. Bot. 9: 348 (1971)  |       |
| i                     | <b>Epilobium ciliatum</b> Raf., Med. Repos. 5: 361 (1808)  | 2:233 |
|                       | <b>Epilobium curtisiae</b> Raven, Aliso 5: 249, fig.1 (1963)   | 2:234 |
|                       | <b>Epilobium erosum</b> Hausskn. = <b>Epilobium gunnianum</b>  | 2:230 |
| e                     | <b>Epilobium fugitivum</b> Raven & Engelhorn, N.Z. J. Bot. 9: 346 (1971)   |       |
|                       | <b>Epilobium gunnianum</b> Hausskn., Oesterr. Bot. Z. 29: 149 (1879)   | 2:230 |
|                       | <b>Epilobium hirtigerum</b> A.Cunn., Ann. Nat. Hist. 3: 33 (1839)  | 2:231 |
|                       | <b>Epilobium junceum</b> Forst.f. ex Sprengel = <b>Epilobium billardierianum</b> subsp. <b>cinereum</b>                  | 2:230 |
| i#                    | <b>Epilobium nummulariifolium</b> A.Cunn., Ann. Nat. Hist. 3: 31 (1839)  |       |
| it                    | <b>Epilobium obscurum</b> Schreber, Spicil. Fl. Lipsicae 147, 155 (1771)   |       |
|                       | <b>Epilobium pallidiflorum</b> Sol. ex A.Cunn., Ann. Nat. Hist. 3: 34 (1839)   | 2:231 |

|    |   |       |
|----|---|-------|
| e  | <b>Epilobium perpusillum</b> Hausskn., Monogr. Epilobium 301, t.21, fig.90 (1884) | 2:233 |
| it | <b>Epilobium rotundifolium</b> Forst.f., Fl. Ins. Austral. 27 (1786)              |       |
|    | <b>Epilobium sarmentaceum</b> Hausskn., Oesterr. Bot. Z. 29: 149 (1879)           | 2:229 |
|    | <b>Epilobium tasmanicum</b> Hausskn., Monogr. Epilobium 296, t.20, fig.84 (1884)  | 2:233 |
|    | <b>Epilobium willisii</b> Raven & Engelhorn, N.Z. J. Bot. 9: 347 (1971)           |       |
| i  | <b>Fuchsia magellanica</b> Lam., Encycl. 2: 565 (1788)                            | 2:235 |
| i# | <b>Oenothera biennis</b> L., Sp. Pl. 1: 346 (1753)                                |       |
| i  | <b>Oenothera glazioviana</b> M.Micheli in C.P.Martius, Fl. Brasil 13: 178 (1882)  |       |
| i  | <b>Oenothera stricta</b> Ledeb. ex Link, Enum. Pl. Hort. Berol. 1: 377 (1821)     | 2:235 |

#### OROBANCHACEAE

Orobanche australiana F.Muell., not present in Tasmania, formerly recorded in error

|   |  |       |
|---|--|-------|
| i | <b>Orobanche minor</b> Smith, Engl. Bot. 6: t.422 (1797) | 3:533 |
|---|--|-------|

#### OXALIDACEAE

|   |   |       |
|---|---|-------|
| i | <b>Oxalis articulata</b> Savigny in Lam., Encycl. Méth. Bot. 4: 686 (1798)  | 1:102 |
| i | <b>Oxalis corniculata</b> L., Sp. Pl. 1: 435 (1753)                         | 1:101 |
| i | subsp. <b>corniculata</b>   |       |
|   | <b>Oxalis exilis</b> A.Cunn., Ann. Nat. Hist. 3: 316 (1839)                 |       |
| i | <b>Oxalis incarnata</b> L., Sp. Pl. 1: 433 (1753)                           | 1:102 |
|   | <b>Oxalis lactea</b> Hook. = <b>Oxalis magellanica</b>                      | 1:101 |
| i | <b>Oxalis latifolia</b> Kunth, Nov. Gen. Sp. 5: 237, t.467 (1822)           | 1:101 |
|   | <b>Oxalis magellanica</b> Forst.f., Comm. Soc. Reg. Sci. Gött. 9: 33 (1789) | 1:101 |
|   | <b>Oxalis perennans</b> Haw., Miscel. 181 (1803)                            |       |
| i | <b>Oxalis pes-caprae</b> L., Sp. Pl. 1: 434 (1753)                          | 1:101 |
| i | <b>Oxalis purpurea</b> L., Sp. Pl. 1: 433 (1753)                            | 1:102 |
|   | <b>Oxalis radicata</b> A.Rich., Tent. Fl. Abyss. 2: 123 (1847)              |       |

#### PAPAVERACEAE

|    |  |      |
|----|--|------|
|    | <b>Argemone mexicana</b> sensu Curtis & Morris (1975) = <b>Argemone ochroleuca</b> | 1:28 |
| i  | <b>Argemone ochroleuca</b> Sweet, Brit. Fl. Gard. 3, t.242 (1828)                  | 1:28 |
| i  | <b>Eschscholzia californica</b> Cham. in Nees, Horae Phys. Berol. 74, t.15 (1820)  | 1:29 |
| i  | <b>Glaucium flavum</b> Crantz, Stirp. Austriac. ed.1, fasc.2: 133 (1763)           | 1:28 |
| ix | <b>Papaver aculeatum</b> Thunb., Prodr. Pl. Cap. 92 (1800)                         | 1:27 |
| i  | <b>Papaver argemone</b> L., Sp. Pl. 1: 506 (1753)                                  | 1:27 |
| i  | <b>Papaver dubium</b> L., Sp. Pl. 2: 1196 (1753)                                   | 1:27 |
| i  | <b>Papaver hybridum</b> L., Sp. Pl. 1: 506 (1753)                                  | 1:27 |
| i  | <b>Papaver rhoeas</b> L., Sp. Pl. 1: 507 (1753)                                    | 1:27 |
| i  | <b>Papaver somniferum</b> L., Sp. Pl. 1: 508 (1753)                                | 1:28 |
| i  | subsp. <b>setigerum</b> (DC.) Corb., Nouv. Fl. Normandie 30 (1894)                 |      |
| i  | subsp. <b>somniferum</b>   |      |

#### PASSIFLORACEAE

|    |  |  |
|----|--|--|
| i# | <b>Passiflora cinnabarina</b> Lindley, Gard. Chron. 23: 724 (1855) |  |
|----|--|--|



*Passiflora mollissima* sensu Census (1999) = **Passiflora tarminiana**

i **Passiflora tarminiana** Coppens & V.E.Barney, Novon 11: 9 (2001)

#### PITTOSPORACEAE

*Billardiera alpina* (McGillivray) E.Bennett = **Rhytidosporum inconspicuum**

i **Billardiera heterophylla** (Lindley) L.Cayzer & Crisp, Aust. Syst. Bot. 17(1): 119 (2004) 1:59

e **Billardiera longiflora** Labill., Nov. Holl. Pl. 1: 64, t.89 (1805) 1:58

var. *alpina* Rodway = **Billardiera longiflora** 1:58

var. *ovalis* (Lindley) E.Bennett = **Billardiera ovalis**

n **Billardiera macrantha** Hook.f., Fl. Tasm. 1: 37 (1855)

n **Billardiera mutabilis** Salisb. in Salisb. & Hook., Paradis. Lond. 1: t.48 (1806) 1:59

ne **Billardiera nesophila** L.Cayzer & D.L.Jones, Aust. Syst. Bot. 17(1): 105 (2004)

e **Billardiera ovalis** Lindley, Edwards's Bot. Reg. 20: t.1719 (1834)

*Billardiera procumbens* (Hook.) E.Bennett = **Rhytidosporum procumbens** 1:58

*Billardiera scandens* sensu Curtis & Morris (1975) = **Billardiera mutabilis** 1:59

ne **Billardiera viridiflora** L.Cayzer & D.L.Jones, Aust. Syst. Bot. 17(1): 105 (2004)

**Bursaria spinosa** Cav., Icon. 4: 30, t.350 (1797) 1:57

*Marianthus procumbens* (Hook.) Benth. = **Rhytidosporum procumbens** 1:58

**Pittosporum bicolor** Hook., J. Bot. (Hooker) 1: 249 (1834) 1:56

i **Pittosporum undulatum** Vent., Descr. Pl. Nouv. Jard. Cels. 76, t.76 (1802)

subsp. **Xemmettii** W.M.Curtis in Morris & Curtis, Rec. Q. Vic. Mus. 50: 3 (1974) 1:57

i subsp. **undulatum**

*Rhytidosporum alpinum* sensu Census 1999 = **Rhytidosporum inconspicuum**

**Rhytidosporum inconspicuum** L.Cayzer, Crisp & I.Telford, Aust. Syst. Bot. 12(5): 700 (1999)

**Rhytidosporum procumbens** (Hook.) F.Muell., Pl. Vict. 1: 75 (1860) 1:58

*Sollya heterophylla* Lindley = **Billardiera heterophylla** 1:59

#### PLANTAGINACEAE

**Plantago antarctica** Decne. in DC., Prodr. 13: 703 (1852) 3:562

i **Plantago australis** Lam., Tabl. Encycl. 1: 339 (1792)

e **Plantago bellidioides** Decne. in DC., Prodr. 13: 701 (1852) 3:561

i **Plantago coronopus** L., Sp. Pl. 1: 115 (1753) 3:560

ni subsp. **commutata** (Guss.) Pilg., Repert. Spec. Nov. Regni Veg. 28:287 (1930)

e **Plantago daltonii** Decne. in DC., Prodr. 13: 703 (1852) 3:563

**Plantago debilis** R.Br., Prodr. 425 (1810) 3:560

**Plantago gaudichaudii** Barnéoud, Monogr. Plantag. 15 (1845)

e **Plantago glabrata** Hook.f., Hook. Lond. J. Bot. 6: 276 (1847) 3:562

**Plantago glacialis** B.G.Briggs, Carolin & Pulley, Contr. N.S.W. Natl. Herb. 4: 395 (1973) 3:564

e **Plantago gunnii** Hook.f., Hook. Lond. J. Bot. 5: 446 (1846) 3:564

**Plantago hispida** R.Br., Prodr. 425 (1810) 3:561

i **Plantago lanceolata** L., Sp. Pl. 1: 113 (1753) 3:560

i **Plantago major** L., Sp. Pl. 1: 112 (1753) 3:559

*Plantago muelleri* sensu W.M.Curtis (1967) = **Plantago glacialis** 3:564

e **Plantago paradoxa** Hook.f., Hook. Lond. J. Bot. 6: 277 (1847) 3:564

|   |   |       |
|---|---|-------|
| e | <b>Plantago tasmanica</b> Hook.f., Hook. Lond. J. Bot. 6: 276 (1847)    | 3:562 |
| e | var. <b>archeri</b> (Hook.f.) W.M.Curtis, Stud. Fl. Tasm. 3: 650 (1967) | 3:563 |
| e | var. <b>tasmanica</b>   | 3:562 |
| t | <b>Plantago triantha</b> Sprengel, Syst. Veg. 1: 439 (1824)             | 3:563 |
|   | <b>Plantago varia</b> R.Br., Prodr. 425 (1810)                          | 3:561 |

#### PLUMBAGINACEAE

|   |  |       |
|---|--|-------|
|   | <b>Limonium australe</b> (R.Br.) Kuntze, Revis. Gen. Pl. 1: 395 (1891) | 3:465 |
| e | <b>Limonium baudinii</b> Lincz., Nov. Syst. Pl. Vasc. 23: 107 (1986)   |       |

#### POLEMONIACEAE

|     |   |       |
|-----|---|-------|
| it# | <b>Collomia linearis</b> (Cav.) Nutt., Gen. N.Am. Pl. 1: 126 (1818)               | 3:485 |
| i   | <b>Navarretia squarrosa</b> (Eschsch.) Hook. & Arn., Bot. Beechey Voy. 368 (1839) | 3:485 |

#### POLYGALACEAE

|   |  |      |
|---|--|------|
|   | <b>Comesperma calymega</b> Labill., Nov. Holl. Pl. 2: 23, t.162 (1806) | 1:62 |
|   | <b>Comesperma defoliatum</b> F.Muell., Pl. Victoria 1: 189 (1862)      | 1:63 |
|   | <b>Comesperma ericinum</b> DC., Prodr. 1: 334 (1824)                   | 1:62 |
|   | <b>Comesperma retusum</b> Labill., Nov. Holl. Pl. 2: 22, t.160 (1806)  | 1:62 |
|   | <b>Comesperma volubile</b> Labill., Nov. Holl. Pl. 2: 24, t.163 (1806) | 1:62 |
| i | <b>Polygala myrtifolia</b> L., Sp. Pl. 2: 703 (1753)                   | 1:63 |
| i | <b>Polygala vulgaris</b> L., Sp. Pl. 2: 702 (1753)                     |      |

#### POLYGONACEAE

|    |   |       |
|----|---|-------|
| i  | <b>Acetosa sagittata</b> (Thunb.) L.A.S.Johnson & B.G.Briggs, Contr. N.S.W. Natl. Herb. 3: 166 (1962) |       |
| i  | <b>Acetosella vulgaris</b> Fourr., Ann. Soc. Linn. Lyon ser.2, 17: 145 (1869)                         | 3:587 |
| i  | <b>Emex australis</b> Steinh., Ann. Sci. Nat. Bot. ser.2, 9: 195 (1838)                               | 3:583 |
| i  | <b>Fallopia convolvulus</b> (L.) A.Löve, Taxon 19: 300 (1970)   | 3:591 |
| i  | <b>Fallopia japonica</b> (Houtt.) Ronse Decraene, Bot. J. Linn. Soc. 98(4): 369 (1988)                |       |
|    | <b>Muehlenbeckia adpressa</b> (Labill.) Meisn., Pl. Vasc. Gen. 227 (1841)                             | 3:592 |
|    | <b>Muehlenbeckia axillaris</b> (Hook.f.) Endl., Gen. Pl. Suppl. 4(2): 51 (1848)                       | 3:593 |
|    | <b>Muehlenbeckia gunnii</b> (Hook.f.) Endl., Gen. Pl. Suppl. 4(2): 51 (1848)                          | 3:593 |
|    | <b>Persicaria decipiens</b> (R.Br.) K.L.Wilson, Telopea 3: 178 (1988)                                 | 3:590 |
|    | <b>Persicaria hydropiper</b> (L.) Spach, Hist. Nat. Vég. 10: 536 (1841)                               | 3:590 |
| i  | <b>Persicaria lapathifolia</b> (L.) S.F.Gray, Nat. Arr. Brit. Pl. 2: 270 (1821)                       | 3:591 |
| i  | <b>Persicaria maculosa</b> S.F.Gray, Nat. Arr. Brit. Pl. 2: 269 (1821)                                | 3:591 |
|    | <b>Persicaria praetermissa</b> (Hook.f.) H.Hara, Fl. E. Himal. 73 (1966)                              | 3:589 |
| ?i | <b>Persicaria prostrata</b> (R.Br.) Soják, Preslia 46: 154 (1974)                                     | 3:590 |
|    | <b>Persicaria subsessilis</b> (R.Br.) K.L.Wilson, Telopea 3: 180 (1988)                               | 3:591 |
| i  | <b>Polygonum arenastrum</b> Jord. ex Boreau, Fl. Cent. Fr. ed.3 2: 559 (1857)                         | 3:589 |
| i  | <b>Polygonum aviculare</b> L., Sp. Pl. 1: 362 (1753)  | 3:588 |
|    | <b>Polygonum convolvulus</b> L. = <b>Fallopia convolvulus</b>   | 3:591 |
|    | <b>Polygonum decipiens</b> R.Br. = <b>Persicaria decipiens</b>  | 3:590 |
|    | <b>Polygonum hydropiper</b> L. = <b>Persicaria hydropiper</b>   | 3:590 |

|                          |  |       |
|--------------------------|--|-------|
|                          | Polygonum lapathifolium L. = <b>Persicaria lapathifolia</b>                      | 3:591 |
|                          | Polygonum persicaria L. = <b>Persicaria maculosa</b>                             | 3:591 |
|                          | Polygonum plebeium sensu W.M.Curtis (1967) = <b>Polygonum arenastrum</b>         | 3:589 |
|                          | Polygonum prostratum R.Br. = <b>Persicaria prostrata</b>                         | 3:590 |
|                          | Polygonum strigosum sensu W.M.Curtis (1967) = <b>Persicaria praetermissa</b>     | 3:589 |
|                          | Polygonum subsessile R.Br. = <b>Persicaria subsessilis</b>                       | 3:591 |
|                          | Reynoutria japonica Houtt. = <b>Fallopia japonica</b>                            |       |
|                          | Rumex acetosella L. = <b>Acetosella vulgaris</b>                                 | 3:587 |
|                          | Rumex angiocarpus Murb. = <b>Acetosella vulgaris</b>                             | 3:586 |
|                          | <b>Rumex bidens</b> R.Br., Prodr. 421 (1810)                                     | 3:586 |
|                          | <b>Rumex brownii</b> Campderá, Monogr. Rumex 64: 81 (1819)                       | 3:585 |
| i                        | <b>Rumex conglomeratus</b> Murray, Prodr. Desig. Stirp. Gott. 52 (1770)          | 3:585 |
| i                        | <b>Rumex crispus</b> L., Sp. Pl. 1: 335 (1753)                                   | 3:584 |
|                          | Rumex dumosiformis Rech. = <b>Rumex dumosus</b>                                  | 3:586 |
|                          | <b>Rumex dumosus</b> A.Cunn. ex Meisn. in DC., Prodr. 14: 62 (1856)              | 3:586 |
|                          | var. dumosiformis (Rech.f.) Rech.f. = <b>Rumex dumosus</b>                       | 3:586 |
| i                        | <b>Rumex obtusifolius</b> L., Sp. Pl. 1: 335 (1753)                              | 3:585 |
| i                        | subsp. <b>obtusifolius</b>   |       |
| i                        | subsp. <b>transiens</b> (Simonkai) Rech.f., Beih. Bot. Cent. 49(2): 52 (1932)    |       |
| i                        | <b>Rumex pulcher</b> L., Sp. Pl. 1: 336 (1753)                                   | 3:585 |
| i                        | subsp. <b>pulcher</b>  |       |
|                          | Rumex sagittatus Thunb. = <b>Acetosa sagittata</b>                               |       |
| <br><b>PORTULACACEAE</b> |  |       |
|                          | <b>Calandrinia calyptрата</b> Hook.f., Icon. Pl. 3: t.296 (1840)                 | 1:78  |
|                          | Calandrinia caulescens sensu Curtis & Morris (1975) = <b>Calandrinia ciliata</b> | 1:79  |
| ni                       | <b>Calandrinia ciliata</b> (Ruiz & Pav.) DC., Prodr. 3: 359 (1828)               | 1:79  |
|                          | <b>Calandrinia eremaea</b> Ewart, Fl. Vict. 486 (1931)                           |       |
|                          | <b>Calandrinia granulifera</b> Benth., Fl. Austral. 1: 176 (1863)                | 1:78  |
|                          | Calandrinia menziesii sensu Census (1999) = <b>Calandrinia ciliata</b>           | 1:79  |
|                          | Calandrinia neesiana H.Eichler = <b>Calandrinia granulifera</b>                  | 1:78  |
|                          | Claytonia australasica Hook.f. = <b>Neopaxia australasica</b>                    | 1:79  |
|                          | Montia australasica (Hook.f.) Pax & Hoffm. = <b>Neopaxia australasica</b>        | 1:79  |
|                          | <b>Montia fontana</b> L., Sp. Pl. 1: 87 (1753)                                   | 1:79  |
|                          | subsp. <b>chondrosperma</b> (Fenzl) Walters, Watsonia 3: 4 (1953)                |       |
|                          | <b>Neopaxia australasica</b> (Hook.f.) Ö.Nilss., Grana Palyn. 7: 331 (1967)      | 1:79  |
|                          | <b>Portulaca oleracea</b> L., Sp. Pl. 1: 445 (1753)                              | 1:78  |
| <br><b>PRIMULACEAE</b>   |  |       |
| i                        | <b>Anagallis arvensis</b> L., Sp. Pl. 1: 148 (1753)                              | 3:467 |
|                          | subsp. foemina sensu W.M.Curtis (1967) = var. <b>coerulea</b>                    | 3:468 |
| i                        | var. <b>arvensis</b>   | 3:468 |
| i                        | var. <b>coerulea</b> Gouan, Fl. Monsp. 30 (1764)                                 | 3:468 |
| i                        | <b>Lysimachia nummularia</b> L., Sp. Pl. 1: 148 (1753)                           | 3:467 |

|                   |   |       |
|-------------------|---|-------|
|                   | <b>Samolus repens</b> (Forst. & Forst.f.) Pers., Syn. Pl. 1: 171 (1805)                 | 3:468 |
| <b>PROTEACEAE</b> |   |       |
| e                 | <b>Agastachys odorata</b> R.Br., Trans. Linn. Soc. Bot. 10: 158 (1810)                  | 3:601 |
| x                 | <b>Banksia integrifolia</b> L.f., Suppl. Pl. 127 (1782)                                 | 3:616 |
|                   | var. <b>integrifolia</b>  |       |
|                   | <b>Banksia marginata</b> Cav., Anales Hist. Nat. 1(3): 227, t.13 (1800)                 | 3:615 |
|                   | <b>Banksia serrata</b> L.f., Suppl. Pl. 126 (1782)                                      | 3:616 |
| e                 | <b>Bellenden montana</b> R.Br., Trans. Linn. Soc. Bot. 10: 166 (1810)                   | 3:600 |
| e                 | <b>Cenarrhenes nitida</b> Labill., Nov. Holl. Pl. 1: 36, t.50 (1805)                    | 3:601 |
| e                 | <b>Conospermum hookeri</b> (Meisn.) E.M.Bennett, Flora Australia 16: 485 (1995)         | 3:600 |
|                   | Conospermum taxifolium Gaertner (Tasmanian plants) = <b>Conospermum hookeri</b>         | 3:600 |
|                   | <b>Grevillea australis</b> R.Br., Trans. Linn. Soc. Bot. 10: 171 (1810)                 | 3:606 |
|                   | var. <b>australis</b>   |       |
| e                 | var. <b>brevifolia</b> Hook.f., Hook. Lond. J. Bot. 6: 282 (1847)                       | 3:606 |
| e                 | var. <b>erecta</b> Hook.f., Hook. Lond. J. Bot. 6: 282 (1847)                           | 3:606 |
| e                 | var. <b>linearifolia</b> Hook.f., Hook. Lond. J. Bot. 6: 282 (1847)                     | 3:606 |
| e                 | var. <b>montana</b> Hook.f., Hook. Lond. J. Bot. 6: 282 (1847)                          | 3:606 |
| e                 | var. <b>planifolia</b> Hook.f., Hook. Lond. J. Bot. 6: 282 (1847)                       | 3:606 |
| e                 | var. <b>subulata</b> Hook.f., Hook. Lond. J. Bot. 6: 282 (1847)                         | 3:606 |
| e                 | var. <b>tenuifolia</b> (R.Br.) Meisn. in DC., Prodr. 14: 360 (1856)                     | 3:607 |
| ni                | <b>Grevillea</b> hybrids  |       |
|                   | <b>Hakea decurrens</b> R.Br., Suppl. Prodr. 27 (1830)                                   |       |
|                   | subsp. <b>physocarpa</b> W.R.Barker, J. Adelaide Bot. Gard. 17: 193 (1996)              | 3:610 |
|                   | subsp. <b>platyaenia</b> W.R.Barker, J. Adelaide Bot. Gard. 17: 196 (1996)              |       |
| e                 | <b>Hakea epiglottis</b> Labill., Nov. Holl. Pl. 1: 30, t.40 (1805)                      | 3:609 |
| e                 | subsp. <b>epiglottis</b>  |       |
| e                 | subsp. <b>milliganii</b> (Meisn.) R.M.Barker, Fl. Australia 17B: 394 (1999)             |       |
|                   | <b>Hakea lissosperma</b> R.Br., Trans. Linn. Soc. Bot. 10: 180 (1810)                   | 3:611 |
| e                 | <b>Hakea megadenia</b> R.M.Barker in Banks et al., Aspects Tasm. Bot. 83 (1991)         |       |
|                   | <b>Hakea microcarpa</b> R.Br., Trans. Linn. Soc. Bot. 10: 182 (1810)                    | 3:611 |
|                   | <b>Hakea nodosa</b> R.Br., Trans. Linn. Soc. Bot. 10: 179 (1810)                        | 3:610 |
|                   | Hakea rostrata sensu W.M.Curtis (1967) = <b>Hakea epiglottis</b> & <b>H. megadenia</b>  | 3:609 |
|                   | Hakea rugosa sensu W.M.Curtis (1967) = <b>Hakea epiglottis</b> & <b>H. megadenia</b>    | 3:609 |
|                   | Hakea sericea sensu W.M.Curtis (1967) = <b>Hakea decurrens</b> subsp. <b>physocarpa</b> | 3:610 |
|                   | <b>Hakea teretifolia</b> (Salisb.) Britten, J. Bot. 54: 60 (1916)                       | 3:609 |
|                   | subsp. <b>hirsuta</b> (Endl.) R.M.Barker, J. Adelaide Bot. Gard. 13: 105 (1990)         |       |
|                   | <b>Hakea ulicina</b> R.Br., Prodr. Suppl. 1: 29 (1830)                                  | 3:611 |
|                   | <b>Isopogon ceratophyllus</b> R.Br., Trans. Linn. Soc. Bot. 10: 72 (1810)               | 3:599 |
| e                 | <b>Lomatia polymorpha</b> R.Br., Trans. Linn. Soc. Bot. 10: 200 (1810)                  | 3:613 |
| e                 | <b>Lomatia tasmanica</b> W.M.Curtis, Stud. Fl. Tasm. 3: 651 (1967)                      | 3:614 |
| e                 | <b>Lomatia tinctoria</b> (Labill.) R.Br., Trans. Linn. Soc. Bot. 10: 199 (1810)         | 3:613 |
| e                 | <b>Orites acicularis</b> (R.Br.) Roemer & Schultes, Syst. Veg. 3: 427 (1818)            | 3:605 |
| e                 | <b>Orites diversifolia</b> R.Br., Trans. Linn. Soc. Bot. 10: 190 (1810)                 | 3:604 |

|                          |  |       |
|--------------------------|--|-------|
| e                        | <b>Orites milliganii</b> Meisn. in DC., Prodr. 14: 424 (1856)  | 3:604 |
| e                        | <b>Orites revoluta</b> R.Br., Trans. Linn. Soc. Bot. 10: 190 (1810)  | 3:605 |
| e                        | <b>Persoonia gunnii</b> Hook.f., Hook. Lond. J. Bot. 6: 283 (1847)   | 3:603 |
| e                        | var. <b>gunnii</b>   |       |
| e                        | var. <b>oblanceolata</b> Orch., Brunonia 6: 236 (1984)   |       |
|                          | <b>Persoonia juniperina</b> Labill., Nov. Holl. Pl. 1: 33, t.45 (1805)   | 3:603 |
| e                        | var. <b>brevifolia</b> Meisn. in DC., Prodr. 14: 336 (1856)  |       |
|                          | var. <b>juniperina</b>   |       |
| e                        | var. <b>mollis</b> Orch., Brunonia 6: 225 (1984)   |       |
| e                        | var. <b>ulicina</b> Meisn. in DC., Prodr. 14: 336 (1856)   |       |
| e                        | <b>Persoonia moscalii</b> Orch., Brunonia 6: 238, fig.12 (1984)  |       |
| e                        | <b>Persoonia muelleri</b> (P.Parm.) Orch., Brunonia 6: 226 (1984)  |       |
| e                        | subsp. <b>angustifolia</b> (Benth.) L.A.S.Johnson & P.H.Weston, Fl. Australia 16: 472 (1995)                             | 3:603 |
| e                        | subsp. <b>densifolia</b> (Orch.) L.A.S.Johnson & P.H.Weston, Fl. Australia 16: 472 (1995)                                |       |
| e                        | subsp. <b>muelleri</b>   |       |
|                          | <i>Persoonia muelleri</i> var. <i>angustifolia</i> (Benth.) Orch. = <b>Persoonia muelleri</b> subsp. <b>angustifolia</b> |       |
|                          | <i>Persoonia muelleri</i> var. <i>densifolia</i> Orch. = <b>Persoonia muelleri</b> subsp. <b>densifolia</b>              |       |
|                          | <i>Persoonia muelleri</i> var. <i>muelleri</i> = <b>Persoonia muelleri</b> subsp. <b>muelleri</b>                        |       |
| e                        | <b>Telopea truncata</b> (Labill.) R.Br., Trans. Linn. Soc. Bot. 10: 198 (1810)   | 3:612 |
| <br><b>RANUNCULACEAE</b> |  |       |
| i#                       | <b>Adonis microcarpa</b> DC., Syst. Nat. 1: 223 (1817)   |       |
| e                        | <b>Anemone crassifolia</b> Hook., Icones Pl. 3: t.257 (1840)   | 1:11  |
| i                        | <b>Batrachium trichophyllum</b> (Chaix) Bosch, Prodr. Fl. Batav. 1: 5 (1850)   | 1:14  |
| e                        | <b>Caltha phylloptera</b> A.W.Hill, Ann. Bot. 32: 433, fig.7 (1918)  | 1:20  |
|                          | <b>Clematis aristata</b> R.Br. ex Ker Gawler, Edwards's Bot. Reg. 3: t.238 (1817)  | 1:9   |
| e                        | <b>Clematis gentianoides</b> DC., Syst. Nat. 1: 159 (1817)   | 1:10  |
|                          | <b>Clematis microphylla</b> DC., Syst. Nat. 1: 147 (1817)  | 1:10  |
| i                        | <b>Clematis vitalba</b> L., Sp. Pl. 1: 544 (1753)  | 1:10  |
| i ?x                     | <b>Myosurus minimus</b> L., Sp. Pl. 1: 284 (1753)  | 1:11  |
| t                        | <b>Ranunculus acaulis</b> Banks & Soland. ex DC., Syst. Nat. 1: 270 (1817)   | 1:17  |
| i                        | <b>Ranunculus acris</b> L., Sp. Pl. 1: 554 (1753)  | 1:18  |
|                          | <b>Ranunculus amphitrichus</b> Colenso, Trans. Proc. N.Z. Inst. 17: 237 (1885)   | 1:16  |
| i#                       | <b>Ranunculus arvensis</b> L., Sp. Pl. 1: 555 (1753)   | 1:18  |
| e                        | <b>Ranunculus collicola</b> Menadue, Brunonia 8(2): 373 (1986)   |       |
|                          | <b>Ranunculus collinus</b> R.Br. ex DC., Syst. Nat. 1: 271 (1817)  | 1:17  |
|                          | <i>Ranunculus concinnus</i> (Hook.f.) Melville = <b>Ranunculus decurvus</b>  | 1:15  |
| e                        | <b>Ranunculus decurvus</b> (Hook.f.) Melville, Kew Bull. 10: 202 (1955)  | 1:15  |
| n                        | <b>Ranunculus diminutus</b> B.G.Briggs, Telopea 5: 583 (1994)  |       |
| i                        | <b>Ranunculus flammula</b> L., Sp. Pl. 1: 548 (1753)   |       |
|                          | <b>Ranunculus glabrifolius</b> Hook., J. Bot. (Hooker) 1: 243 (1834)   | 1:17  |
|                          | <b>Ranunculus gunnianus</b> Hook., J. Bot. (Hooker) 1: 244, t.133 (1834)   | 1:14  |
|                          | <i>Ranunculus inundatus</i> sensu Curtis & Morris (1975) = <b>Ranunculus amphitrichus</b>                                | 1:17  |
| e                        | <b>Ranunculus jugosus</b> Menadue, Brunonia 8(2): 377 (1986)   |       |

|                       |  |       |
|-----------------------|--|-------|
|                       | <b>Ranunculus lappaceus</b> Smith in Rees, Cycl. 29, Ranunculus no.61 (1814)                 | 1:14  |
| i                     | <b>Ranunculus muricatus</b> L., Sp. Pl. 1: 555 (1753)  | 1:18  |
| e                     | <b>Ranunculus nanus</b> Hook., J. Bot. (Hooker) 1: 242 (1834)                                | 1:16  |
| i                     | <b>Ranunculus parviflorus</b> L., Syst. Nat. ed.10: 1087 (1759)                              | 1:19  |
| e                     | <b>Ranunculus pascuinus</b> (Hook.f.) Melville, Kew Bull. 10: 198 (1955)                     | 1:15  |
|                       | <b>Ranunculus pimpinellifolius</b> Hook., J. Bot. (Hooker) 1: 243 (1834)                     | 1:14  |
| e                     | <b>Ranunculus prasinus</b> Menadue, Brunonia 8: 375 (1985)                                   |       |
|                       | <b>Ranunculus pumilio</b> R.Br. ex DC., Syst. Nat. 1: 271 (1817)                             | 1:19  |
|                       | var. <b>pumilio</b>  |       |
| i                     | <b>Ranunculus repens</b> L., Sp. Pl. 1: 554 (1753)   | 1:18  |
|                       | <b>Ranunculus rivularis</b> sensu Curtis & Morris (1975) = <b>Ranunculus amphitrichus</b>    | 1:16  |
| i                     | <b>Ranunculus sardous</b> Crantz, Stirp. Austr. fasc.2: 84 (1763)                            | 1:19  |
|                       | <b>Ranunculus scapiger</b> Hook., J. Bot. (Hooker) 1: 224 (1834)                             | 1:15  |
| i#                    | <b>Ranunculus sceleratus</b> L., Sp. Pl. 1: 551 (1753)                                       | 1:20  |
|                       | <b>Ranunculus sessiliflorus</b> R.Br. ex DC., Syst. Nat. 1: 302 (1817)                       | 1:19  |
|                       | var. <b>sessiliflorus</b>  |       |
| e                     | <b>Ranunculus setaceus</b> Rodway, Pap. Proc. R. Soc. Tasm. for 1900-1901: 107 (1902)        | 1:14  |
|                       | <b>Ranunculus trichophyllus</b> Chaix = <b>Batrachium trichophyllum</b>                      | 1:14  |
| i                     | <b>Ranunculus trilobus</b> Desf., Fl. Atlant. 1: 437, t.113 (1798)                           | 1:19  |
| e                     | <b>Ranunculus triplodontus</b> Melville, Kew Bull. 10: 204 (1955)                            | 1:16  |
| <br><b>RESEDACEAE</b> |  |       |
| i                     | <b>Reseda alba</b> L., Sp. Pl. 1: 449 (1753)   | 1:53  |
| i                     | <b>Reseda lutea</b> L., Sp. Pl. 1: 449 (1753)  | 1:52  |
| i                     | <b>Reseda luteola</b> L., Sp. Pl. 1: 448 (1753)  | 1:52  |
| <br><b>RHAMNACEAE</b> |  |       |
| e                     | <b>Cryptandra alpina</b> Hook.f., Fl. Tasm. 1: 75, t.12B (1855)                              | 1:121 |
|                       | <b>Cryptandra amara</b> Smith in Rees, Cycl. 10: no.2 (1808)                                 | 1:120 |
| e                     | <b>Cryptandra exilis</b> D.I.Morris in Banks et al., Aspects Tasm. Bot. 57 (1991)            |       |
|                       | <b>Cryptandra tomentosa</b> sensu Census Vasc. Pl. Tasm. edn.1 = <b>Cryptandra exilis</b>    |       |
|                       | <b>Discaria pubescens</b> (Brongn.) Druce, Bot. Soc. Exch. Club Brit. Isles 1916: 620 (1917) | 1:121 |
|                       | <b>Pomaderris affinis</b> N.A.Wakef. = <b>Pomaderris intermedia</b>                          | 1:114 |
|                       | <b>Pomaderris apetala</b> Labill., Nov. Holl. Pl. 1: 62, t.87 (1804)                         | 1:115 |
|                       | subsp. <b>apetala</b>  |       |
|                       | subsp. <b>maritima</b> N.G.Walsh & F.Coates, Muellera 10: 34 (1997)                          |       |
|                       | <b>Pomaderris aspera</b> Sieber ex DC., Prodr. 2: 33 (1825)                                  |       |
|                       | <b>Pomaderris elachophylla</b> F.Muell., Fragm. 2: 131 (1861)                                | 1:116 |
|                       | <b>Pomaderris elliptica</b> Labill., Nov. Holl. Pl. 1: 61, t.86 (1804)                       | 1:114 |
| e                     | var. <b>diemenica</b> N.G.Walsh & F.Coates, Muellera 10: 51 (1997)                           |       |
|                       | var. <b>elliptica</b>  |       |
|                       | <b>Pomaderris intermedia</b> Sieber ex DC., Prodr. 2: 33 (1825)                              |       |
|                       | <b>Pomaderris oraria</b> F.Muell. ex Reisseck, Linnaea 29: 268 (1858)                        | 1:116 |
|                       | subsp. <b>oraria</b>   |       |

- Pomaderris paniculosa** F.Muell. ex Reisseck, *Linnaea* 29: 269 (1858)  
 subsp. **paralia** N.G.Walsh, *Muelleria* 7: 276 (1990)
- Pomaderris phyllicifolia** Lodd. ex Link, *Enum. Hort. Berol.* 1: 232 (1821) 1:116  
 subsp. **phyllicifolia**
- Pomaderris pilifera** N.A.Wakef., *Vict. Nat.* 68: 140 (1951) 1:115
- Pomaderris racemosa** Hook., *J. Bot.* 1: 256 (1834-35) 1:116
- i **Rhamnus alaternus** L., *Sp. Pl.* 1: 193 (1753)
- Spyridium eriocephalum** Fenzl in Endl. et al., *Enum. Pl.* 24 (1837) 1:119  
 var. **eriocephalum**
- e **Spyridium gunnii** (Hook.f.) Benth., *Fl. Austral.* 1: 429 (1863) 1:118
- e **Spyridium lawrencei** (Hook.f.) Benth., *Fl. Austral.* 1: 430 (1863) 1:119  
*Spyridium microphyllum* (F.Muell. ex Reisseck) Druce = **Spyridium lawrencei** 1:119
- e **Spyridium obcordatum** (Hook.f.) W.M.Curtis, *Vict. Nat.* 87: 251 (1970) 1:117
- e **Spyridium obovatum** (Hook.) Benth., *Fl. Austral.* 1: 429 (1863) 1:118  
 e var. **obovatum** 1:118  
 e var. **velutinum** (Reisseck) Benth., *Fl. Austral.* 1: 429 (1863) 1:118
- Spyridium parvifolium** (Hook.) F.Muell., *Fragm.* 3: 79 (1862) 1:118
- e var. **molle** (Hook.f.) Benth., *Fl. Austral.* 1: 428 (1863) 1:118  
 var. **parvifolium** 1:118
- e **Spyridium ulicinum** (Hook.) Benth., *Fl. Austral.* 1: 434 (1863) 1:119
- Spyridium vexilliferum** (Hook.) Reisseck, *Linnaea* 29: 285 (1858) 1:119  
 var. **vexilliferum**
- e **Stenanthemum pimeleoides** (Hook.f.) Benth., *Fl. Austral.* 1: 436 (1863) 1:120
- ROSACEAE**
- Acaena agnipila** Gand., *Bull. Soc. Bot. France* 59: 706 (1912) 1:176  
 var. **aequispina** Orch., *Trans. R. Soc. S.Austral.* 93: 96 (1969)  
 var. **tenuispica** (Bitt.) Orch., *Trans. R. Soc. S.Austral.* 93: 95 (1969)
- Acaena Xanserovina** Orch., *Trans. R. Soc. S.Austral.* 93: 104 (1969) 1:176
- Acaena echinata** Nees in Lehm., *Pl. Preiss.* 1: 95 (1844) 1:175  
 var. **retrorsumpilosa** (Bitt.) Orch., *Trans. R. Soc. S.Austral.* 93: 101 (1969)  
 var. **subglabricalyx** (Bitt.) Orch., *Trans. R. Soc. S.Austral.* 93: 99 (1969)  
 var. **tylacantha** Orch., *Trans. R. Soc. S.Austral.* 93: 101 (1969)
- e **Acaena montana** Hook.f., *Lond. J. Bot.* 6: 476 bis (1847) 1:175
- Acaena novae-zelandiae** Kirk, *Trans. Proc. N.Z. Inst.* 3: 177 (1871) 1:175
- Acaena ovina** A.Cunn. in Field, *Geog. Mem. N.S.W.* 358 (1825) 1:175  
 var. **velutina** Orch., *Trans. R. Soc. S.Austral.* 93: 103 (1969)
- Acaena pallida** (Kirk) Allan, *Fl. New Zealand* 1: 360 (1961)
- i **Aphanes arvensis** L., *Sp. Pl.* 1: 123 (1753) 1:174
- Aphanes australiana** (Rothm.) Rothm., *Kew Bull.* 1938: 270 (1838)  
*Aphanes inexpectata* sensu Census 1999 = **Aphanes microcarpa**
- i **Aphanes microcarpa** (Boiss. & Reut.) Rothm., *Repert. Spec. Nov. Regni Veg.* 42: 172 (1937)
- i **Cotoneaster franchetii** Bois, *Rev. Hort.* 1902: 379 (1902)
- i **Cotoneaster glaucophyllus** Franchet, *Pl. Delav.* 222 (1890)

|    |   |       |
|----|---|-------|
| i  | <b>Cotoneaster pannosus</b> Franchet, Pl. Delav. 223 (1890)                   |       |
| ni | <b>Cotoneaster simonsii</b> Baker, Refug. Bot. 1:t.55 (1869)                  |       |
| i  | <b>Crataegus monogyna</b> Jacq., Fl. Austriac. 3: 50 (1775)                   | 1:178 |
|    | <i>Geum renifolium</i> F.Muell. = <b>Geum talbotianum</b>                     |       |
| e  | <b>Geum talbotianum</b> W.M.Curtis, Rec. Q. Vic. Mus. 50: 4 (1974)            | 1:172 |
| x  | <b>Geum urbanum</b> L., Sp. Pl. 1: 501 (1753)                                 | 1:172 |
|    | var. <b>strictum</b> (Aiton) Hook.f., Fl. Tasm. 1: 114 (1856)                 | 1:172 |
| i  | <b>Malus Xdomestica</b> Borkh., Theor. Prakt. Handb. Forstbot. 2: 1272 (1803) |       |
| i  | <b>Potentilla anglica</b> Laich., Veg. Eur. 1: 475 (1790)                     | 1:173 |
| i  | <b>Potentilla anserina</b> L., Sp. Pl. 1: 494 (1753)                          | 1:173 |
| i  | <b>Potentilla recta</b> L., Sp. Pl. 1: 497 (1753)                             | 1:173 |
| i  | <b>Potentilla reptans</b> L., Sp. Pl. 1: 499 (1753)                           | 1:173 |
|    | <i>Poterium polygamum</i> Waldst. & Kit. = <b>Sanguisorba minor</b>           | 1:176 |
| i  | <b>Prunus domestica</b> L., Sp. Pl. 1: 475 (1753)                             | 1:178 |
| i  | subsp. <b>insititia</b> (L.) C.Schneider, Illus. Hand. Bot. (1905)            | 1:178 |
| i  | <b>Prunus spinosa</b> L., Sp. Pl. 1: 475 (1753)                               | 1:178 |
| i  | <b>Rosa canina</b> L., Sp. Pl. 1: 491 (1753)                                  | 1:177 |
| i  | <b>Rosa rubiginosa</b> L., Mant. Pl. 2: 564 (1771)                            | 1:177 |
| i  | <b>Rubus fruticosus</b> [aggregate] L., Sp. Pl. 1: 493 (1753)                 | 1:171 |
| e  | <b>Rubus gunnianus</b> Hook., Icon. Pl. 3: t.291 (1840)                       | 1:171 |
| i  | <b>Rubus laciniatus</b> Willd., Hort. Berol. 2: 82 (1806)                     | 1:171 |
|    | <b>Rubus parvifolius</b> L., Sp. Pl. 2: 1197 (1753)                           | 1:171 |
| i  | <b>Sanguisorba minor</b> Scop., Fl. Carn. ed.2 1: 110 (1772)                  | 1:176 |

## RUBIACEAE

|   |   |       |
|---|---|-------|
|   | <i>Asperula charophyton</i> Airy Shaw & Turrill. Attributed to Tasmania in error              | 2:276 |
|   | <b>Asperula conferta</b> Hook.f., Hook. Lond. J. Bot. 6: 464 bis (1847)                       | 2:275 |
|   | var. <b>abbreviata</b> Airy Shaw & Turrill, Bull. Misc. Inform. Kew 87 (1928)                 | 2:275 |
|   | var. <b>conferta</b>  |       |
|   | var. <b>scoparioides</b> Airy Shaw & Turrill, Bull. Misc. Inform. Kew 87 (1928)               | 2:275 |
|   | <b>Asperula gunnii</b> Hook.f., Hook. Lond. J. Bot. 6: 464 bis (1847)                         | 2:275 |
| e | var. <b>curta</b> (Hook.f.) Airy Shaw & Turrill, Bull. Misc. Inform. Kew 90 (1928)            | 2:276 |
|   | var. <b>gunnii</b>  |       |
|   | <b>Asperula minima</b> Hook.f., Hook. Lond. J. Bot. 6: 464 bis (1847)                         | 2:276 |
|   | <b>Asperula pusilla</b> Hook.f., Hook. Lond. J. Bot. 6: 464 bis (1847)                        | 2:276 |
|   | <b>Asperula scoparia</b> Hook.f., Hook. Lond. J. Bot. 6: 463 bis (1847)                       | 2:275 |
|   | var. <b>scoparia</b>  |       |
|   | <b>Asperula subsimplex</b> Hook.f., Hook. Lond. J. Bot. 6: 463 bis (1847)                     | 2:276 |
|   | <b>Coprosma hirtella</b> Labill., Nov. Holl. Pl. 1: 70, t.95 (1805)                           | 2:268 |
|   | <b>Coprosma moorei</b> F.Muell. ex Rodway, Pap. Proc. R. Soc. Tasm. for 1893: 179, t.I (1894) | 2:269 |
|   | <b>Coprosma nitida</b> Hook.f., Hook. Lond. J. Bot. 6: 465 bis (1847)                         | 2:269 |
|   | <b>Coprosma perpusilla</b> Colenso, Trans. N.Z. Inst. 22: 466 (1890)                          |       |
|   | subsp. <b>perpusilla</b>  |       |
|   | <b>Coprosma pumila</b> Hook.f., Fl. Antart. 543 (1847)  | 2:269 |



|                     |  |       |
|---------------------|--|-------|
|                     | <b>Coprosma quadrifida</b> (Labill.) Robinson, Proc. Amer. Acad. Arts 45: 409 (1910)       | 2:268 |
| i                   | <b>Coprosma repens</b> A.Rich., Essai Fl. N.Z. 264 (1832)                                  |       |
| i                   | <b>Coprosma robusta</b> Raoul, Ann. Sci. Nat. Bot. ser.3 2: 121 (1844)                     |       |
|                     | <b>Galium albescens</b> Hook.f. = <b>Galium australe</b>                                   | 2:273 |
| i                   | <b>Galium aparine</b> L., Sp. Pl. 1: 108 (1753)  | 2:273 |
|                     | <b>Galium australe</b> DC., Prodr. 4: 608 (1830)   | 2:273 |
|                     | <b>Galium ciliare</b> Hook.f., Hook. Lond. J. Bot. 6: 461 bis (1847)                       | 2:272 |
| i                   | <b>Galium divaricatum</b> Pourret ex Lam., Encycl. 2: 580 (1788)                           |       |
|                     | <b>Galium gaudichaudii</b> DC., Prodr. 4: 607 (1830)                                       | 2:272 |
| i                   | <b>Galium murale</b> (L.) All., Fl. Pedem. 1: 8 (1785)                                     | 2:273 |
| it                  | <b>Galium palustre</b> L., Sp. Pl. 1: 105 (1753)   |       |
| i                   | <b>Galium verum</b> L., Sp. Pl. 1: 107 (1753)  |       |
|                     | <i>Nertera depressa</i> Banks & Soland. ex Gaertner = <b>Nertera granadensis</b>           | 2:270 |
|                     | <b>Nertera granadensis</b> (L.f.) Druce, Bot. Soc. Exch. Club Brit. Isles 1916: 637 (1917) | 2:270 |
|                     | <b>Opercularia ovata</b> Hook.f., Hook. Lond. J. Bot. 6: 465 bis (1847)                    | 2:270 |
|                     | <b>Opercularia varia</b> Hook.f., Hook. Lond. J. Bot. 6: 466 bis (1847)                    | 2:271 |
| i                   | <b>Sherardia arvensis</b> L., Sp. Pl. 1: 102 (1753)  | 2:277 |
| <br><b>RUTACEAE</b> |  |       |
| e                   | <b>Acradenia frankliniae</b> Milligan ex Kippist, Proc. Linn. Soc. Lond. 2: 201 (1853)     | 1:106 |
|                     | <b>Boronia anemonifolia</b> A.Cunn. in Field, Geogr. Mem. N.S.W. 330 (1825)                | 1:105 |
|                     | subsp. <b>variabilis</b> (Hook.) P.G.Neish, Muelleria 14: 11 (2000)                        |       |
| e                   | <b>Boronia citriodora</b> Gunn ex Hook.f., Fl. Tasm. 1: 68 (1855)                          | 1:105 |
| e                   | subsp. <b>citriodora</b>   |       |
| e                   | subsp. <b>orientalis</b> Duretto, Muelleria 17: 83 (2003)                                  |       |
| e                   | subsp. <b>paulwilsonii</b> Duretto, Muelleria 17: 81 (2003)                                |       |
| e                   | <b>Boronia elisabethiae</b> Duretto, Muelleria 17: 88 (2003)                               |       |
| e                   | <b>Boronia gunnii</b> Hook.f., Fl. Tasm. 1: 68, t.10 (1855)                                |       |
| e                   | <b>Boronia hemichiton</b> Duretto, Muelleria 17: 87 (2003)                                 |       |
| e                   | <b>Boronia hippopala</b> Duretto, Muelleria 17: 84 (2003)                                  |       |
|                     | <b>Boronia nana</b> Hook., Icon. Pl. 3: t.270 (1840)                                       | 1:105 |
|                     | var. <b>hyssopifolia</b> Melville, Kew Bull. 9: 463 (1954)                                 |       |
|                     | var. <b>nana</b>   |       |
|                     | <b>Boronia parviflora</b> Smith, Tracts Nat. Hist. 295: t.6 (1798)                         | 1:105 |
|                     | <b>Boronia pilosa</b> Labill., Nov. Holl. Pl. 1: 97, t.124 (1805)                          | 1:104 |
|                     | subsp. <b>pilosa</b>   | 1:105 |
| e                   | subsp. <b>tasmanensis</b> Duretto, Muelleria 17: 97 (2003)                                 |       |
|                     | var. <i>floribunda</i> Hook. = <b>Boronia pilosa</b> subsp. <b>pilosa</b>                  | 1:105 |
|                     | var. <i>laricifolia</i> (Hook.) Hook.f. = <b>Boronia pilosa</b> subsp. <b>pilosa</b>       | 1:105 |
|                     | <b>Boronia rhomboidea</b> Hook., Icon. Pl. 8: t.722 (1845)                                 | 1:105 |
| e                   | <b>Boronia rozefeldsii</b> Duretto, Muelleria 17: 101 (2003)                               |       |
|                     | <b>Correa alba</b> Andrews, Bot. Repos. 1: t.18 (1798)                                     | 1:109 |
|                     | var. <b>alba</b>   |       |
|                     | <b>Correa backhouseana</b> Hook., J. Bot. 1: 253 (1834)                                    | 1:110 |

|                        |  |       |
|------------------------|--|-------|
|                        | var. <b>backhouseana</b>   |       |
|                        | <b>Correa lawrenceana</b> Hook., J. Bot. 1: 254 (1834)                                       | 1:111 |
| e                      | var. <b>lawrenceana</b>  | 1:111 |
|                        | <b>Correa reflexa</b> (Labill.) Vent., Jard. Malmaison 1: t.13 (1803)                        | 1:110 |
|                        | var. <b>nummulariifolia</b> (Hook.f.) Paul G.Wilson, Trans. R. Soc. S.Austral. 85: 30 (1961) | 1:110 |
|                        | var. <b>reflexa</b>  | 1:110 |
|                        | Eriostemon verrucosus A.Rich. = <b>Philotheca verrucosa</b>                                  | 1:107 |
|                        | Eriostemon virgatus Hook.f. = <b>Philotheca virgata</b>                                      | 1:107 |
|                        | <b>Leionema bilobum</b> (Lindley) Paul G.Wilson, Nuytsia 12(2): 271 (1998)                   | 1:108 |
| e                      | <b>Leionema montanum</b> (Hook.) Paul G.Wilson, Nuytsia 12(2): 275 (1998)                    | 1:108 |
| e                      | <b>Leionema oldfieldii</b> (F.Muell.) Paul G.Wilson, Nuytsia 12(2): 276 (1998)               | 1:108 |
|                        | <b>Nematolepis squamea</b> (Labill.) Paul G.Wilson, Nuytsia 12(2): 279 (1998)                | 1:109 |
| e                      | subsp. <b>retusa</b> (Hook.) Paul G.Wilson, Nuytsia 12(2): 279 (1998)                        | 1:109 |
|                        | subsp. <b>squamea</b>  | 1:109 |
|                        | Phebalium bilobum Lindley in Mitchell = <b>Leionema bilobum</b>                              | 1:108 |
| e                      | <b>Phebalium daviesii</b> Hook.f., Fl. Tasm. 2: 358 (1859)                                   | 1:108 |
|                        | Phebalium montanum Hook. = <b>Leionema montanum</b>  | 1:108 |
|                        | Phebalium oldfieldii (F.Muell.) Benth. = <b>Leionema oldfieldii</b>                          | 1:108 |
|                        | Phebalium squameum (Labill.) Engl. in Engler & Prantl = <b>Nematolepis squamea</b>           | 1:109 |
| e                      | <b>Philotheca freyciana</b> Rozefelds, Muelleria 15: 23 (2001)                               |       |
|                        | <b>Philotheca verrucosa</b> (A.Rich.) Paul G.Wilson, Nuytsia 12(2): 260 (1998)               | 1:107 |
|                        | <b>Philotheca virgata</b> (Hook.f.) Paul G.Wilson, Nuytsia 12(2): 260 (1998)                 | 1:107 |
|                        | <b>Zieria arborescens</b> Sims, Bot. Mag. 34: t.1395 (1811)                                  | 1:103 |
|                        | subsp. <b>arborescens</b>  | 1:103 |
|                        | Zieria cytisoides sensu W.M.Curtis (1975) = <b>Zieria littoralis</b>                         | 1:103 |
|                        | <b>Zieria littoralis</b> J.A.Armstrong, Aust. Syst. Bot. 15(3): 389 (2002)                   | 1:103 |
|                        | <b>Zieria veronicea</b> (F.Muell.) Benth., Fl. Austral. 1: 305 (1863)                        | 1:103 |
|                        | subsp. <b>veronicea</b>  | 1:103 |
| <br><b>SALICACEAE</b>  |  |       |
| i                      | <b>Populus alba</b> L., Sp. Pl. 2: 1034 (1753)   |       |
| i                      | <b>Salix alba</b> L., Sp. Pl. 2: 1021 (1753)   |       |
|                        | Salix alba L. x fragilis L. = <b>Salix Xrubens</b>   | 3:648 |
|                        | Salix atrocinerea Brot. = <b>Salix cinerea</b>   | 3:649 |
| i                      | <b>Salix babylonica</b> L., Sp. Pl. 2: 1017 (1753)   | 3:648 |
| i                      | <b>Salix cinerea</b> L., Sp. Pl. 2: 1021 (1753)  | 3:649 |
| i                      | <b>Salix fragilis</b> L., Sp. Pl. 2: 1017 (1753)   |       |
| i                      | <b>Salix Xrubens</b> Schrank, Baiers. Fl. 1: 226 (1789)                                      | 3:648 |
| <br><b>SANTALACEAE</b> |  |       |
|                        | <b>Exocarpos cupressiformis</b> Labill., Voy. Rech. Perouse 1: 155, t.14 (1800)              | 3:628 |
| e                      | <b>Exocarpos humifusus</b> R.Br., Prodr. 356 (1810)  | 3:629 |
|                        | <b>Exocarpos nanus</b> Hook.f., Lond. J. Bot. 6: 281 (1847)                                  | 3:629 |
|                        | <b>Exocarpos strictus</b> R.Br., Prodr. 357 (1810)   | 3:628 |

|                             |  |       |
|-----------------------------|--|-------|
|                             | <b>Exocarpos syrticola</b> (F.Muell. ex Miq.) Stauffer, Mitt. Bot. Mus. Univ. Zurich 213: 173 (1959) | 3:628 |
|                             | <b>Leptomeria drupacea</b> (Labill.) Druce, Bot. Exch. Club Soc. Brit. Isles 1916: 632 (1917)        | 3:626 |
| e                           | <b>Leptomeria glomerata</b> F.Muell. ex Hook.f., Fl. Tasm. 2: 370 (1859)                             | 3:627 |
| x                           | <b>Thesium australe</b> R.Br., Prodr. 353 (1810)   | 3:626 |
| <br><b>SAPINDACEAE</b>      |  |       |
|                             | <i>Dodonaea ericifolia</i> G.Don = <b>Dodonaea filiformis</b>  | 1:122 |
| e                           | <b>Dodonaea filiformis</b> Link, Enum. Hort. Berol. 1: 381 (1821)                                    | 1:122 |
|                             | <b>Dodonaea viscosa</b> Jacq., Enum. Sys. Pl. 19 (1760)  | 1:122 |
|                             | subsp. <b>spatulata</b> (Smith) J.West, Brunonia 7: 43 (1984)  |       |
| <br><b>SCROPHULARIACEAE</b> |  |       |
| i                           | <b>Bellardia trixago</b> (L.) All., Fl. Pedem. 1: 61 (1785)  |       |
|                             | <i>Celsia cretica</i> L. = <b>Verbascum creticum</b>   | 3:513 |
| t                           | <b>Chionohebe ciliolata</b> (Hook.f.) B.G.Briggs & Ehrend., Contr. Herb. Austral. 25: 2 (1976)       |       |
| i                           | <b>Cymbalaria muralis</b> P.Gaertner, B.Meyer & Scherb., Fl. Wett. 2: 397 (1800)                     | 3:514 |
|                             | <b>Derwentia derwentiana</b> (Andrews) B.G.Briggs & Ehrend., Telopea 5: 265 (1992)                   | 3:523 |
|                             | subsp. <b>derwentiana</b>  |       |
|                             | <b>Derwentia nivea</b> (Lindley) B.G.Briggs & Ehrend., Telopea 5: 260 (1992)                         | 3:523 |
| i                           | <b>Digitalis purpurea</b> L., Sp. Pl. 2: 621 (1753)  | 3:519 |
| e                           | <b>Euphrasia amphisysepala</b> W.R.Barker, J. Adelaide Bot. Gard. 10(2): 204 (1987)                  |       |
|                             | <i>Euphrasia brownii</i> var. <i>striata</i> (R.Br.) Rodw. = <b>Euphrasia striata</b>                | 3:530 |
|                             | <b>Euphrasia collina</b> R.Br., Prodr. 436 (1810)  | 3:527 |
|                             | subsp. <b>collina</b>  |       |
| e                           | subsp. <b>deflexifolia</b> (Gand.) W.R.Barker in Stones & Curtis, End. Fl. Tasm. 6: 477 (1978)       |       |
| e                           | subsp. <b>diemenica</b> (Sprengel) W.R.Barker in Stones & Curtis, End. Fl. Tasm. 6: 477 (1978)       | 3:528 |
| e                           | subsp. <b>gunnii</b> (Du Rietz) W.R.Barker, J. Adelaide Bot. Gard. 5: 199 (1982)                     | 3:528 |
|                             | subsp. <b>tetragona</b> (R.Br.) W.R.Barker, J. Adelaide Bot. Gard. 5: 189 (1982)                     |       |
|                             | <i>Euphrasia cuspidata</i> Hook.f. = <b>Euphrasia hookeri</b>  | 3:532 |
|                             | <i>Euphrasia diemenica</i> Sprengel = <b>Euphrasia collina</b> subsp. <b>diemenica</b>               | 3:528 |
| e                           | <b>Euphrasia fragosa</b> W.R.Barker, J. Adelaide Bot. Gard. 17: 217 (1996)                           |       |
|                             | <b>Euphrasia gibbsiae</b> Du Rietz, Sv. Bot. Tidskr. 42: 104 (1948)                                  | 3:531 |
| e                           | subsp. <b>comberi</b> (Du Rietz) W.R.Barker, J. Adelaide Bot. Gard. 5: 122 (1982)                    | 3:531 |
| e                           | subsp. <b>discolor</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 128 (1982)                              |       |
| e                           | subsp. <b>gibbsiae</b>   |       |
| e                           | subsp. <b>kingii</b> (W.M.Curtis) W.R.Barker in Stones & Curtis, End. Fl. Tasm. 6: 477 (1978)        | 3:530 |
| e                           | subsp. <b>microdonta</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 126 (1982)                            |       |
| e                           | subsp. <b>psilantherea</b> (F.Muell.) W.R.Barker, J. Adelaide Bot. Gard. 5: 119 (1982)               |       |
| e                           | subsp. <b>pulvinestris</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 131-2 (1982)                        |       |
|                             | subsp. aff. <b>subglabrifolia</b> (Du Rietz) W.R.Barker, J. Adelaide Bot. Gard. 5: 127 (1982)        | 3:531 |
| e                           | subsp. <b>wellingtonensis</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 120 (1982)                       |       |
|                             | <i>Euphrasia gunnii</i> Du Rietz = <b>Euphrasia collina</b> subsp. <b>gunnii</b>                     | 3:528 |
| e                           | <b>Euphrasia hookeri</b> Wettst., Monogr. Gatt. Euphrasia 268 (1896)                                 | 3:532 |
|                             | <i>Euphrasia kingii</i> W.M.Curtis = <b>Euphrasia gibbsiae</b> subsp. <b>kingii</b>                  | 3:530 |

|     |   |       |
|-----|---|-------|
| e   | <b>Euphrasia phragmostoma</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 103 (1982)            |       |
|     | <b>Euphrasia scabra</b> R.Br., Prodr. 437 (1810)  | 3:531 |
| e   | <b>Euphrasia semipicta</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 139 (1982)               |       |
| e   | <b>Euphrasia striata</b> R.Br., Prodr. 436 (1810)   | 3:530 |
|     | <b>Glossostigma elatinoides</b> (Benth.) Benth. ex Hook.f., Fl. Nov.-Zel. 1: 189 (1852)   | 3:518 |
|     | <i>Gratiola latifolia</i> R.Br. = <b>Gratiola peruviana</b>                               | 3:517 |
|     | <b>Gratiola nana</b> Benth. in DC., Prodr. 10: 404 (1846)                                 | 3:517 |
|     | <b>Gratiola peruviana</b> L., Sp. Pl. 1: 17 (1753)  | 3:517 |
|     | <b>Gratiola pubescens</b> R.Br., Prodr. 435 (1810)  | 3:517 |
|     | <b>Gratiola pumilo</b> F.Muell., Linnaea 25: 431 (1853)                                   |       |
| i   | <b>Kickxia elatine</b> (L.) Dumort., Fl. Belg. 35 (1827)                                  | 3:514 |
| i   | <b>Kickxia spuria</b> (L.) Dumort., Fl. Belg. 35 (1827)                                   | 3:514 |
|     | <b>Limosella australis</b> R.Br., Prodr. 443 (1810)                                       | 3:519 |
|     | <i>Limosella lineata</i> Gluck = <b>Limosella australis</b>                               | 3:519 |
| i   | <b>Linaria vulgaris</b> Mill., Gard. Dict. ed.8, no.6 (1768)                              | 3:513 |
|     | <b>Mazus pumilio</b> R.Br., Prodr. 439 (1810)   | 3:516 |
| i   | <b>Mimulus moschatus</b> Douglas ex Lindley, Bot. Reg. 13: t.1118 (1828)                  | 3:516 |
|     | <b>Mimulus repens</b> R.Br., Prodr. 439 (1810)  | 3:515 |
| e   | <b>Ourisia integrifolia</b> R.Br., Prodr. 439 (1810)                                      | 3:520 |
|     | <i>Parahebe derwentiana</i> (Andrews) B.G.Briggs & Ehrend. = <b>Derwentia derwentiana</b> | 3:523 |
| i   | <b>Parentucellia latifolia</b> (L.) Caruel in Parl., Fl. Ital. 6: 480 (1885)              | 3:532 |
| i   | <b>Parentucellia viscosa</b> (L.) Caruel in Parl., Fl. Ital. 6: 482 (1885)                | 3:533 |
|     | <i>Pygmea ciliolata</i> Hook.f. = <b>Chionohebe ciliolata</b>                             |       |
|     | <i>Scrophularia aquatica</i> sensu W.M.Curtis (1967) = <b>Scrophularia auriculata</b>     | 3:515 |
| it  | <b>Scrophularia auriculata</b> L., Sp. Pl. 2: 620 (1753)                                  | 3:515 |
| i   | <b>Verbascum blattaria</b> L., Sp. Pl. 1: 178 (1753)                                      | 3:512 |
| i   | <b>Verbascum creticum</b> (L.) Cav., Elench. Pl. Horti Matr. 39 (1803)                    | 3:513 |
| i   | <b>Verbascum thapsus</b> L., Sp. Pl. 1: 177 (1753)  | 3:512 |
| i   | <b>Verbascum virgatum</b> Stokes in Withering, Arr. Brit. Pl. ed.2 227 (1796)             | 3:512 |
| i   | <b>Veronica arvensis</b> L., Sp. Pl. 1: 13 (1753)   | 3:525 |
|     | <b>Veronica calycina</b> R.Br., Prodr. 435 (1810)   | 3:524 |
|     | <i>Veronica derwentiana</i> Andrews = <b>Derwentia derwentiana</b>                        | 3:523 |
|     | <i>Veronica distans</i> var. <i>pubescens</i> Benth. = <b>Veronica novae-hollandiae</b>   | 3:523 |
| e   | <b>Veronica formosa</b> R.Br., Prodr. 434 (1810)  | 3:522 |
|     | <b>Veronica gracilis</b> R.Br., Prodr. 435 (1810)   | 3:523 |
| i   | <b>Veronica hederifolia</b> L., Sp. Pl. 1: 13 (1753)                                      | 3:525 |
|     | <i>Veronica nivea</i> Lindley = <b>Derwentia nivea</b>                                    | 3:523 |
| ?i# | <b>Veronica notabilis</b> F.Muell. ex Benth., Fl. Austral. 4: 511 (1868)                  | 3:524 |
|     | <b>Veronica novae-hollandiae</b> Poirlet in Lam., Encycl. Méth. Bot. 8: 526 (1808)        | 3:523 |
| i   | <b>Veronica peregrina</b> L., Sp. Pl. 1: 14 (1753)  | 3:525 |
| i   | <b>Veronica persica</b> Poirlet in Lam., Encycl. Méth. Bot. 8: 542 (1808)                 | 3:526 |
|     | <b>Veronica plebeia</b> R.Br., Prodr. 435 (1810)  | 3:524 |
| it  | <b>Veronica scutellata</b> L., Sp. Pl. 1: 12 (1753)                                       |       |
| i   | <b>Veronica serpyllifolia</b> L., Sp. Pl. 1: 12 (1753)                                    | 3:525 |

## SOLANACEAE

- Anthocercis tasmanica Hook.f. = **Cyphanthera tasmanica** 3:509
- e **Cyphanthera tasmanica** Miers, Ann. Mag. Nat. Hist. ser.2, 11: 377 (1853) 3:509
- i **Datura ferox** L., Amoen. Acad. 3: 403 (1756) 3:508
- i **Datura stramonium** L., Sp. Pl. 1: 179 (1753) 3:508
- i# **Hyoscyamus albus** L., Sp. Pl. 1: 180 (1753) 3:509
- Hyoscyamus niger sensu W.M.Curtis (1967) = **Hyoscyamus albus** 3:509
- i **Lycium barbarum** L., Sp. Pl. 1: 192 (1753)
- i **Lycium ferocissimum** Miers, Ann. Mag. Nat. Hist. ser.2, 14: 187 (1854) 3:507
- i **Physalis peruviana** L., Sp. Pl. ed.2, 2: 1670 (1763)
- i **Salpichroa organifolia** (Lam.) Baill., Hist. Pl. 9: 288 (1888) 3:506
- ni# **Solanum americanum** Mill., Gard. Dict. edn.8 no.5 (1768)
- it **Solanum dulcamara** L., Sp. Pl. 1: 185 (1753) 3:504
- i **Solanum furcatum** Dunal ex Poir., Encycl. Suppl. 3: 750 (1814) 3:504
- Solanum laciniatum** Aiton, Hort. Kew. 1: 247 (1789) 3:505
- i **Solanum marginatum** L.f., Suppl. 147 (1781) 3:506
- i **Solanum nigrum** L., Sp. Pl. 1: 186 (1753) 3:504
- Solanum nitidibaccatum** Bitter = **Solanum physalifolium** var. **nitidibaccatum** 3:505
- Solanum opacum** A.Braun & Bouché, Ind. Sem. Hort. Berol. App. 8: 18 no.38 (1853)
- i **Solanum physalifolium** Rusby, Mem. Torrey Bot. Club 6: 88 (1896)
- i var. **nitidibaccatum** (Bitter) Edmonds, Bot. J. Linn. Soc. 92(1): 27 (1986) 3:505
- i **Solanum pseudocapsicum** L., Sp. Pl. 1: 184 (1753) 3:506
- Solanum sarrachoides** Sendtner = **Solanum physalifolium** var. **nitidibaccatum**
- i **Solanum triflorum** Nutt., Gen. N.Amer. Pl. 1: 128 (1818)
- Solanum vescum** F.Muell., Trans. Proc. Vict. Inst. Adv. Sci. 1: 69 (1855) 3:506
- var. **dauidii** Herasim. This is a cultivar name and is deleted from the Census

## STACKHOUSIACEAE

- Stackhousia flava Hook. = **Stackhousia viminea** 1:112
- e **Stackhousia gunnii** Hook.f., Fl. Tasm. 1: 79 (1855) [nom. illeg. but retained here pending resolution] 1:112
- Stackhousia monogyna** Labill., Nov. Holl. Pl. 1: 77 t.104 (1805) 1:112
- Stackhousia pulvinaris** F.Muell., Trans. Phil. Soc. Vict. 1: 101 (1855) 1:112
- Stackhousia spathulata** Sieber ex Sprengel, Syst. Veg. 4: 124 (1827) 1:112
- Stackhousia viminea** Smith in Rees, Cycl. 33 (1816) 1:112

## STERCULIACEAE

- Lasiopetalum baueri** Steetz in Lehm., Pl. Preiss. 2: 339 (1848) 1:90
- Lasiopetalum dasyphyllum** Sieber ex Hook.f. = **Lasiopetalum macrophyllum** 1:89
- Lasiopetalum discolor** Hook., Comp. Bot. Mag. 1: 276 (1835) 1:89
- Lasiopetalum macrophyllum** Graham, Bot. Mag. 68: t.3908 (1841) 1:89
- e **Lasiopetalum micranthum** Hook.f., Fl. Tasm. 1: 51 (1855) 1:90

## STYLIDIACEAE

|                          |  |       |
|--------------------------|--|-------|
| e                        | <b>Forstera bellidifolia</b> Hook., Icon. Pl. 9: t.851 (1851)  | 2:395 |
| x                        | <b>Levenhookia dubia</b> Sonder in Lehm., Pl. Preiss. 1: 392 (1845)  | 2:395 |
| t                        | <b>Phyllachne colensoi</b> (Hook.f.) Bergg., Minn. Kong. Fys. Säll. Lund 8: 11, t.3 figs.1-27 (1877)           | 2:396 |
|                          | <b>Stylidium armeria</b> (Labill.) Labill., Nov. Holl. Pl. 2. 66 (1806) (sensu Jackson & Wiltshire, loc. cit.) |       |
|                          | <i>Stylidium brachyphyllum</i> Sonder = <b>Stylidium inundatum</b>   | 2:394 |
|                          | <b>Stylidium despectum</b> R.Br., Prodr. 571 (1810)  | 2:394 |
|                          | <b>Stylidium dilatatum</b> W.D.Jackson & R.J.E.Wiltshire, Aust. Syst. Bot. 14(6): 961 (2001)                   |       |
|                          | <b>Stylidium graminifolium</b> Swartz ex Willd., Sp. Pl. ed.5, 4: 146 (1805)                                   | 2:393 |
|                          | <b>Stylidium inundatum</b> R.Br., Prodr. 571 (1810)  | 2:394 |
|                          | <b>Stylidium perpusillum</b> Hook.f., London J. Bot. 6: 266 (1847)   | 2:394 |
| <br><b>THYMELAEACEAE</b> |  |       |
|                          | <i>Drapetes tasmanicus</i> Hook.f. = <b>Kelleria dieffenbachii</b>   | 3:625 |
|                          | <b>Kelleria dieffenbachii</b> (Hook.) Endl., Gen. Pl. Suppl. 4(2): 61 (1848)                                   | 3:625 |
|                          | <b>Pimelea axiflora</b> F.Muell. ex Meisn., Linnaea 26: 345 (1854)   | 3:622 |
|                          | subsp. <b>axiflora</b>   |       |
| e                        | <b>Pimelea cinerea</b> R.Br., Prodr. 361 (1810)  | 3:619 |
|                          | <b>Pimelea curviflora</b> R.Br., Prodr. 362 (1810)   | 3:624 |
|                          | var. <i>gracilis</i> (R.Br.) Threlfall = <b>Pimelea curviflora</b>   | 3:624 |
|                          | var. <i>sericea</i> Benth. = <b>Pimelea curviflora</b>   |       |
|                          | <b>Pimelea drupacea</b> Labill., Nov. Holl. Pl. 1: 10 (1805)   | 3:622 |
| e                        | <b>Pimelea filiformis</b> Hook.f., London J. Bot. 6: 280 (1847)  | 3:622 |
|                          | <b>Pimelea flava</b> R.Br., Prodr. 361 (1810)  | 3:623 |
|                          | subsp. <b>flava</b>  |       |
|                          | <b>Pimelea glauca</b> R.Br., Prodr. 360 (1810)   | 3:619 |
|                          | <b>Pimelea humilis</b> R.Br., Prodr. 361 (1810)  | 3:620 |
|                          | <b>Pimelea ligustrina</b> Labill., Nov. Holl. Pl. 1: 9, f.3 (1805)   | 3:620 |
|                          | subsp. <b>ligustrina</b>   |       |
|                          | <i>Pimelea lindleyana</i> Meisn. = <b>Pimelea linifolia</b>  | 3:620 |
|                          | <b>Pimelea linifolia</b> Smith, Bot. Nov. Holl. 31, t.11 (1794)  | 3:620 |
|                          | subsp. <b>linifolia</b>  |       |
|                          | subsp. <b>linoides</b> (Cunn.) Threlfall, Brunonia 5: 142 (1983)   |       |
|                          | <b>Pimelea micrantha</b> F.Muell. ex Meisn., Linnaea 26: 351 (1854)  | 3:624 |
| e                        | <b>Pimelea milliganii</b> Meisn. in A.P. de Candolle, Prodr. 14: 509 (1857)                                    | 3:619 |
| e                        | <b>Pimelea nivea</b> Labill., Nov. Holl. Pl. 1: 10 (1805)  | 3:621 |
|                          | <b>Pimelea pauciflora</b> R.Br., Prodr. 360 (1810)   | 3:622 |
|                          | <b>Pimelea phycoides</b> Meisn. in Lehm., Pl. Preiss. 2: 271 (1848)  |       |
| e                        | <b>Pimelea pygmaea</b> F.Muell. & C.Stuart ex Meisn., Linnaea 26: 346 (1854)                                   | 3:623 |
| e                        | <b>Pimelea sericea</b> R.Br., Prodr. 361 (1810)  | 3:621 |
|                          | <b>Pimelea serpyllifolia</b> R.Br., Prodr. 360 (1810)  | 3:623 |
|                          | subsp. <b>serpyllifolia</b>  |       |
|                          | <i>Pimelea stricta</i> sensu W.M.Curtis (1967) = <b>Pimelea linifolia</b> subsp. <b>linifolia</b>              | 3:624 |
| <br><b>TREMANDRACEAE</b> |  |       |

|                          |   |       |
|--------------------------|---|-------|
|                          | <b>Tetradthea ciliata</b> Lindley in T.Mitch., Three Exped. Int. E. Australia 2: 205 (1838)         | 1:60  |
|                          | Tetradthea glandulosa Labill. = <b>Tetradthea labillardierei</b>                                    | 1:60  |
| e                        | <b>Tetradthea gunnii</b> Hook.f., Fl. Tasm. 1: 36, t.7B (1855)                                      |       |
|                          | <b>Tetradthea labillardierei</b> Joy Thomps., Telopea 1: 189 (1976)                                 | 1:60  |
|                          | <b>Tetradthea pilosa</b> Labill., Nov. Holl. Pl. 1: 95, t.122 (1806)                                | 1:60  |
|                          | subsp. <b>latifolia</b> Joy Thomps., Telopea 1: 213 (1976)  |       |
|                          | subsp. <b>pilosa</b>  |       |
|                          | <b>Tetradthea procumbens</b> Gunn ex Hook.f., Fl. Tasm. 1: 35, t.7A (1855)                          | 1:61  |
| <br><b>ULMACEAE</b>      |   |       |
| i                        | <b>Ulmus Xhollandica</b> Mill., Gard. Dict. ed.8, no.5 (1768)                                       | 3:642 |
|                          | var. <b>hollandica</b>  | 3:642 |
| <br><b>URTICACEAE</b>    |   |       |
|                          | Australina muelleri Wedd. = <b>Australina pusilla</b> subsp. <b>muelleri</b>                        | 3:641 |
|                          | <b>Australina pusilla</b> (Desf. ex Poir.) Gaudich., Voy. Bonite Bot. 3: t.114A (1852)              | 3:641 |
|                          | subsp. <b>muelleri</b> (Wedd.) Friis & Wilmot-Dear in Friis et al., Nord. J. Bot. 7: 126 (1987)     | 3:641 |
|                          | subsp. <b>pusilla</b>   | 3:641 |
|                          | Helxine soleirolii Req. = <b>Soleirolia soleirolii</b>  | 3:640 |
|                          | <b>Parietaria cardiostegia</b> Greuter, Fl. Australia 3: 190 (1989)                                 |       |
|                          | <b>Parietaria debilis</b> Forst.f., Fl. Ins. Austral. Prodr. 73 (1786)                              | 3:640 |
| i                        | <b>Parietaria judaica</b> L., Sp. Pl. edn.2, 2: 1492 (1763)   |       |
| i                        | <b>Soleirolia soleirolii</b> (Req.) Dandy, Feddes Repert. 70: 4 (1965)                              | 3:640 |
|                          | <b>Urtica incisa</b> Poir. in Lam., Encyc. Supp. 4: 224 (1816)                                      | 3:639 |
| i                        | <b>Urtica urens</b> L., Sp. Pl. 2: 984 (1753)   | 3:639 |
| <br><b>VALERIANACEAE</b> |   |       |
| i                        | <b>Centranthus ruber</b> (L.) Lam. & DC., Fl. Fr. ed.3 4: 239 (1805)                                | 2:278 |
| i                        | <b>Valerianella eriocarpa</b> Desv., J. Bot. 2: 314, t.11 fig.2 (1809)                              | 2:278 |
| i                        | <b>Valerianella locusta</b> (L.) Betteke, Animadv. Bot. Valer. 10 (1826)                            |       |
| it                       | <b>Valerianella rimosa</b> Bast., J. Bot. 3: 20 (1814)  |       |
| <br><b>VERBENACEAE</b>   |   |       |
| i                        | <b>Verbena bonariensis</b> L., Sp. Pl. 1: 20 (1753)   |       |
|                          | Verbena incompta P.W.Michael = <b>Verbena bonariensis</b>   |       |
| i                        | <b>Verbena officinalis</b> L., Sp. Pl. 1: 20 (1753)   | 3:541 |
| <br><b>VIOLACEAE</b>     |   |       |
|                          | <b>Hymenanchera dentata</b> R.Br. ex DC., Prodr. 1: 315 (1824)                                      | 1:55  |
|                          | var. <b>angustifolia</b> (R.Br. ex DC.) Benth. = <b>Hymenanchera dentata</b>                        |       |
| i                        | <b>Viola arvensis</b> Murray, Prodr. Stirp. Gott. 73 (1770)   | 1:55  |
|                          | <b>Viola betonicifolia</b> Smith in Rees, Cycl. 37, no.7 (1817)                                     | 1:54  |
|                          | <b>Viola caleyana</b> G.Don, Gen. Hist. 1: 329 (1831)   | 1:55  |
|                          | <b>Viola cleistogamoides</b> (L.G.Adams) Seppelt in Jessop & Toelken, Fl. S.Australia 2: 870 (1986) |       |

|   |  |      |
|---|--|------|
| t | <b>Viola cunninghamii</b> Hook.f., Fl. N.Z. 1: 16 (1852)                 | 1:54 |
|   | <b>Viola fuscoviolacea</b> (L.G.Adams) T.A.James, Muelleria 9: 35 (1996) |      |
|   | <b>Viola hederacea</b> Labill., Nov. Holl. Pl. 1: 66, t.91 (1805)        | 1:54 |
|   | subsp. <i>cleistogamoides</i> L.Adams = <b>Viola cleistogamoides</b>     |      |
| e | subsp. <b>curtisiae</b> L.G.Adams, Fl. Australia 8: 386 (1982)           |      |
|   | subsp. <i>fuscoviolacea</i> L.G.Adams = <b>Viola fuscoviolacea</b>       |      |
|   | subsp. <b>hederacea</b>  |      |
|   | <b>Viola sieberiana</b> Spreng., Syst. Veg. 4: Cur. Post. 96 (1827)      | 1:54 |

#### WINTERACEAE

|  |   |      |
|--|---|------|
|  | <i>Drimys lanceolata</i> (Poiret) Baillon = <b>Tasmannia lanceolata</b> | 1:25 |
|  | <b>Tasmannia lanceolata</b> (Poiret) A.C.Smith, Taxon 18: 287 (1969)    | 1:25 |

#### ZYGOPHYLLACEAE

|  |   |      |
|--|---|------|
|  | <i>Zygophyllum apiculatum</i> F.Muell., not present in Tasmania, previously recorded in error | 1:93 |
|  | <b>Zygophyllum billardi</b> DC., Prodr. 1: 705 (1824)   | 1:93 |

## Monocotyledonae

#### AGAVACEAE

|   |   |        |
|---|---|--------|
| i | <b>Phormium tenax</b> Forst. & Forst.f., Char. Gen. Pl. 48, t.24 (1775) | 4b:418 |
|---|---|--------|

#### ALISMATACEAE

|   |   |      |
|---|---|------|
| i | <b>Alisma plantago-aquatica</b> L., Sp. Pl. 1: 342 (1753)                             | 4b:4 |
|   | <i>Damasonium australe</i> Salisb. = <b>Damasonium minus</b>                          | 4b:6 |
|   | <b>Damasonium minus</b> (R.Br.) Buchenau, Abh. Naturwiss. Vereine Bremen 2: 20 (1869) | 4b:6 |

#### ALOEACEAE

|   |  |  |
|---|--|--|
| i | <b>Aloe saponaria</b> (Aiton) Haw., Trans. Linn. Soc. Lond. 7: 17 (1804) |  |
|---|--|--|

#### APONOGETONACEAE

|   |  |       |
|---|--|-------|
| i | <b>Aponogeton distachyos</b> L.f., Suppl. Pl. 215 (1782) | 4b:13 |
|---|--|-------|

#### ARACEAE

|   |   |       |
|---|---|-------|
| i | <b>Zantedeschia aethiopica</b> (L.) Sprengel, Sys. Veg. 3: 765 (1826) | 4b:30 |
|---|---|-------|

#### BURMANNIACEAE (THISMIACEAE)

|  |   |        |
|--|---|--------|
|  | <b>Thismia rodwayi</b> F.Muell., Vict. Nat. 7: 115 (1890) | 4b:423 |
|--|---|--------|

#### CENTROLEPIDACEAE

|  |   |       |
|--|---|-------|
|  | <b>Aphelia gracilis</b> Sonder, Linnaea 28: 227 (1856)            | 4b:49 |
|  | <b>Aphelia pumilio</b> F.Muell. ex Sonder, Linnaea 28: 226 (1856) | 4b:50 |



|   |  |       |
|---|--|-------|
|   | <b>Centrolepis aristata</b> (R.Br.) Roemer & Schultes, Syst. Veg. 1: 44 (1817)                         | 4b:54 |
|   | <b>Centrolepis fascicularis</b> Labill., Nov. Holl. Pl. 1: 7, t.1 (1804)                               | 4b:55 |
|   | <b>Centrolepis glabra</b> (F.Muell. ex Sonder) Hieron., Abh. Naturf. Ges. Halle 12: 209 (1873)         | 4b:52 |
| e | <b>Centrolepis monogyna</b> (Hook.f.) Benth., Fl. Austral. 7: 205 (1878)                               | 4b:53 |
| e | subsp. <b>monogyna</b>   | 4b:53 |
| e | subsp. <b>paludicola</b> (W.M.Curtis) D.A.Cooke, J. Adelaide Bot. Gard. 15(1): 22 (1992)               | 4b:54 |
| e | <b>Centrolepis muscoides</b> (Hook.f.) Hieron., Abh. Naturf. Ges. Halle 12: 209 (1873)                 | 4b:53 |
|   | Centrolepis paludicola W.M.Curtis = <b>Centrolepis monogyna</b> subsp. <b>paludicola</b>               | 4b:54 |
| e | <b>Centrolepis pedderensis</b> W.M.Curtis, Brunonia 7: 299 (1984)                                      | 4b:53 |
|   | <b>Centrolepis polygyna</b> (R.Br.) Hieron., Abh. Naturf. Ges. Halle 12: 210 (1873)                    | 4b:51 |
|   | Centrolepis pulvinata (R.Br.) Roemer & Schultes = <b>Centrolepis strigosa</b> subsp. <b>pulvinata</b>  | 4b:55 |
|   | <b>Centrolepis strigosa</b> (R.Br.) Roemer & Schultes, Syst. Veg. 1: 43 (1817)                         | 4b:55 |
| e | subsp. <b>pulvinata</b> (R.Br.) D.A.Cooke, J. Adelaide Bot. Gard. 15(1): 28 (1992)                     | 4b:55 |
|   | subsp. <b>strigosa</b>   | 4b:55 |
| e | <b>Gaimardia amblyphylla</b> W.M.Curtis, Brunonia 7: 299 (1984)  | 4b:57 |
| e | <b>Gaimardia fitzgeraldii</b> F.Muell.&Rodway in Rodway, Pap. Proc. R. Soc. Tasm. for 1894-5:55 (1896) | 4b:57 |
| t | <b>Gaimardia setacea</b> Hook.f., Fl. Nov.-Zel. 1: 267 (1853)  | 4b:57 |

#### COMMELINACEAE

|   |   |       |
|---|---|-------|
| i | <b>Tradescantia albiflora</b> Kunth, Enum. Pl. 4: 84 (1843) | 4b:36 |
|---|---|-------|

#### CYMODOCEACEAE

|  |  |       |
|--|--|-------|
|  | <b>Amphibolis antarctica</b> (Labill.) Sonder & Asch. ex Asch., Linnaea 35: 164 (1867) | 4b:27 |
|  | Cymodocea antarctica (Labill.) Endl. = <b>Amphibolis antarctica</b>                    | 4b:27 |

#### CYPERACEAE

|     |  |        |
|-----|--|--------|
|     | <b>Baumea acuta</b> (Labill.) Palla, Allg. Bot. Zeitschr. 8: 69 (1902)                   | 4b:131 |
|     | <b>Baumea arthrophylla</b> (Nees) Boeck., Linnaea 38: 242 (1874)                         | 4b:129 |
|     | <b>Baumea articulata</b> (R.Br.) S.T.Blake, Contr. Qld. Herb. 8: 28 (1969)               | 4b:128 |
|     | <b>Baumea gunnii</b> (Hook.f.) S.T.Blake, Contr. Qld. Herb. 8: 27 (1969)                 | 4b:130 |
|     | <b>Baumea juncea</b> (R.Br.) Palla, Allg. Bot. Zeitschr. 15: 113 (1909)                  | 4b:129 |
|     | <b>Baumea planifolia</b> (Benth.) K.L.Wilson, Telopea 5: 589 (1994)                      |        |
|     | <b>Baumea rubiginosa</b> (Forst.f.) Boeck., Linnaea 38: 241 (1874)                       | 4b:129 |
|     | <b>Baumea tetragona</b> (Labill.) S.T.Blake, Contr. Qld. Herb. 8: 30 (1969)              | 4b:131 |
|     | <b>Bolboschoenus caldwellii</b> (V.Cook) Soják, Cas. Nár. Mus. Odd. Prir. 141: 62 (1972) | 4b:92  |
|     | <b>Bolboschoenus medianus</b> (V.Cook) Soják, Cas. Nár. Mus. Odd. Prir. 141: 63 (1972)   | 4b:93  |
|     | Carex acicularis sensu Rodway (1903) = <b>Carex archeri</b>                              | 4b:148 |
| it# | <b>Carex albula</b> Allan, Trans. R. Soc. N.Z. 76: 589 (1947)                            | 4b:164 |
|     | <b>Carex appressa</b> R.Br., Prodr. 242 (1810)   | 4b:152 |
| t   | var. <b>virgata</b> (Boott) Kük., Pflanzenr. 38: 179 (1909)                              | 4b:153 |
| e   | forma <b>minor</b> Kük., Pflanzenr. 38: 179 (1909)                                       | 4b:152 |
|     | <b>Carex archeri</b> Boott in Hook.f., Fl. Tasm. 2: 98 (1858)                            | 4b:148 |
| e   | <b>Carex barbata</b> Boott, Ill. Gen. Car. 1: 68, t.187 (1858)                           | 4b:160 |
|     | <b>Carex bichenoviana</b> Boott ex Hook.f., Fl. Tasm. 2: 101 (1858)                      | 4b:167 |

|    |   |        |
|----|---|--------|
|    | <b>Carex breviculmis</b> R.Br., Prodr. 242 (1810)   | 4b:158 |
| i# | <b>Carex buxbaumii</b> Wahlenb., Kongl. Vetensk. Acad. Nya Handl. 24: 163 (1803)  |        |
|    | Carex canescens auctt. non L. = <b>Carex curta</b>  | 4b:155 |
|    | <b>Carex capillacea</b> Boott, Ill. Gen. Carex 1: 44, t.110 (1858)  | 4b:148 |
| e  | <b>Carex cataractae</b> R.Br., Prodr. 242 (1810)  | 4b:162 |
|    | <b>Carex cephalotes</b> F.Muell., Trans. Phil. Soc. Vict. 1: 110 (1855)   | 4b:149 |
|    | <b>Carex chlorantha</b> R.Br., Prodr. 242 (1810)  | 4b:149 |
|    | <b>Carex</b> aff. <b>cirrhosa</b> S.Berggren, Minn. Fys. Sallsk. Lund. 29, t.7, fig.27-34 (1878)                              | 4b:163 |
|    | <b>Carex curta</b> Gooden., Trans. Linn. Soc. Lond. 2: 145 (1794)   | 4b:155 |
| it | <b>Carex demissa</b> Hornem., Dansk. Oekon. Plantel. ed.3, 1: 939 (1821)  | 4b:161 |
|    | <b>Carex</b> aff. <b>diandra</b> Schrank, Cent. Bot. Anmerk. 57: 49 (1781)  | 4b:150 |
| i  | <b>Carex divisa</b> Hudson, Fl. Angl. 1: 348 (1762)   | 4b:149 |
| i  | <b>Carex divulsa</b> Stokes in With., Bot. Arr. Veg. Gt. Brit. ed.2, 2: 1035 (1787)   | 4b:150 |
|    | <b>Carex fascicularis</b> Sol. ex Boott in Hook.f., Fl. Nov.-Zel. 1: 283 (1853)   | 4b:166 |
| i  | <b>Carex flacca</b> Schreber, Spicil. Fl. Lips. 178 (1771)  | 4b:157 |
| it | <b>Carex flagellifera</b> Colenso, Trans. N.Z. Inst. 16: 342 (1884)   | 4b:163 |
|    | Carex flava sensu Rodway (1903) = <b>Carex cataractae</b> , <b>C. demissa</b> , <b>C. flaviformis</b> & <b>C. lepidocarpa</b> |        |
| t  | <b>Carex flaviformis</b> Nelmes, Kew Bull. for 1955: 84 (1955)  | 4b:162 |
|    | <b>Carex gaudichaudiana</b> Kunth, Enum. Pl. 2: 417 (1837)  | 4b:156 |
|    | Carex gaudichaudiana sensu Rodway (1903) = <b>Carex gaudichaudiana</b> , <b>C. polyantha</b> & <b>C. hypandra</b>             |        |
|    | <b>Carex gunniana</b> Boott, Proc. Linn. Soc. Lond. 1: 258 (1845)   | 4b:160 |
|    | <b>Carex hypandra</b> F.Muell. ex Benth., Fl. Austral. 7: 439 (1878)  | 4b:157 |
|    | <b>Carex inversa</b> R.Br., Prodr. 242 (1810)   | 4b:153 |
|    | <b>Carex iynx</b> Nelmes, Proc. Linn. Soc. Lond. 155: 279 (1944)  | 4b:159 |
| it | <b>Carex lepidocarpa</b> Tausch, Flora (Regensb.) 17: 179 (1834)  | 4b:161 |
|    | <b>Carex longebrachiata</b> Boeck., Linnaea 41: 282 (1877)  | 4b:158 |
|    | Carex longifolia sensu Rodway (1903) = <b>Carex iynx</b> & <b>C. longebrachiata</b>   |        |
| i  | <b>Carex ovalis</b> Gooden., Trans. Linn. Soc. Lond. 2: 145 (1794)  | 4b:155 |
|    | Carex paniculata sensu Rodway (1903) = <b>Carex appressa</b>  | 4b:152 |
| i  | <b>Carex pilulifera</b> L., Sp. Pl. 2: 976 (1753)   |        |
|    | <b>Carex polyantha</b> F.Muell., Trans. Phil. Soc. Vict. 1: 110 (1855)  | 4b:156 |
|    | Carex pseudocyperus sensu Rodway (1903) = <b>Carex fascicularis</b>   | 4b:166 |
|    | <b>Carex pumila</b> Thunb. ex Murr., Syst. Veg. 844 (1784)  | 4b:166 |
|    | <b>Carex raleighii</b> Nelmes, Kew Bull. 310 (1939)   | 4b:154 |
| i  | <b>Carex scoparia</b> Schk. ex Willd., Sp. Pl. 4: 230 (1805)  | 4b:154 |
|    | <b>Carex tasmanica</b> Kük., Bull. Herb. Boiss. ser.2, 4: 59 (1904)   | 4b:165 |
|    | <b>Carex tereticaulis</b> F.Muell., Fragm. 8: 256 (1874)  | 4b:151 |
| i  | <b>Carex testacea</b> Sol. ex Boott in Hook.f., Fl. Nov.-Zel. 1: 282 (1853)   | 4b:164 |
|    | Carex virgata Sol. ex Boott = <b>Carex appressa</b> var. <b>virgata</b>   | 4b:153 |
|    | Carex vulgaris sensu Rodway (1903) = <b>Carex gaudichaudiana</b>  | 4b:156 |
|    | <b>Carpha alpina</b> R.Br., Prodr. 230 (1810)   | 4b:87  |
| e  | <b>Carpha curvata</b> W.M.Curtis, Brunonia 7(2): 303 (1984)   | 4b:88  |
| e  | <b>Carpha rodwayi</b> W.M.Curtis, Brunonia 7(2): 302 (1984)   | 4b:88  |
|    | <b>Caustis pentandra</b> R.Br., Prodr. 240 (1810)   | 4b:137 |

|     |   |        |
|-----|---|--------|
|     | <b>Chorizandra australis</b> K.L.Wilson, <i>Telopea</i> 5(4): 594 (1994)                      | 4b:138 |
|     | <i>Chorizandra cymbaria</i> sensu Rodway (1903) = <b>Chorizandra australis</b>                | 4b:138 |
|     | <b>Chorizandra enodis</b> Nees in Lehm., <i>Pl. Preiss.</i> 2: 73 (1846)                      | 4b:138 |
|     | <i>Cladium filum</i> (Labill.) R.Br. = <b>Gahnia filum</b>                                    | 4b:136 |
|     | <i>Cladium glomeratum</i> sensu Rodway (1903) = <b>Baumea rubiginosa</b>                      | 4b:129 |
|     | <i>Cladium gunnii</i> Hook.f. = <b>Baumea gunnii</b>  | 4b:130 |
|     | <i>Cladium junceum</i> R.Br. = <b>Baumea juncea</b>   | 4b:129 |
| i   | <b>Cladium procerum</b> S.T.Blake, <i>Trans. R. Soc. S.Austral.</i> 67: 57 (1943)             | 4b:428 |
|     | <i>Cladium schoenoides</i> R.Br. = <b>Baumea acuta</b>  | 4b:131 |
|     | <i>Cladium tetraquetrum</i> Hook.f. = <b>Baumea tetragona</b>                                 | 4b:131 |
| i   | <b>Cyperus congestus</b> M.Vahl, <i>Enum. Pl.</i> 2: 358 (1805)                               | 4b:91  |
| i   | <b>Cyperus eragrostis</b> Lam., <i>Tabl. Encycl.</i> 1: 146 (1791)                            | 4b:90  |
|     | <b>Cyperus gunnii</b> Hook.f., <i>Fl. Tasm.</i> 2: 80, t.140, fig.A (1858)                    | 4b:91  |
|     | <b>Cyperus lucidus</b> R.Br., <i>Prodr.</i> 218 (1810)  | 4b:91  |
| ?i# | <b>Cyperus sanguinolentus</b> M.Vahl, <i>Enum. Pl.</i> 2: 351 (1805)                          | 4b:90  |
| ?i  | <b>Cyperus tenellus</b> L.f., <i>Suppl.</i> 103 (1782)  | 4b:90  |
|     | <i>Eleocharis acicularis</i> sensu Rodway (1903) = <b>Eleocharis pusilla</b>                  | 4b:108 |
|     | <b>Eleocharis acuta</b> R.Br., <i>Prodr.</i> 224 (1810)                                       | 4b:107 |
|     | <b>Eleocharis gracilis</b> R.Br., <i>Prodr.</i> 224 (1810)                                    | 4b:107 |
|     | <b>Eleocharis pusilla</b> R.Br., <i>Prodr.</i> 225 (1810)                                     | 4b:108 |
|     | <b>Eleocharis sphacelata</b> R.Br., <i>Prodr.</i> 224 (1810)                                  | 4b:107 |
|     | <i>Elynanthus capillaceus</i> Benth. = <b>Tetraria capillaris</b>                             | 4b:132 |
|     | <b>Gahnia filum</b> (Labill.) F.Muell., <i>Key Syst. Vict. Pl.</i> 1: 456 (1888)              | 4b:136 |
|     | <i>Gahnia fitzgeraldi</i> Rodway = <b>Gahnia microstachya</b>                                 | 4b:134 |
|     | <i>Gahnia graminifolia</i> Rodway = <b>Gahnia rodwayi</b>                                     | 4b:134 |
|     | <b>Gahnia grandis</b> (Labill.) S.T.Blake, <i>Contr. Qld. Herb.</i> 8: 33 (1969)              | 4b:135 |
|     | <i>Gahnia melanocarpa</i> R.Br., not present in Tasmania, formerly recorded in error.         |        |
|     | <b>Gahnia microstachya</b> Benth., <i>Fl. Austral.</i> 7: 414 (1878)                          | 4b:134 |
|     | <i>Gahnia psittacorum</i> Labill. = <b>Gahnia grandis</b>                                     | 4b:135 |
|     | <b>Gahnia radula</b> (R.Br.) Benth., <i>Fl. Austral.</i> 7: 417 (1878)                        | 4b:135 |
| e   | <b>Gahnia rodwayi</b> F.Muell. ex Rodway, <i>Pap. Proc. R. Soc. Tasm.</i> for 1892: 93 (1893) | 4b:134 |
|     | <b>Gahnia sieberiana</b> Kunth, <i>Enum. Pl.</i> 2: 332 (1837)                                | 4b:135 |
|     | <b>Gahnia trifida</b> Labill., <i>Nov. Holl. Pl.</i> 1: 89, t.116 (1805)                      | 4b:136 |
|     | <b>Gymnoschoenus sphaerocephalus</b> (R.Br.) Hook.f., <i>Fl. Tasm.</i> 2: 83, t.142 (1858)    | 4b:119 |
|     | <i>Heleocharis</i> , see <b>Eleocharis</b>  | 4b:106 |
|     | <b>Isolepis alpina</b> Hook.f., <i>Fl. Tasm.</i> 2: 68, t.143 (1858)                          | 4b:103 |
|     | <b>Isolepis aucklandica</b> Hook.f., <i>Fl. Antarct.</i> 1: 88 (1844), t.50 (1845)            | 4b:102 |
|     | <b>Isolepis cernua</b> (M.Vahl) Roemer & Schultes, <i>Syst. Veg.</i> 2: 106 (1817)            | 4b:99  |
|     | <b>Isolepis crassiuscula</b> Hook.f., <i>Fl. Tasm.</i> 2: 86, t.143, fig.A (1858)             | 4b:98  |
|     | <b>Isolepis fluitans</b> (L.) R.Br., <i>Prodr.</i> 221 (1810)                                 | 4b:98  |
|     | <b>Isolepis habra</b> (Edgar) Soják, <i>Cas. Nár. Mus. Odd. Prir.</i> 148: 194 (1979)         | 4b:101 |
|     | <b>Isolepis hookeriana</b> Boeck., <i>Flora</i> 41: 418 (1858)                                | 4b:104 |
| i#  | <b>Isolepis hystrix</b> (Thunb.) Nees, <i>Linnaea</i> 7: 496 (1832)                           |        |
|     | <b>Isolepis inundata</b> R.Br., <i>Prodr.</i> 222 (1810)                                      | 4b:100 |

|    |   |        |
|----|---|--------|
| e  | <b>Isolepis limbata</b> W.M.Curtis, Brunonia 7: 300 (1984)  | 4b:103 |
|    | <b>Isolepis marginata</b> (Thunb.) A.Dietr., Sp. Pl. ed.6, 1(2): 110 (1832)                           | 4b:104 |
|    | <b>Isolepis montivaga</b> (S.T.Blake) K.L.Wilson, Telopea 2(2): 168 (1981)                            | 4b:103 |
|    | <b>Isolepis nodosa</b> (Rottb.) R.Br., Prodr. 221 (1810)  | 4b:105 |
|    | <b>Isolepis platycarpa</b> (S.T.Blake) Soják, Cas. Nár. Mus. Odd. Prir. 148: 194 (1979)               | 4b:99  |
|    | <b>Isolepis producta</b> (C.B.Clarke) K.L.Wilson, Telopea 2(2): 168 (1981)                            | 4b:98  |
| i  | <b>Isolepis prolifera</b> (Rottb.) R.Br., Prodr. 223 (1810)   |        |
| it | <b>Isolepis setacea</b> (L.) R.Br., Prodr. 222 (1810)   | 4b:105 |
|    | <b>Isolepis stellata</b> (C.B.Clarke) K.L.Wilson, Telopea 2(2): 169 (1981)                            | 4b:104 |
|    | <b>Isolepis subtilissima</b> Boeck., Flora 41: 416 (1858)   | 4b:101 |
| e  | <b>Isolepis tasmanica</b> (S.T.Blake) K.L.Wilson, Telopea 2(2): 170 (1981)                            | 4b:102 |
|    | <b>Isolepis wakefieldiana</b> (S.T.Blake) K.L.Wilson, Telopea 2(2): 170 (1981)                        | 4b:100 |
|    | <b>Lepidosperma concavum</b> R.Br., Prodr. 234 (1810)   | 4b:126 |
|    | <b>Lepidosperma curtisiae</b> K.L.Wilson & D.I.Morris, Telopea 5(2): 423 (1993)                       | 4b:123 |
|    | <i>Lepidosperma elatior</i> sensu Rodway (1903) = <b>Lepidosperma elatius</b>                         | 4b:124 |
|    | <b>Lepidosperma elatius</b> Labill., Nov. Holl. Pl. 1: 15, t.11 (1805)                                | 4b:124 |
|    | var. <i>ensiforme</i> Rodway = <b>Lepidosperma ensiforme</b>  | 4b:125 |
|    | var. <i>oldfieldii</i> (Hook.f.) Rodway = <b>Lepidosperma oldfieldii</b>                              | 4b:125 |
|    | <b>Lepidosperma ensiforme</b> (Rodway) D.I.Morris, Stud. Fl. Tasm. 4b: 425 (1994)                     | 4b:125 |
|    | <b>Lepidosperma filiforme</b> Labill., Nov. Holl. Pl. 1: 17, t.15 (1805)                              | 4b:121 |
|    | <b>Lepidosperma forsythii</b> A.A.Hamilton, Proc. Linn. Soc. N.S.W. 35: 411 (1910)                    | 4b:122 |
|    | <b>Lepidosperma gladiatum</b> Labill., Nov. Holl. Pl. 1: 15, t.12 (1805)                              | 4b:125 |
| e  | <b>Lepidosperma globosum</b> Labill., Nov. Holl. Pl. 1: 16, t.14 (1805)                               | 4b:123 |
|    | <b>Lepidosperma gunnii</b> Boeck., Linnaea 38: 325 (1874)   | 4b:123 |
| e  | <b>Lepidosperma inops</b> F.Muell. ex Rodway, Pap. Proc. R. Soc. Tasm. for 1892: 93 (1893)            | 4b:122 |
|    | <b>Lepidosperma laterale</b> R.Br., Prodr. 234 (1810)   | 4b:124 |
|    | <i>Lepidosperma lineare</i> sensu Rodway (1903) = <b>Lepidosperma gunnii</b>                          | 4b:123 |
|    | <i>Lepidosperma lineare</i> var. <i>inops</i> (F.Muell. ex Rodway) Rodway = <b>Lepidosperma inops</b> | 4b:122 |
|    | <b>Lepidosperma longitudinale</b> Labill., Nov. Holl. Pl. 1: 16, t.13 (1805)                          | 4b:124 |
|    | <b>Lepidosperma neesii</b> Kunth, Enum. Pl. 2: 319 (1837)   | 4b:127 |
| e  | <b>Lepidosperma oldfieldii</b> Hook.f., Fl. Tasm. 2: 91, t.146A (1858)                                | 4b:125 |
|    | <i>Lepidosperma squamata</i> sensu Rodway (1903) = <b>Lepidosperma concavum</b>                       | 4b:126 |
|    | <b>Lepidosperma tortuosum</b> F.Muell., Fragm. 9: 23 (1875)   | 4b:121 |
|    | <b>Lepidosperma viscidum</b> R.Br., Prodr. 234 (1810)   | 4b:126 |
|    | <i>Mesomelaena sphaerocephala</i> (R.Br.) Benth. = <b>Gymnoschoenus sphaerocephalus</b>               | 4b:119 |
| e  | <b>Oreobolus acutifolius</b> S.T.Blake, Contr. Qld. Herb. 8: 39 (1969)                                | 4b:109 |
|    | <b>Oreobolus distichus</b> F.Muell., Defin. Austral. Pl. 33 (1855)                                    | 4b:109 |
| e  | <b>Oreobolus oligocephalus</b> W.M.Curtis, Brunonia 7(2): 301 (1985)                                  | 4b:110 |
|    | <b>Oreobolus oxycarpus</b> S.T.Blake, Contr. Qld. Herb. 8: 40 (1969)                                  | 4b:110 |
| e  | subsp. <b>brownii</b> Seberg, Bot. J. Linn. Soc. 96: 178 (1988)                                       | 4b:110 |
|    | <b>Oreobolus pumilio</b> R.Br., Prodr. 236 (1810)   | 4b:109 |
|    | subsp. <b>pumilio</b>   |        |
| e  | <b>Oreobolus tholicarpus</b> D.I.Morris, Muelleria 15: 28 (2001)                                      |        |
|    | <i>Schoenoides oligocephalus</i> (W.M.Curtis) Seberg = <b>Oreobolus oligocephalus</b>                 | 4b:110 |

|   |  |        |
|---|--|--------|
|   | <b>Schoenoplectus pungens</b> (M.Vahl) Palla, Bot. Jahrb. 10: 299 (1889)                           | 4b:94  |
|   | <b>Schoenoplectus validus</b> (M.Vahl) A. & D.Löve, Bull. Torrey Bot. Club 81: 33 (1954)           | 4b:95  |
| e | <b>Schoenus absconditus</b> Kük., Feddes Repert. 44: 99 (1938)                                     | 4b:117 |
|   | <b>Schoenus apogon</b> Roemer & Schultes, Syst. Veg. 2: 77 (1817)                                  | 4b:114 |
|   | Schoenus axillaris (R.Br.) Poiret = <b>Schoenus maschalinus</b>                                    | 4b:116 |
| e | <b>Schoenus biglumis</b> Kük., Feddes Repert. 44: 164 (1938)                                       | 4b:118 |
|   | <b>Schoenus brevifolius</b> R.Br., Prodr. 231 (1810)   | 4b:113 |
|   | Schoenus brownii Hook.f. = <b>Schoenus apogon</b>  | 4b:114 |
|   | <b>Schoenus calyptratus</b> Kük., Feddes Repert. 48: 248 (1940)                                    | 4b:117 |
|   | <b>Schoenus carsei</b> Cheeseman, Man. N.Z. Fl. 781 (1906)   | 4b:113 |
|   | <b>Schoenus fluitans</b> Hook.f., Fl. Tasm. 2: 81 (1858)   | 4b:114 |
| # | <b>Schoenus latelaminatus</b> Kük., Feddes Repert. 44: 88 (1938)                                   | 4b:116 |
|   | <b>Schoenus lepidosperma</b> (F.Muell.) K.L.Wilson, Telopea 5(4): 619 (1994)                       | 4b:114 |
|   | subsp. <b>lepidosperma</b>   |        |
|   | <b>Schoenus maschalinus</b> Roemer & Schultes, Syst. Veg. 2: 77 (1817)                             | 4b:116 |
|   | <b>Schoenus nitens</b> (R.Br.) Roemer & Schultes, Syst. Veg. 2: 77 (1817)                          | 4b:118 |
| e | <b>Schoenus pygmaeus</b> S.T.Blake, Contr. Qld. Herb. 8: 41 (1969)                                 | 4b:116 |
|   | Schoenus tenuissimus sensu Rodway (1903) = <b>Schoenus lepidosperma</b> subsp. <b>lepidosperma</b> | 4b:114 |
|   | <b>Schoenus tesquorum</b> J.Black, Trans. Proc. R. Soc. S.Austral. 46: 565 (1922)                  | 4b:115 |
|   | <b>Schoenus turbinatus</b> (R.Br.) Roemer & Schultes, Syst. Veg. 2: 76 (1817)                      | 4b:118 |
|   | Scirpus americanus sensu J.H.Willis (1973) = <b>Schoenoplectus pungens</b>                         | 4b:94  |
|   | Scirpus antarcticus sensu J.H.Willis (1973) = <b>Isolepis marginata</b>                            | 4b:104 |
|   | Scirpus aucklandicus (Hook.f.) Boeck. = <b>Isolepis aucklandica</b>                                | 4b:102 |
|   | Scirpus caldwellii V.Cook = <b>Bolboschoenus caldwellii</b>  | 4b:92  |
|   | Scirpus cartilagineus (R.Br.) Sprengel = <b>Isolepis marginata</b>                                 | 4b:104 |
|   | Scirpus cernuus Vahl = <b>Isolepis cernua</b>  | 4b:99  |
|   | Scirpus crassiusculus (Hook.f.) Benth. = <b>Isolepis crassiuscula</b>                              | 4b:98  |
|   | Scirpus fluitans L. = <b>Isolepis fluitans</b>   | 4b:98  |
|   | Scirpus fluviatilis sensu J.H.Willis (1973) = <b>Bolboschoenus medianus</b>                        | 4b:93  |
|   | Scirpus hookerianus (Boeck.) S.T.Blake = <b>Isolepis hookeriana</b>                                | 4b:104 |
|   | Scirpus inundatus (R.Br.) Poiret = <b>Isolepis inundata</b>  | 4b:100 |
|   | Scirpus lacustris sensu Rodway (1903) = <b>Schoenoplectus validus</b>                              | 4b:95  |
|   | Scirpus lenticularis sensu Rodway (1903) = <b>Isolepis fluitans</b>                                | 4b:98  |
|   | Scirpus maritimus sensu Rodway (1903) = <b>Bolboschoenus caldwellii</b>                            | 4b:92  |
|   | Scirpus montivagus S.T.Blake = <b>Isolepis montivaga</b>   | 4b:103 |
|   | Scirpus nodosus Rottb. = <b>Isolepis nodosa</b>  | 4b:105 |
|   | Scirpus platycarpus S.T.Blake = <b>Isolepis platycarpa</b>   | 4b:99  |
|   | Scirpus productus C.B.Clarke = <b>Isolepis producta</b>  | 4b:98  |
|   | Scirpus pungens Vahl = <b>Schoenoplectus pungens</b>   | 4b:94  |
|   | Scirpus riparius sensu Rodway (1903) = <b>Isolepis cernua</b>                                      | 4b:99  |
|   | Scirpus setaceus L. = <b>Isolepis setacea</b>  | 4b:105 |
|   | Scirpus stellatus C.B.Clarke = <b>Isolepis stellata</b>  | 4b:104 |
|   | Scirpus subtilissimus (Boeck.) S.T.Blake = <b>Isolepis subtilissima</b>                            | 4b:101 |
|   | Scirpus validus Vahl = <b>Schoenoplectus validus</b>   | 4b:95  |

|                             |   |        |
|-----------------------------|---|--------|
|                             | <i>Scirpus wakefieldianus</i> S.T.Blake = <b>Isolepis wakefieldiana</b>                           | 4b:100 |
|                             | <b>Tetraria capillaris</b> (F.Muell.) J.Black, Trans. R. Soc. S.Austral. 58: 169 (1934)           | 4b:132 |
|                             | <b>Tricostularia pauciflora</b> (F.Muell.) Benth., Fl. Austral. 7: 383 (1878)                     | 4b:127 |
|                             | <b>Uncinia compacta</b> R.Br., Prodr. 241 (1810)  | 4b:142 |
| t                           | <b>Uncinia elegans</b> (Kük.) Hamlin, Bull. Dom. Mus. Wellington 19: 11 (1959)                    | 4b:140 |
|                             | <b>Uncinia flaccida</b> S.T.Blake, Proc. R. Soc. Qld. 51: 49 (1940)                               | 4b:143 |
| t                           | <b>Uncinia nervosa</b> Boott in Hook.f., Fl. Tasm. 2: 102 (1858)                                  | 4b:142 |
|                             | <b>Uncinia riparia</b> R.Br., Prodr. 241 (1810)   | 4b:140 |
|                             | <b>Uncinia tenella</b> R.Br., Prodr. 241 (1810)   | 4b:143 |
| <br><b>HAEMODORACEAE</b>    |   |        |
| e                           | <b>Haemodorum distichophyllum</b> Hook., Icon. Pl. 9: t.866 (1851)                                | 4b:364 |
| <br><b>HYDATELLACEAE</b>    |   |        |
| e                           | <b>Hydatella filamentosa</b> (Rodway) W.M.Curtis, Rec. Q. Vic. Mus. 50: 5 (1974)                  | 4b:361 |
|                             | <i>Trithuria filamentosa</i> Rodway = <b>Hydatella filamentosa</b>                                | 4b:361 |
|                             | <b>Trithuria submersa</b> Hook.f., Fl. Tasm. 2: 79 (1858)   | 4b:360 |
| <br><b>HYDROCHARITACEAE</b> |   |        |
| i                           | <b>Egeria densa</b> Planchon, Ann. Sci. Nat. ser.3, 11: 80 (1849)                                 | 4b:8   |
| i                           | <b>Elodea canadensis</b> Michaux, Fl. Bor. Amer. 1: 20 (1803)                                     | 4b:8   |
|                             | <i>Elodea densa</i> (Planchon) Casp. = <b>Egeria densa</b>  | 4b:8   |
|                             | <b>Halophila australis</b> Doty & Stone, Brittonia 18: 306, fig.2 (1966)                          | 4b:13  |
|                             | <i>Halophila ovalis</i> sensu Rodway (1903) = <b>Halophila australis</b>                          | 4b:13  |
| i                           | <b>Lagarosiphon major</b> (Ridley) Moss, Trans. Roy. Soc. S.Afr. 16: 193 (1928)                   | 4b:9   |
|                             | <b>Vallisneria americana</b> Mich., Fl. Bor.-Amer. 2: 220 (1803)                                  | 4b:11  |
|                             | var. <b>americana</b>   |        |
|                             | <i>Vallisneria gigantea</i> Graebner = <b>Vallisneria americana</b>                               | 4b:11  |
|                             | <i>Vallisneria spiralis</i> sensu Rodway (1903) = <b>Vallisneria americana</b>                    | 4b:11  |
| <br><b>IRIDACEAE</b>        |   |        |
| i                           | <b>Chasmanthe floribunda</b> (Salisb.) N.E.Br., Trans. R. Soc. S.Af. 20: 274 (1932)               |        |
| i                           | <b>Crocosmia Xcrocosmiiflora</b> (Lemoine ex Morren) N.E.Br., Trans. R. Soc. S.Af. 20: 264 (1932) | 4b:417 |
| e                           | <b>Diplarrena latifolia</b> Benth., Fl. Austral. 6: 400 (1873)                                    | 4b:405 |
|                             | <b>Diplarrena moraea</b> Labill., Voy. Rech. Pérouse 1: 157 (1800)                                | 4b:405 |
| i                           | <b>Freesia</b> hybrid   | 4b:416 |
| i                           | <b>Gladiolus communis</b> L., Sp. Pl. 1: 36 (1753)  | 4b:413 |
| i                           | subsp. <b>byzantinus</b> (Miller) A.P.Ham., Bot. J. Linn. Soc. 76: 358 (1978)                     | 4b:413 |
|                             | <i>Gladiolus cuspidatus</i> Jacq. = <b>Gladiolus undulatus</b>                                    | 4b:413 |
| i                           | <b>Gladiolus tristis</b> L., Sp. Pl. ed.2, 1: 53 (1762)   | 4b:411 |
| i                           | <b>Gladiolus undulatus</b> L., Mant. Pl. 1: 27 (1767)   | 4b:413 |
|                             | <i>Hewardia tasmanica</i> Hook. = <b>Isophysis tasmanica</b>                                      | 4b:401 |
|                             | <i>Homeria breyniana</i> sensu J.H.Willis (1973) = <b>Homeria flaccida</b>                        | 4b:408 |
| i                           | <b>Homeria flaccida</b> Sweet, Hort. Brit. 395 (1826)   | 4b:408 |

|                      |  |        |
|----------------------|--|--------|
| it                   | <b>Iris foetidissima</b> L., Sp. Pl. 1: 39 (1753)  | 4b:406 |
| i                    | <b>Iris germanica</b> L., Sp. Pl. 1: 38 (1753)   | 4b:406 |
| e                    | <b>Isophysis tasmanica</b> (Hook.) T.Moore, Proc. Linn. Soc. London 2: 212 (1853)                | 4b:401 |
| i                    | <b>Ixia flexuosa</b> L., Sp. Pl. ed.2, 1: 51 (1762)  | 4b:414 |
| i                    | <b>Ixia maculata</b> L., Sp. Pl. ed.2, 2: 1664 (1763)  | 4b:414 |
| i                    | <b>Ixia polystachya</b> L., Sp. Pl. ed.2, 1: 51 (1762)   | 4b:414 |
|                      | <b>Libertia pulchella</b> Sprengel, Syst. Veg. 1: 69 (1824)                                      | 4b:401 |
|                      | var. <b>pulchella</b>  | 4b:402 |
| e                    | var. <b>pygmaea</b> D.I.Morris, Stud. Fl. Tasm. 4b: 425 (1994)                                   | 4b:402 |
|                      | <b>Patersonia fragilis</b> (Labill.) Aschers & Graebner, Syn. Mitteleur. Fl. 3: 532 (1906)       | 4b:403 |
|                      | <i>Patersonia glauca</i> R.Br. = <b>Patersonia fragilis</b>                                      | 4b:403 |
|                      | <i>Patersonia longiscapa</i> Sweet = <b>Patersonia occidentalis</b>                              | 4b:403 |
|                      | <b>Patersonia occidentalis</b> R.Br., Prodr. 304 (1810)  | 4b:403 |
|                      | <i>Romulea longifolia</i> (Salisb.) Baker = <b>Romulea rosea</b> var. <b>australis</b>           | 4b:409 |
| i                    | <b>Romulea minutiflora</b> Klatt, Abh. Naturf. Ges. Halle 15: 399 (1882)                         |        |
| i                    | <b>Romulea rosea</b> (L.) Ecklon, Topogr. Verz. Pflanzensamml. Ecklon 19 (1827)                  | 4b:409 |
| i                    | var. <b>australis</b> (Ewart) De Vos, J. S.Af. Bot. Supp. 9: 254 (1972)                          | 4b:409 |
| i                    | <b>Schizostylis coccinea</b> Backh. & Harvey, Curtis's Bot. Mag. t.5422 (1864)                   | 4b:417 |
| i                    | <b>Sisyrinchium iridifolium</b> Kunth in Humboldt, Bonpland & Kunth, Nov. Gen. Sp. 1: 324 (1816) | 4b:402 |
| i                    | <b>Sparaxis bulbifera</b> (L.) Ker Gawler, Ann. Bot. (König & Sims) 1: 225 (1804)                | 4b:415 |
| i                    | <b>Tritonia lineata</b> (Salisb.) Ker Gawler, Ann. Bot. (König & Sims) 1: 229 (1804)             | 4b:416 |
|                      | <i>Watsonia bulbifera</i> J.Mathews & L.Bolus = <b>Watsonia meriana</b> var. <b>bulbillifera</b> | 4b:408 |
| i                    | <b>Watsonia meriana</b> (L.) Miller, Gard. Dict. ed.8 no.1 (1768)                                | 4b:409 |
| i                    | var. <b>bulbillifera</b> (J.Mathews & L.Bolus) D.A.Cooke, J. Adelaide Bot. Gard. 18(1): 6 (1998) | 4b:408 |
| i                    | var. <b>meriana</b>  | 4b:409 |
| i                    | <b>Watsonia versfeldii</b> J.Mathews & L.Bolus, Ann. Bolus Herb. 3: 140 (1922)                   | 4b:409 |
| i                    | var. <b>alba</b> J.Mathews & L.Bolus, Ann. Bolus Herb. 3: 140 (1922)                             | 4b:409 |
| <br><b>JUNCACEAE</b> |  |        |
| i                    | <b>Juncus acuminatus</b> Michx., Fl. Bor. Am. 1: 192 (1803)                                      | 4b:75  |
| i                    | <b>Juncus acutiflorus</b> Ehrh. ex Hoffm., Deutschl. Fl. 125 (1791)                              | 4b:77  |
| i                    | <b>Juncus acutus</b> L., Sp. Pl. 1: 325 (1753)   |        |
|                      | <b>Juncus amabilis</b> Edgar, N.Z. J. Bot. 2: 186, figs.7, 21 (1964)                             | 4b:67  |
|                      | <b>Juncus antarcticus</b> Hook.f., Fl. Antarct. 1: 79, t.46 (1844)                               | 4b:71  |
| i                    | <b>Juncus articulatus</b> L., Sp. Pl. 1: 327 (1753)  | 4b:77  |
| e                    | <b>Juncus astreptus</b> L.A.S.Johnson in Banks et al., Aspects Tasm. Bot. 41 (1991)              | 4b:66  |
|                      | <b>Juncus australis</b> Hook.f., Fl. Tasm. 2: 66, t.134a (1858)                                  | 4b:67  |
|                      | <b>Juncus bassianus</b> L.A.S.Johnson in Banks et al., Aspects Tasm. Bot. 40 (1991)              | 4b:66  |
|                      | <b>Juncus bufonius</b> L., Sp. Pl. 1: 328 (1753)   | 4b:63  |
| i                    | <b>Juncus bulbosus</b> L., Sp. Pl. 1: 327 (1753)   | 4b:73  |
|                      | <b>Juncus caespiticus</b> E.Meyer in Lehm., Pl. Preiss. 2: 47 (1846)                             | 4b:72  |
|                      | <i>Juncus capillaceus</i> sensu Rodway (1903) = <b>Juncus sandwithii</b>                         | 4b:74  |
| i                    | <b>Juncus capitatus</b> Weigel, Observ. Bot. 28 (1772)   | 4b:70  |
|                      | <i>Juncus communis</i> sensu Rodway (1903) = <b>Juncus australis</b>                             | 4b:67  |

|    |  |       |
|----|--|-------|
| i  | <b>Juncus conglomeratus</b> L., Sp. Pl. 1: 326 (1753)  | 4b:69 |
| e  | <b>Juncus curtisiae</b> L.A.S.Johnson in Banks et al., Aspects Tasm. Bot. 44 (1991)              | 4b:74 |
| i  | <b>Juncus effusus</b> L., Sp. Pl. 1: 326 (1753)  | 4b:69 |
|    | <b>Juncus falcatus</b> E.Meyer, Synops. Luzul. 34 (1823)   | 4b:71 |
|    | <b>Juncus filicaulis</b> Buchenau, Proc. Linn. Soc. N.S.W. 28: 913 (1904)                        | 4b:68 |
|    | <b>Juncus fockei</b> Buchenau in Engl., Bot. Jahrb. 12: 358 (1890)                               | 4b:75 |
| i  | <b>Juncus gerardii</b> Loisel, J. Bot. (Desvaux) 2: 284 (1809)                                   |       |
|    | <b>Juncus gregiflorus</b> L.A.S.Johnson, Contr. N.S.W. Natl. Herb. 3: 243 (1963)                 | 4b:66 |
|    | <b>Juncus holoschoenus</b> R.Br., Prodr. 259 (1810)  | 4b:75 |
| i  | <b>Juncus indescryptus</b> Steudel, Syn. Pl. Glum. 2: 304 (1855)                                 | 4b:72 |
|    | <b>Juncus kraussii</b> Hochst., Flora 28: 342 (1845)   | 4b:70 |
|    | subsp. <b>australiensis</b> (Buchenau) Snogerup, Willdenowia 23: 61 (1993)                       |       |
|    | <i>Juncus maritimus</i> sensu Rodway (1903) = <b>Juncus kraussii</b> subsp. <b>australiensis</b> | 4b:70 |
| i  | <b>Juncus microcephalus</b> Kunth in Humboldt, Bonpland & Kunth, Nov. Gen. Sp. 1: 237 (1816)     | 4b:75 |
|    | <b>Juncus pallidus</b> R.Br., Prodr. 258 (1810)  | 4b:64 |
|    | <b>Juncus pauciflorus</b> R.Br., Prodr. 259 (1810)   | 4b:65 |
|    | <b>Juncus planifolius</b> R.Br., Prodr. 259 (1810)   | 4b:71 |
|    | <b>Juncus prismatocarpus</b> R.Br., Prodr. 259 (1810)  | 4b:73 |
|    | <b>Juncus procerus</b> E.Meyer, Linnaea 3: 367 (1827)  | 4b:65 |
|    | <i>Juncus pusillus</i> sensu J.H.Willis (1973) = <b>Juncus sandwithii</b>                        | 4b:74 |
| e  | <b>Juncus ratkowskyanus</b> L.A.S.Johnson in Banks et al., Aspects Tasm. Bot. 43 (1991)          | 4b:74 |
|    | <b>Juncus revolutus</b> R.Br., Prodr. 259 (1810)   | 4b:63 |
|    | <b>Juncus sandwithii</b> Lourteig, Com. Nat. Fr. Rec. Antarct. 23: 44 (1968)                     | 4b:74 |
|    | <b>Juncus sarophorus</b> L.A.S.Johnson, Contr. N.S.W. Natl. Herb. 3: 242 (1963)                  | 4b:68 |
| it | <b>Juncus squarrosus</b> L., Sp. Pl. 1: 327 (1753)   | 4b:62 |
|    | <b>Juncus subsecundus</b> N.A.Wakef., Vict. Nat. 73: 211 (1957)                                  | 4b:69 |
| i  | <b>Juncus tenuis</b> Willd., Sp. Pl. ed.4, 2: 214 (1799)   | 4b:62 |
|    | <b>Juncus vaginatus</b> R.Br., Prodr. 258 (1810)   | 4b:65 |
|    | <b>Luzula acutifolia</b> Nordensk., Bot. Not. 122: 85 (1969)                                     | 4b:84 |
|    | <b>Luzula atrata</b> Edgar, N.Z. J. Bot. 13: 794 (1975)  | 4b:83 |
|    | <b>Luzula australasica</b> Steudel, Syn. Pl. Glum. 2: 294 (1855)                                 | 4b:82 |
| e  | subsp. <b>australasica</b>   | 4b:82 |
| it | <b>Luzula campestris</b> (L.) DC. in Lam. & DC., Fl. Fr. ed.3, 3: 161 (1805)                     | 4b:81 |
| it | <b>Luzula congesta</b> (Thuill.) Lej., Fl. Spa 1: 168 (1811)                                     | 4b:81 |
|    | <b>Luzula densiflora</b> (Nordensk.) Edgar, N.Z. J. Bot. 13: 786 (1975)                          | 4b:80 |
|    | <b>Luzula flaccida</b> (Buchenau) Edgar, N.Z. J. Bot. 13: 786 (1975)                             | 4b:80 |
|    | <b>Luzula meridionalis</b> Nordensk., Bot. Not. 122: 71 (1969)                                   | 4b:78 |
|    | <b>Luzula modesta</b> Buchenau in Engler, Pflanzenz. 25: 80, fig.51 (1906)                       | 4b:82 |
| it | <b>Luzula multiflora</b> (Ehrh.) Lej., Fl. Spa 1: 169 (1811)                                     | 4b:81 |
|    | <b>Luzula novae-cambriae</b> Gandoger, Bull. Soc. Bot. Fr. 46: 392 (1900)                        | 4b:82 |
|    | <i>Luzula oldfieldii</i> Hook.f. = <b>Luzula australasica</b> subsp. <b>australasica</b>         | 4b:82 |
| e  | <b>Luzula poimena</b> W.M.Curtis, Brunonia 7: 297 (1985)   | 4b:83 |

## JUNCAGINACEAE



|                      |   |        |
|----------------------|---|--------|
|                      | <b>Triglochin alcockiae</b> H.Aston, Muellera 8(1): 85 (1993)   | 4b:427 |
|                      | Triglochin centropurpureum (Tasmanian plants) = <b>Triglochin nanum</b>                                     | 4b:16  |
|                      | <b>Triglochin minutissimum</b> F.Muell., Fragm. 6: 82 (1867)  | 4b:16  |
|                      | <b>Triglochin mucronatum</b> R.Br., Prodr. 343 (1810)   | 4b:16  |
|                      | <b>Triglochin nanum</b> F.Muell., Fragm. 6: 82 (1867)   |        |
|                      | <b>Triglochin procerum</b> R.Br., Prodr. 343 (1810)   | 4b:14  |
|                      | <b>Triglochin rheophilum</b> H.Aston, Muellera 8(1): 94 (1993)  | 4b:427 |
|                      | <b>Triglochin striatum</b> Ruíz Lopez & Pavón, Fl. Peruv. 3: 72 (1802)                                      | 4b:16  |
| <br><b>LEMNACEAE</b> |   |        |
|                      | <b>Lemna disperma</b> Hegelm., Bot. Zeitung 29: 655 (1871)  | 4b:32  |
|                      | Lemna minor sensu Rodway (1903) = <b>Lemna disperma</b>   | 4b:32  |
|                      | <b>Lemna trisulca</b> L., Sp. Pl. 2: 970 (1753)   | 4b:32  |
|                      | Spirodela polyrhiza (L.) Schleiden, not present in Tasmania, formerly recorded in error                     |        |
|                      | Wolffia arrhiza sensu J.H. Willis (1973) = <b>Wolffia australiana</b>                                       | 4b:32  |
|                      | <b>Wolffia australiana</b> (Benth.) Hartog & Plas, Blumea 20: 151 (1972)                                    | 4b:32  |
| <br><b>LILIACEAE</b> |   |        |
| i                    | <b>Allium ampeloprasum</b> L., Sp. Pl. 1: 294 (1753)  |        |
| i                    | <b>Allium neapolitanum</b> Cirillo, Pl. Rar. Neapol. 1: 13, t.4 (1788)                                      | 4b:392 |
| i                    | <b>Allium triquetrum</b> L., Sp. Pl. 1: 300 (1753)  | 4b:392 |
| i                    | <b>Allium vineale</b> L., Sp. Pl. 1: 299 (1753)   | 4b:392 |
|                      | Anguillaria dioica R.Br. pro spec. Tasm. = <b>Wurmbea uniflora</b> , <b>W. dioica</b> & <b>W. latifolia</b> |        |
|                      | <b>Arthropodium milleflorum</b> (DC.) J.F. Macbride, Contr. Gray Herb. n.ser. 3(56): 2 (1918)               | 4b:384 |
|                      | <b>Arthropodium minus</b> R.Br., Prodr. 276 (1810)  | 4b:385 |
|                      | Arthropodium paniculatum (Haw.) R.Br. = <b>Arthropodium milleflorum</b>                                     | 4b:384 |
|                      | <b>Arthropodium pendulum</b> DC., Cat. Pl. Horti Monsp. 80 (1813)   |        |
|                      | <b>Arthropodium strictum</b> R.Br., Prodr. 276 (1810)   | 4b:388 |
| i                    | <b>Asparagus asparagoides</b> (L.) W. Wight, Cent. Dict. Cycl. 12: 845 (1909)                               |        |
| i                    | <b>Asparagus officinalis</b> L., Sp. Pl. 1: 313 (1753)  | 4b:370 |
| i                    | <b>Asparagus scandens</b> Thunb., Prod. Pl. Cap. 66 (1794)  |        |
| i                    | <b>Asphodelus fistulosus</b> L., Sp. Pl. 1: 309 (1753)  | 4b:380 |
|                      | <b>Astelia alpina</b> R.Br., Prodr. 291 (1810)  | 4b:370 |
| e                    | var. <b>alpina</b>  | 4b:370 |
|                      | Blandfordia marginata Herbert = <b>Blandfordia punicea</b>  | 4b:374 |
| e                    | <b>Blandfordia punicea</b> (Labill.) Sweet, Hort. Brit. ed.2 517 (1830)                                     | 4b:374 |
|                      | <b>Bulbine bulbosa</b> (R.Br.) Haw., Rev. Pl. Succ. 33 (1821)   | 4b:383 |
|                      | <b>Bulbine glauca</b> (Raf.) E.M. Watson, Fl. Australia 45: 469 (1987)                                      | 4b:382 |
|                      | <b>Bulbine semibarbata</b> (R.Br.) Haw., Rev. Pl. Succ. 33 (1821)   | 4b:382 |
|                      | <b>Burchardia umbellata</b> R.Br., Prodr. 273 (1810)  | 4b:396 |
|                      | <b>Caesia alpina</b> Hook.f., Fl. Tasm. 2: 373 (1859)   | 4b:386 |
|                      | <b>Caesia calliantha</b> R. Henderson, Fl. Australia 45: 471 (1987)   | 4b:387 |
|                      | <b>Caesia parviflora</b> R.Br., Prodr. 277 (1810)   | 4b:386 |
|                      | var. <b>minor</b> R. Henderson, Fl. Australia 45: 471 (1987)  | 4b:386 |

|     |   |        |
|-----|---|--------|
|     | var. <b>parviflora</b>  | 4b:386 |
|     | var. <b>vittata</b> (R.Br.) R.Henderson, Fl. Australia 45: 472 (1987)   | 4b:386 |
|     | <i>Caesia vittata</i> sensu Rodway (1903) = <b>Caesia calliantha</b>  | 4b:387 |
| e   | <b>Campynema lineare</b> Labill., Nov. Holl. Pl. 1: 93, t.121 (1805)  | 4b:395 |
|     | <b>Chamaescilla corymbosa</b> (R.Br.) F.Muell. ex Benth., Fl. Austral. 7: 48 (1878)                               | 4b:387 |
|     | var. <b>corymbosa</b>   | 4b:387 |
|     | <i>Chlorophytum alpinum</i> (Hook.f.) Benth. ex Baker = <b>Caesia alpina</b>                                      | 4b:386 |
|     | <b>Dianella amoena</b> G.W.Carr & P.F.Horsfall, Muelleria 8(3): 369 (1995)  |        |
|     | <b>Dianella brevicaulis</b> (Ostenf.) G.W.Carr & P.F.Horsfall, Muelleria 8(3): 375 (1995)                         | 4b:378 |
|     | <i>Dianella caerulea</i> sensu Henderson, Fl. Australia 45: 211 (1987)(Tasmanian plants) = <b>Dianella amoena</b> |        |
|     | <i>Dianella laevis</i> R.Br. = <b>Dianella amoena</b>   | 4b:377 |
|     | <i>Dianella longifolia</i> sensu Curtis & Morris (1994) = <b>Dianella amoena</b>                                  | 4b:377 |
|     | <b>Dianella revoluta</b> R.Br., Prodr. 280 (1810)   | 4b:378 |
|     | var. <b>brevicaulis</b> Ostenf. = <b>Dianella brevicaulis</b>   | 4b:378 |
|     | <b>Dianella tasmanica</b> Hook.f., Fl. Tasm. 2: 57, t.133A (1858)   | 4b:378 |
|     | <i>Dichopogon strictus</i> (R.Br.) Baker = <b>Arthropodium strictum</b>   | 4b:388 |
|     | <b>Drymophila cyanocarpa</b> R.Br., Prodr. 292 (1810)   | 4b:369 |
|     | <b>Herpolorion novae-zelandiae</b> Hook.f., Fl. Nov.-Zel. 1: 258 (1853)   | 4b:383 |
|     | <b>Hypoxis glabella</b> R.Br., Prodr. 289 (1810)  | 4b:375 |
|     | var. <b>glabella</b>  | 4b:375 |
|     | <b>Hypoxis hygrometrica</b> Labill., Nov. Holl. Pl. 1: 82 t.108 (1805)  | 4b:376 |
|     | var. <b>hygrometrica</b>  | 4b:376 |
|     | var. <b>villosispala</b> R.Henderson, Fl. Australia 45: 488 (1987)  | 4b:377 |
|     | <i>Hypoxis pusilla</i> Hook.f. = <b>Hypoxis glabella</b> var. <b>glabella</b>                                     | 4b:375 |
|     | <b>Hypoxis vaginata</b> Schldtl., Linnaea 20: 568 (1847)  | 4b:376 |
|     | var. <b>brevistigmata</b> R.Henderson, Fl. Australia 45: 490 (1987)   | 4b:376 |
|     | var. <b>vaginata</b>  | 4b:376 |
|     | <i>Laxmannia morrisii</i> Keighery. Not a Tasmanian plant   |        |
|     | <b>Laxmannia orientalis</b> Keighery, Fl. Australia 45: 492 (1987)  | 4b:388 |
|     | <i>Laxmannia sessiliflora</i> sensu Rodway (1903) = <b>Laxmannia orientalis</b>                                   | 4b:388 |
| e   | <b>Milligania densiflora</b> Hook.f., J. Bot. Kew Gard. Misc. 5: 298 (1853)                                       | 4b:374 |
| e   | <b>Milligania johnstonii</b> F.Muell. ex Benth., Fl. Austral. 7: 26 (1878)  | 4b:371 |
| e   | <b>Milligania lindoniana</b> Rodway ex W.M.Curtis, Rec. Q. Vic. Mus. 45: 1 (1972)                                 | 4b:372 |
| e   | <b>Milligania longifolia</b> Hook.f., J. Bot. Kew Gard. Misc. 5: 297 (1853)                                       | 4b:373 |
| e   | <b>Milligania stylosa</b> (F.Muell. ex Hook.f.) F.Muell. ex Benth., Fl. Austral. 7: 27 (1878)                     | 4b:373 |
|     | <i>Myrsiphyllum asparagoides</i> (L.) Willd. = <b>Asparagus asparagoides</b>                                      |        |
|     | <i>Myrsiphyllum scandens</i> (Thunb.) Oberm. = <b>Asparagus scandens</b>  |        |
| i   | <b>Narcissus pseudonarcissus</b> L., Sp. Pl. 1: 289 (1753)  | 4b:394 |
| i   | <b>Nothoscordum borbonicum</b> Kunth, Enum. Pl. 4: 463 (1843)   | 4b:394 |
|     | <i>Nothoscordum gracile</i> sensu Curtis & Morris (1994) = <b>Nothoscordum borbonicum</b>                         | 4b:394 |
| it# | <b>Ornithogalum angustifolium</b> Boreau., Fl. Centre Fr. ed.3, 2: 625 (1857)                                     |        |
| i   | <b>Ornithogalum umbellatum</b> L., Sp. Pl. 1: 307 (1753)  | 4b:369 |
| i#  | <b>Scilla peruviana</b> L., Sp. Pl. 1: 309 (1753)   | 4b:391 |
|     | <b>Sowerbaea juncea</b> Andrews, Bot. Repos. 2: t.81 (1800)   | 4b:389 |

|                    |   |        |
|--------------------|---|--------|
|                    | <i>Stypandra caespitosa</i> R.Br. = <b><i>Thelionema caespitosum</i></b>  | 4b:379 |
|                    | <b><i>Thelionema caespitosum</i></b> (R.Br.) R.Henderson, <i>Austrobaileya</i> 2: 109 (1985)                                | 4b:379 |
|                    | <b><i>Thelionema umbellatum</i></b> (R.Br.) R.Henderson, <i>Austrobaileya</i> 2: 109 (1985)                                 | 4b:380 |
|                    | <b><i>Thysanotus patersonii</i></b> R.Br., <i>Prodr.</i> 284 (1810)   | 4b:389 |
|                    | <b><i>Tricoryne elatior</i></b> R.Br., <i>Prodr.</i> 278 (1810)   | 4b:391 |
|                    | <b><i>Wurmbea dioica</i></b> (R.Br.) F.Muell., <i>Fragm.</i> 10: 119 (1877)   | 4b:396 |
|                    | subsp. <b><i>dioica</i></b>   |        |
| ?x                 | <b><i>Wurmbea latifolia</i></b> T.Macfarlane, <i>Brunonia</i> 3: 170 (1980)   | 4b:398 |
|                    | <b><i>Wurmbea uniflora</i></b> (R.Br.) T.Macfarlane, <i>Brunonia</i> 3: 194 (1980)  | 4b:398 |
| <b>ORCHIDACEAE</b> |   |        |
|                    | <i>Acianthus caudatus</i> R.Br. = <b><i>Nemacianthus caudatus</i></b>   | 4a:117 |
|                    | <i>Acianthus exsertus</i> sensu W.M.Curtis (1979) = <b><i>Acianthus pusillus</i></b>  | 4a:117 |
|                    | <b><i>Acianthus pusillus</i></b> D.L.Jones, <i>Austral. Orchid Res.</i> 2: 7 (1991)   | 4a:117 |
|                    | <i>Acianthus reniformis</i> (R.Br.) Schltr. = <b><i>Cyrtostylis reniformis</i></b>  | 4a:119 |
|                    | <i>Acianthus viridis</i> Hook.f. = <b><i>Townsonia viridis</i></b>  | 4a:118 |
|                    | <b><i>Anzybas fordhamii</i></b> (Rupp) D.L.Jones & M.A.Clem., <i>Orchadian</i> 13(10): 443 (2002)                           | 4a:124 |
|                    | <b><i>Anzybas unguiculatus</i></b> (R.Br.) D.L.Jones & M.A.Clem., <i>Orchadian</i> 13(10): 443 (2002)                       | 4a:124 |
|                    | <i>Arachnorchis anthracina</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b><i>Caladenia anthracina</i></b>                       |        |
|                    | <i>Arachnorchis australis</i> (G.W.Carr) D.L.Jones & M.A.Clem. = <b><i>Caladenia australis</i></b>                          |        |
|                    | <i>Arachnorchis brachyscapa</i> (G.W.Carr) D.L.Jones & M.A.Clem. = <b><i>Caladenia brachyscapa</i></b>                      |        |
|                    | <i>Arachnorchis cardiochila</i> (Tate) D.L.Jones & M.A.Clem. = <b><i>Caladenia cardiochila</i></b>                          |        |
|                    | <i>Arachnorchis caudata</i> (Nicholls) D.L.Jones & M.A.Clem. = <b><i>Caladenia caudata</i></b>                              | 4a:101 |
|                    | <i>Arachnorchis clavigera</i> (A.Cunn. ex Lindley) D.L.Jones & M.A.Clem. = <b><i>Caladenia clavigera</i></b>                | 4a:105 |
|                    | <i>Arachnorchis dienema</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b><i>Caladenia dienema</i></b>                             |        |
|                    | <i>Arachnorchis dilatata</i> (R.Br.) D.L.Jones & M.A.Clem. = <b><i>Caladenia dilatata</i></b>                               | 4a:100 |
|                    | <i>Arachnorchis echidnachila</i> (Nicholls) D.L.Jones & M.A.Clem. = <b><i>Caladenia echidnachila</i></b>                    | 4a:101 |
|                    | <i>Arachnorchis helvina</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b><i>Caladenia helvina</i></b>                             |        |
|                    | <i>Arachnorchis lindleyana</i> (Rchb.f.) D.L.Jones & M.A.Clem. = <b><i>Caladenia lindleyana</i></b>                         |        |
|                    | <i>Arachnorchis pallida</i> (Lindley) D.L.Jones & M.A.Clem. = <b><i>Caladenia pallida</i></b>                               | 4a:104 |
|                    | <i>Arachnorchis patersonii</i> (R.Br.) D.L.Jones & M.A.Clem. = <b><i>Caladenia patersonii</i></b>                           | 4a:100 |
|                    | <i>Arachnorchis saggicola</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b><i>Caladenia saggicola</i></b>                         |        |
|                    | <i>Arthrochilus huntianus</i> (F.Muell.) Blaxell  |        |
|                    | subsp. <i>huntianus</i> = <b><i>Thynninorchis huntiana</i></b>  | 4a:89  |
|                    | subsp. <i>nothofagicola</i> D.L.Jones = <b><i>Thynninorchis nothofagicola</i></b>   |        |
|                    | <b><i>Bunochilus melagrammus</i></b> (D.L.Jones) D.L.Jones & M.A.Clem., <i>Austral. Orchid Res.</i> 4: 66 (2002)            | 4a:27  |
| e                  | <b><i>Bunochilus stenochilus</i></b> (D.L.Jones) D.L.Jones & M.A.Clem., <i>Austral. Orchid Res.</i> 4: 67 (2002)            |        |
|                    | <b><i>Bunochilus tunstallii</i></b> (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem., <i>Austral. Orchid Res.</i> 4: 66 (2002) |        |
| e                  | <b><i>Bunochilus williamsonii</i></b> (D.L.Jones) D.L.Jones & M.A.Clem., <i>Austral. Orchid Res.</i> 4: 66 (2002)           |        |
|                    | <b><i>Burnettia cuneata</i></b> Lindley, <i>Gen. Sp. Orchid. Pl.</i> 518 (1840)   | 4a:92  |
|                    | <b><i>Caladenia alata</i></b> R.Br., <i>Prodr.</i> 324 (1810)   |        |
|                    | <i>Caladenia alba</i> R.Br., not present in Tasmania, formerly recorded in error  |        |
|                    | <b><i>Caladenia alpina</i></b> R.S.Rogers, <i>Trans. Roy. Soc. S.Austral.</i> 51: 12 (1927)                                 | 4a:111 |
| e                  | <b><i>Caladenia angustata</i></b> Lindley, <i>Gen. Sp. Orchid. Pl.</i> 420 (1840)   | 4a:108 |

- e **Caladenia anthracina** D.L.Jones, Austral. Orchid Res. 3: 21 (1998)  
*Caladenia atkinsonii* Rodway = **Caladenia carnea** 4a:107
- e **Caladenia atrata** D.L.Jones, Muelleria 8(2): 178 (1994)
- e **Caladenia atrochila** D.L.Jones, Austral. Orchid Res. 3: 23 (1998)  
**Caladenia aurantiaca** (R.S.Rogers) Rupp, Proc. Linn. Soc. N.S.W. 71: 280 (1947)  
**Caladenia australis** G.W.Carr, Indig. Flora & Fauna Assn. Misc. Pap. 1: 2 (1991)  
**Caladenia brachyscapa** G.W.Carr, Muelleria 6(6): 439 (1988)
- e **Caladenia campbellii** D.L.Jones, Austral. Orchid Res. 3: 25 (1998)  
**Caladenia cardiochila** Tate, Trans. Roy. Soc. S.Austral. 9: 60, t.2 (1887)  
**Caladenia carnea** R.Br., Prodr. 324 (1810) 4a:106  
*Caladenia catenata*  
var. *catenata* sensu W.M.Curtis (1979) = **Caladenia carnea** & **C. sylvicola** 4a:106  
var. *exigua* sensu W.M.Curtis (1979) = **Caladenia alata** 4a:107  
var. *gigantea* sensu W.M.Curtis (1979), identity unresolved 4a:107  
var. *minor* sensu W.M.Curtis (1979) = **Caladenia mentiensi** 4a:106
- e **Caladenia caudata** Nicholls, Vict. Nat. 64: 231 (1948) 4a:101  
**Caladenia clavigera** A.Cunn. ex Lindley, Gen. Sp. Orchid. Pl. 422 (1840) 4a:105  
**Caladenia congesta** R.Br., Prodr. 324 (1810) 4a:112  
*Caladenia corynepetala* D.L.Jones = **Caladenia dilatata** 4a:100
- e **Caladenia cracens** D.L.Jones, Muelleria 9: 46 (1996)  
*Caladenia cucullata* sensu W.M.Curtis (1979) = **Caladenia atrata** & **C. gracilis** 4a:111  
*Caladenia deformis* R.Br. = **Pheladenia deformis** 4a:113
- e **Caladenia dienema** D.L.Jones, Austral. Orchid Res. 3: 28 (1998)  
**Caladenia dilatata** R.Br., Prodr. 325 (1810) 4a:100  
*Caladenia dimorpha* sensu W.M.Curtis (1979) = ?**Caladenia gracilis** 4a:109
- e **Caladenia echidnachila** Nicholls, Pap. Proc. R. Soc. Tasm. for 1932: 13, t.6 (1933) 4a:101  
**Caladenia filamentosa** R.Br., Prodr. 324 (1810) 4a:99  
**Caladenia fuscata** (Rchb.f.) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 25 (1989)  
**Caladenia gracilis** R.Br., Prodr. 324 (1810) 4a:110
- e **Caladenia helvina** D.L.Jones, Austral. Orchid Res. 2: 26, f.30 (1991)  
*Caladenia iridescens* sensu W.M.Curtis (1979) = **Caladenia transitoria** 4a:112  
**Caladenia latifolia** R.Br., Prodr. 324 (1810) 4a:105
- e **Caladenia lindleyana** (Rchb.f.) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 27 (1989)  
*Caladenia longii* R.S.Rogers = **Caladenia angustata** 4a:108  
*Caladenia lyallii* sensu W.M.Curtis (1979) = **Caladenia cracens** & **C. alpina** 4a:111  
**Caladenia mentiensi** D.L.Jones, Austral. Orchid Res. 3: 35 (1998)  
*Caladenia menziesii* R.Br. = **Leptoceras menziesii** 4a:99
- e **Caladenia pallida** Lindley, Gen. Sp. Orchid. Pl. 421 (1840) 4a:104  
**Caladenia patersonii** R.Br., Prodr. 324 (1810) 4a:100  
*Caladenia praecox* sensu W.M.Curtis (1979) = ?**Caladenia angustata** 4a:109  
**Caladenia prolata** D.L.Jones, Austral. Orchid Res. 2: 30 (1991)  
**Caladenia pusilla** W.M.Curtis, Stud. Fl. Tasm. 4a: 133 (1980) 4a:107  
*Caladenia reticulata* sensu W.M.Curtis (1979) = identity unresolved (D.L.Jones pers. comm.) 4a:104
- e **Caladenia saggicola** D.L.Jones, Austral. Orchid Res. 3: 39 (1998)

|   |  |        |
|---|--|--------|
|   | Caladenia suaveolens (R.Br.) Reichb.f. = <b>Lyperanthus suaveolens</b>                                   | 4a:94  |
| e | <b>Caladenia sylvicola</b> D.L.Jones, Austral. Orchid Res. 3: 40 (1998)                                  |        |
|   | Caladenia testacea sensu Rodway (1903) = <b>Caladenia atrata</b>   |        |
| e | <b>Caladenia tonellii</b> D.L.Jones, Austral. Orchid Res. 3: 41 (1998)                                   |        |
|   | <b>Caladenia transitoria</b> D.L.Jones, Austral. Orchid Res. 3: 42 (1998)                                |        |
|   | Caladenia tutelata R.S.Rogers = <b>XCalassodia tutelata</b>  | 4a:113 |
|   | Caladenia venusta sensu Census Vasc. Pl. Tasm. edn.2 = <b>Caladenia saggicola</b>                        |        |
|   | <b>Caladenia vulgaris</b> D.L.Jones, Austral. Orchid Res. 2: 34 (1991)                                   |        |
|   | <b>XCalassodia tutelata</b> (R.S.Rogers) M.A.Clem., Austral. Orchid Res. 1: 33 (1989)                    | 4a:113 |
|   | <b>Caleana major</b> R.Br., Prodr. 329 (1810)  | 4a:87  |
|   | Caleana minor R.Br. = <b>Paracaleana minor</b>   | 4a:88  |
|   | <b>Calochilus campestris</b> R.Br., Prodr. 320 (1810)  | 4a:53  |
|   | <b>Calochilus herbaceus</b> Lindley, Gen. Sp. Orchid. Pl. 459 (1840)                                     | 4a:55  |
|   | <b>Calochilus imberbis</b> R.S.Rogers, Trans. Proc. R. Soc. S.Austral. 51: 4 (1927)                      | 4a:53  |
|   | <b>Calochilus paludosus</b> R.Br., Prodr. 320 (1810)   | 4a:56  |
|   | <b>Calochilus robertsonii</b> Benth., Fl. Austral. 6: 315 (1873)   | 4a:55  |
|   | Calonema filamentosum (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia filamentosa</b>                       | 4a:99  |
|   | Calonemorchis filamentosa (R.Br.) Szlachetko = <b>Caladenia filamentosa</b>                              | 4a:99  |
|   | Chiloglottis cornuta Hook.f. = <b>Simpliglottis cornuta</b>  | 4a:85  |
|   | Chiloglottis diphylla sensu Rodway (1903) = <b>Chiloglottis reflexa</b>                                  | 4a:86  |
|   | Chiloglottis grammata G.W.Carr = <b>Simpliglottis grammata</b>   |        |
|   | Chiloglottis gunnii Lindley = <b>Simpliglottis gunnii</b>  | 4a:84  |
|   | Chiloglottis platychila G.W.Carr = <b>Simpliglottis gunnii</b>   |        |
|   | <b>Chiloglottis reflexa</b> (Labill.) Druce, Bot. Soc. Exch. Club Brit. Isles 1916, Suppl. 2: 614 (1917) | 4a:86  |
|   | <b>Chiloglottis trapeziformis</b> Fitzg., Austral. Orchids 1(3): t.9 (1877)                              | 4a:86  |
|   | Chiloglottis Xpescottiana sensu W.M.Curtis (1979) = <b>Simpliglottis grammata</b>                        | 4a:86  |
|   | Chiloglottis triceratops D.L.Jones = <b>Simpliglottis triceratops</b>                                    |        |
|   | Chiloglottis trilabra Fitzg. = <b>Chiloglottis reflexa</b>   |        |
|   | Chiloglottis valida D.L.Jones = <b>Simpliglottis valida</b>  |        |
|   | <b>Corunastylis archeri</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13(10): 460 (2002)                | 4a:82  |
| e | <b>Corunastylis brachystachya</b> (Lindley) D.L.Jones & M.A.Clem., Orchadian 13(10): 460 (2002)          | 4a:79  |
|   | <b>Corunastylis despectans</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13(10): 460 (2002)             | 4a:76  |
| e | <b>Corunastylis firthii</b> (L.Cady) D.L.Jones & M.A.Clem., Orchadian 13(10): 461 (2002)                 | 4a:80  |
|   | <b>Corunastylis morrisii</b> (Nicholls) D.L.Jones & M.A.Clem., Orchadian 13(10): 461 (2002)              | 4a:83  |
|   | <b>Corunastylis nuda</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13(10): 461 (2002)                   | 4a:82  |
| x | <b>Corunastylis nudiscapa</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13(10): 461 (2002)              | 4a:79  |
|   | <b>Corunastylis pumila</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13(10): 461 (2002)                 | 4a:80  |
| e | <b>Corunastylis tasmanica</b> (D.L.Jones) D.L.Jones & M.A.Clem., Orchadian 13(10): 462 (2002)            | 4a:77  |
|   | <b>Corybas aconitiflorus</b> Salisb. in Salisb. & Hook., Paradis. Lond. 2: t.83 (1807)                   | 4a:123 |
|   | Corybas diemenicus (Lindley) Rchb.f. = <b>Corysanthes diemenica</b>                                      | 4a:121 |
|   | Corybas diemenicus sensu W.M.Curtis (1979) = <b>Corysanthes incurva</b>                                  | 4a:121 |
|   | Corybas dilatatus (Rupp & Nicholls) Rupp & Nicholls ex Rupp = <b>Corysanthes diemenica</b>               | 4a:121 |
|   | Corybas fimbriatus (R.Br.) Rchb.f. = <b>Corysanthes fimbriata</b>  | 4a:121 |
|   | Corybas fordhamii (Rupp) Rupp = <b>Anzybas fordhamii</b>   | 4a:124 |

|   |  |        |
|---|--|--------|
|   | <i>Corybas incurvus</i> D.L.Jones & M.A.Clem. = <b>Corysanthes incurva</b>                                     | 4a:121 |
|   | <i>Corybas unguiculatus</i> (R.Br.) Rchb.f. = <b>Anzybas unguiculatus</b>                                      | 4a:124 |
|   | <i>Corysanthes bicalcarata</i> R.Br. = <b>Corybas aconitiflorus</b>  | 4a:123 |
|   | <b>Corysanthes diemenica</b> Lindley, Gen. Sp. Orchid. Pl. 393 (1840)  | 4a:121 |
|   | <b>Corysanthes fimbriata</b> R.Br., Prodr. 328 (1810)  | 4a:121 |
|   | <b>Corysanthes incurva</b> (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem., Orchadian 13(10): 447 (2002)         | 4a:121 |
|   | <i>Corysanthes pruinosa</i> sensu Rodway (1903) = <b>Corysanthes incurva</b>                                   | 4a:121 |
|   | <b>Crangonorchis pedoglossa</b> (Fitzg.) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 67 (2002)              | 4a:13  |
|   | <b>Cryptostylis leptochila</b> F.Muell. ex Benth., Fl. Austral. 6: 334 (1873)                                  | 4a:126 |
|   | <i>Cryptostylis longifolia</i> R.Br. = <b>Cryptostylis subulata</b>  | 4a:125 |
|   | <b>Cryptostylis subulata</b> (Labill.) Rchb.f., Beitr. Syst. Pflanzenk. 15 (1871)                              | 4a:125 |
|   | <b>Cyrtostylis reniformis</b> R.Br., Prodr. 322 (1810)   | 4a:119 |
|   | <b>Cyrtostylis robusta</b> D.L.Jones & M.A.Clem., Lindleyana 2(3): 157 (1987)                                  |        |
|   | <i>Dendrobium striolatum</i> Rchb.f. = <b>Dockrillia striolata</b>   | 4a:129 |
| e | <b>Diplodium alatum</b> (Labill.) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 70 (2002)                     | 4a:20  |
|   | <b>Diplodium atrans</b> (D.L.Jones) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 70 (2002)                   | 4a:22  |
|   | <b>Diplodium decurvum</b> (R.S.Rogers) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 71 (2002)                | 4a:22  |
|   | <b>Diplodium grandiflorum</b> (R.Br.) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 71 (2002)                 | 4a:20  |
|   | <i>Dipodium punctatum</i> sensu W.M.Curtis (1979) = <b>Dipodium roseum</b>                                     | 4a:131 |
|   | <b>Dipodium roseum</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 2: 51 (1991)                                | 4a:131 |
|   | <b>Diuris chryseopsis</b> D.L.Jones, Austral. Orchid Res. 3: 74 (1998)   |        |
|   | <i>Diuris corymbosa</i> sensu W.M.Curtis (1979) = <b>Diuris orientis</b>                                       | 4a:36  |
|   | <i>Diuris flavopurpurea</i> sensu W.M.Curtis (1979) = <b>Diuris pardina</b>                                    | 4a:35  |
| e | <b>Diuris lanceolata</b> Lindley, Gen. Sp. Orchid. Pl. 508 (1840)  | 4a:34  |
|   | <i>Diuris longifolia</i> sensu W.M.Curtis (1979) = <b>Diuris orientis</b>                                      | 4a:36  |
|   | <i>Diuris maculata</i> sensu W.M.Curtis (1979) = <b>Diuris pardina</b>   | 4a:34  |
|   | <b>Diuris monticola</b> D.L.Jones, Austral. Orchid Res. 3: 76 (1998)   |        |
|   | <b>Diuris orientis</b> D.L.Jones, Austral. Orchid Res. 3: 77 (1998)  | 4a:36  |
|   | <i>Diuris palachila</i> sensu W.M.Curtis (1979) = hybrid between <b>Diuris pardina</b> & <b>D. chryseopsis</b> | 4a:35  |
|   | <b>Diuris palustris</b> Lindley, Gen. Sp. Orchid. Pl. 507 (1840)   | 4a:32  |
|   | <b>Diuris pardina</b> Lindley, Gen. Sp. Orchid. Pl. 507 (1840)   | 4a:34  |
|   | <i>Diuris pedunculata</i> sensu W.M.Curtis (1979) = <b>Diuris chryseopsis</b> pro parte maj.                   | 4a:34  |
|   | <b>Diuris sulphurea</b> R.Br., Prodr. 316 (1810)   | 4a:36  |
|   | <b>Dockrillia striolata</b> (Rchb.f.) Rauschert, Feddes Repert. 94(7-8): 447 (1983)                            | 4a:129 |
| e | subsp. <b>chrysantha</b> D.L.Jones, Austral. Orchid Res. 3: 9 (1998)   |        |
|   | subsp. <b>striolata</b>  |        |
|   | <i>Eriochilus autumnalis</i> sensu Rodway (1903) = <b>Eriochilus cucullatus</b>                                | 4a:91  |
|   | <b>Eriochilus cucullatus</b> (Labill.) Rchb.f., Beitr. Syst. Pflanzenk. 27 (1871)                              | 4a:91  |
|   | <b>Gastrodia procera</b> G.W.Carr, Indig. Flora & Fauna Assoc. Misc. Pap. 1: 22 (1991)                         |        |
|   | <b>Gastrodia sesamoides</b> R.Br., Prodr. 330 (1810)   | 4a:127 |
|   | <i>Genoplesium archeri</i> (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis archeri</b>                       | 4a:82  |
|   | <i>Genoplesium brachystachyum</i> (Lindley) D.L.Jones & M.A.Clem. = <b>Corunastylis brachystachya</b>          | 4a:79  |
|   | <i>Genoplesium despectans</i> (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis despectans</b>                 | 4a:76  |
|   | <i>Genoplesium firthii</i> (L.Cady) D.L.Jones = <b>Corunastylis firthii</b>                                    | 4a:80  |

|   |   |        |
|---|---|--------|
|   | Genoplesium morrisii (Nicholls) D.L.Jones & M.A.Clem. = <b>Corunastylis morrisii</b>                | 4a:83  |
|   | Genoplesium nudiscapum (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis nudiscapa</b>              | 4a:79  |
|   | Genoplesium nudum (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis nuda</b>                        | 4a:82  |
|   | Genoplesium pumilum (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis pumila</b>                    | 4a:80  |
|   | Genoplesium tasmanicum D.L.Jones = <b>Corunastylis tasmanica</b>                                    | 4a:77  |
|   | <b>Glossodia major</b> R.Br., Prodr. 326 (1810)   | 4a:90  |
|   | <b>Hydrorchis orbicularis</b> (R.S.Rogers) D.L.Jones & M.A.Clem., Orchadian 13(10): 462 (2002)      | 4a:60  |
|   | <b>Hymenochilus cynocephalus</b> (Fitzg.) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 74 (2002)  | 4a:26  |
|   | <b>Hymenochilus muticus</b> (R.Br.) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 74 (2002)        | 4a:25  |
| e | <b>Hymenochilus pratensis</b> (D.L.Jones) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 74 (2002)  |        |
| e | <b>Hymenochilus rubenachii</b> (D.L.Jones) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 74 (2002) |        |
| e | <b>Hymenochilus wapstrarum</b> (D.L.Jones) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 74 (2002) |        |
| e | <b>Hymenochilus ziegeleri</b> (D.L.Jones) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 74 (2002)  |        |
|   | Jonesiopsis filamentosa (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia filamentosa</b>                | 4a:99  |
|   | <b>Leptoceras menziesii</b> (R.Br.) Lindley, Gen. Sp. Orchid. Pl. 416 (1840)                        | 4a:99  |
|   | <b>Linguella nana</b> (R.Br.) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 75 (2002)              | 4a:14  |
|   | Lyperanthus nigricans R.Br. = <b>Pyrorchis nigricans</b>  | 4a:93  |
|   | <b>Lyperanthus suaveolens</b> R.Br., Prodr. 325 (1810)  | 4a:94  |
|   | <b>Microtidium atratum</b> (Lindley) D.L.Jones & M.A.Clem., Orchadian 13(10): 463 (2002)            | 4a:60  |
|   | <b>Microtis arenaria</b> Lindley, Gen. Sp. Orchid. Pl. 306 (1840)                                   | 4a:58  |
|   | Microtis atrata Lindley = <b>Microtidium atratum</b>  | 4a:60  |
|   | Microtis biloba Nicholls = <b>Microtis arenaria</b>   | 4a:58  |
|   | Microtis oblonga R.S.Rogers = <b>Microtis rara</b>  | 4a:59  |
|   | Microtis orbicularis R.S.Rogers = <b>Hydrorchis orbicularis</b>                                     | 4a:60  |
|   | <b>Microtis parviflora</b> R.Br., Prodr. 321 (1810)   | 4a:59  |
|   | Microtis porrifolia (Swartz) R.Br. ex Sprengel = <b>Microtis unifolia</b>                           | 4a:58  |
|   | <b>Microtis rara</b> R.Br., Prodr. 321 (1810)   | 4a:59  |
|   | <b>Microtis unifolia</b> (Forst.f.) Rchb.f., Beitr. Syst. Pflanzenk. 62 (1871)                      | 4a:58  |
|   | <b>Nemacianthus caudatus</b> (R.Br.) D.L.Jones & M.A.Clem., Orchadian 13(10): 441 (2002)            | 4a:117 |
| e | <b>Oligochaetochilus commutatus</b> (D.L.Jones) D.L.Szlach., Polish Bot. J. 46(1): 24 (2001)        |        |
|   | <b>Oligochaetochilus squamatus</b> (R.Br.) D.L.Szlach., Polish Bot. J. 46(1): 25 (2001)             | 4a:27  |
|   | <b>Orthoceras strictum</b> R.Br., Prodr. 317 (1810)   | 4a:30  |
|   | <b>Paracaleana minor</b> (R.Br.) Blaxell, Contr. N.S.W. Natl. Herb. 4: 281 (1972)                   | 4a:88  |
|   | Petalochilus alatus (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia alata</b>                          |        |
|   | Petalochilus atrochilus (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia atrochila</b>              |        |
|   | Petalochilus aurantiacus (R.S.Rogers) D.L.Jones & M.A.Clem. = <b>Caladenia aurantiaca</b>           |        |
|   | Petalochilus campbellii (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia campbellii</b>             |        |
|   | Petalochilus carneus (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia carnea</b>                        | 4a:106 |
|   | Petalochilus fuscatus (Rchb.f.) D.L.Jones & M.A.Clem. = <b>Caladenia fuscata</b>                    |        |
|   | Petalochilus mentiensi (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia mentiensi</b>               |        |
|   | Petalochilus prolatus (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia prolata</b>                  |        |
|   | Petalochilus pusillus (W.M.Curtis) D.L.Jones & M.A.Clem. = <b>Caladenia pusilla</b>                 | 4a:107 |
|   | Petalochilus sylvicola (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia sylvicola</b>               |        |
|   | Petalochilus tonellii (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia tonellii</b>                 |        |

|    |  |        |
|----|--|--------|
|    | Petalochilus vulgaris (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia vulgaris</b>                |        |
|    | <b>Pheladenia deformis</b> (R.Br.) D.L.Jones & M.A.Clem., Orchadian 13(9): 411 (2001)              | 4a:113 |
|    | <b>Plumatichilos plumosum</b> (Cady) D.L.Szlach., Polish Bot. J. 46(1): 23 (2001)                  | 4a:24  |
|    | <b>Plumatichilos tasmanicum</b> (D.L.Jones) D.L.Szlach., Polish Bot. J. 46(1): 23 (2001)           |        |
|    | Prasophyllum album sensu W.M.Curtis (1979) = Correct identity unresolved (D.L.Jones 1998)          | 4a:69  |
|    | Prasophyllum alpestre (Tasmanian plants) = <b>Prasophyllum mimulum</b>                             |        |
| e  | <b>Prasophyllum alpinum</b> R.Br., Prodr. 318 (1810)   | 4a:75  |
| e  | <b>Prasophyllum amoenum</b> D.L.Jones, Austral. Orchid Res. 3: 99 (1998)                           |        |
| e  | <b>Prasophyllum apoxychilum</b> D.L.Jones, Austral. Orchid Res. 3: 100 (1998)                      |        |
|    | Prasophyllum archeri Hook.f. = <b>Corunastylis archeri</b>   | 4a:82  |
|    | <b>Prasophyllum australe</b> R.Br., Prodr. 318 (1810)  | 4a:66  |
|    | Prasophyllum beagleholei Nicholls = <b>Corunastylis nuda</b>                                       | 4a:82  |
|    | Prasophyllum brachystachyum Lindley = <b>Corunastylis brachystachya</b>                            | 4a:79  |
|    | Prasophyllum brainei R.S.Rogers = <b>Prasophyllum lindleyanum</b>                                  | 4a:74  |
|    | <b>Prasophyllum brevilabre</b> (Lindley) Hook.f., Fl. Tasm. 2: 11, t.110A (1858)                   | 4a:68  |
|    | Prasophyllum buftonianum J.H.Willis = <b>Corunastylis pumila</b>                                   | 4a:80  |
| e  | <b>Prasophyllum castaneum</b> D.L.Jones, Austral. Orchid Res. 3: 102 (1998)                        |        |
| e  | <b>Prasophyllum concinnum</b> Nicholls, Vict. Nat. 64: 232 (1948)                                  | 4a:73  |
|    | Prasophyllum correctum D.L.Jones (Tasmanian plants) = <b>Prasophyllum incorrectum</b>              |        |
| n  | <b>Prasophyllum crebriflorum</b> D.L.Jones, Muelleria 18: 103 (2003)                               |        |
|    | Prasophyllum despectans Hook.f. = <b>Corunastylis despectans</b>                                   | 4a:76  |
|    | <b>Prasophyllum elatum</b> R.Br., Prodr. 318 (1810)  | 4a:67  |
| e  | <b>Prasophyllum favonium</b> D.L.Jones, Austral. Orchid Res. 3: 104 (1998)                         |        |
|    | Prasophyllum firthii L.Cady = <b>Corunastylis firthii</b>  | 4a:80  |
|    | Prasophyllum fitzgeraldii R.S.Rogers & Maiden, not present in Tasmania, formerly recorded in error |        |
|    | <b>Prasophyllum flavum</b> R.Br., Prodr. 318 (1810)  | 4a:66  |
|    | Prasophyllum frenchii sensu W.M.Curtis (1979), identity unresolved                                 | 4a:73  |
|    | Prasophyllum fuscum sensu W.M.Curtis (1979), identity unresolved (D.L.Jones pers. comm.)           | 4a:75  |
|    | Prasophyllum gracile sensu W.M.Curtis (1979), identity unresolved (D.L.Jones pers. comm.)          | 4a:72  |
| ne | <b>Prasophyllum incorrectum</b> D.L.Jones, Muelleria 18: 107 (2003)                                |        |
| e  | <b>Prasophyllum incurvum</b> D.L.Jones, Austral. Orchid Res. 3: 106 (1998)                         |        |
|    | Prasophyllum intricatum C.Stuart ex Benth. = <b>Corunastylis archeri</b>                           | 4a:82  |
|    | <b>Prasophyllum lindleyanum</b> Rchb.f., Beitr. Syst. Pflanzenk. 3: 58 (1871)                      | 4a:74  |
| e  | <b>Prasophyllum milfordense</b> D.L.Jones, Austral. Orchid Res. 3: 107 (1998)                      |        |
| ne | <b>Prasophyllum mimulum</b> D.L.Jones, Orchadian 14: 372 (2004)                                    |        |
|    | <b>Prasophyllum montanum</b> R.Bates & D.L.Jones, Austral. Orchid Res. 2: 79 (1991)                |        |
|    | Prasophyllum morrisii Nicholls = <b>Corunastylis morrisii</b>                                      | 4a:83  |
|    | Prasophyllum nigricans sensu W.M.Curtis (1979) = <b>Corunastylis tasmanica</b>                     | 4a:77  |
|    | Prasophyllum nudiscapum Hook.f. = <b>Corunastylis nudiscapa</b>                                    |        |
|    | Prasophyllum odoratum sensu W.M.Curtis (1979), identity unresolved                                 | 4a:68  |
| e  | <b>Prasophyllum olidum</b> D.L.Jones, Austral. Orchid Res. 3: 108 (1998)                           |        |
|    | Prasophyllum parviflorum sensu W.M.Curtis (1979), identity unresolved (D.L.Jones pers. comm.)      | 4a:74  |
|    | Prasophyllum patens sensu Rodway (1903) = <b>Prasophyllum truncatum</b>                            | 4a:71  |
| e  | <b>Prasophyllum perangustum</b> D.L.Jones, Austral. Orchid Res. 3: 109 (1998)                      | 4a:72  |



- e **Prasophyllum pulchellum** D.L.Jones, Austral. Orchid Res. 3: 110 (1998)  
**Prasophyllum pyriforme** E.Coleman, Vict. Nat. 49: 195 (1932)
- e **Prasophyllum robustum** (Nicholls) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 117 (1989)  
Prasophyllum rogersii sensu W.M.Curtis (1979) = **Prasophyllum perangustum** 4a:72
- e **Prasophyllum rostratum** Lindley, Gen. Sp. Orchid. Pl. 516 (1840)  
Prasophyllum rufum sensu W.M.Curtis (1979) = **Corunastylis tasmanica** 4a:79
- e **Prasophyllum secutum** D.L.Jones, Austral. Orchid Res. 3: 113 (1998)  
**Prasophyllum sphacelatum** D.L.Jones, Muelleria 9: 59 (1996)
- e **Prasophyllum stellatum** D.L.Jones, Austral. Orchid Res. 3: 115 (1998)  
Prasophyllum suttonii sensu W.M.Curtis (1979) = **Prasophyllum mimulum** 4a:69  
**Prasophyllum tadgellianum** R.S.Rogers, Trans. Proc. R. Soc. S.Austral. 47: 338 (1923)
- ne **Prasophyllum taphanyx** D.L.Jones, Orchadian 14: 373 (2004)
- e **Prasophyllum truncatum** Lindley, Gen. Sp. Orchid. Pl. 513 (1840) 4a:71
- e **Prasophyllum tunbridgense** D.L.Jones, Austral. Orchid Res. 3: 117 (1998)  
Prasophyllum uroglossum sensu W.M.Curtis (1979), identity unresolved (D.L.Jones pers. comm.) 4a:76  
Pterostylis alata (Labill.) Rchb.f. = **Diplodium alatum** 4a:20  
Pterostylis alpina sensu W.M.Curtis (1979) = **Pterostylis scabrida** 4a:19  
Pterostylis aphylla Lindley = **Speculantha aphylla** 4a:24  
Pterostylis atrans D.L.Jones = **Diplodium atrans** 4a:22  
Pterostylis atriola D.L.Jones = **Speculantha atriola**  
Pterostylis barbata sensu Rodway (1903) = **Plumatichilos plumosum** 4a:24  
Pterostylis commutata D.L.Jones = **Oligochaetochilus commutatus**  
Pterostylis concinna R.Br. = **Taurantha concinna** 4a:11  
**Pterostylis cucullata** R.Br., Prodr. 327 (1810) 4a:15  
**Pterostylis curta** R.Br., Prodr. 326 (1810) 4a:12  
Pterostylis cycnocephala Fitzg. = **Hymenochilus cycnocephalus** 4a:26  
Pterostylis decurva R.S.Rogers = **Diplodium decurvum** 4a:22
- e **Pterostylis dubia** R.Br., Prodr. 328 (1810) 4a:16  
**Pterostylis falcata** R.S.Rogers, Proc. R. Soc. Vict. ser.2, 28(1): 106, t.9 (1915) 4a:17  
**Pterostylis foliata** Hook.f. Fl. Nov.-Zel. 1: 249 (1853) 4a:19
- e **Pterostylis furcata** Lindley, Gen. Sp. Orchid. Pl. 390 (1840) 4a:17  
Pterostylis grandiflora R.Br. = **Diplodium grandiflorum** 4a:20  
Pterostylis Xingens sensu W.M.Curtis (1979) = **Pterostylis foliata** 4a:12  
Pterostylis longifolia sensu W.M.Curtis (1979) = **Bunochilus melagrammus** & **B. williamsonii** 4a:27  
Pterostylis melagramma D.L.Jones = **Bunochilus melagrammus** 4a:27  
Pterostylis mutica R.Br. = **Hymenochilus muticus** 4a:25  
Pterostylis nana R.Br. = **Linguella nana** 4a:14  
**Pterostylis nutans** R.Br., Prodr. 327 (1810) 4a:13  
Pterostylis obtusa sensu W.M.Curtis (1979) = **Diplodium atrans** 4a:22  
Pterostylis parviflora R.Br. = **Speculantha parviflora** 4a:23  
Pterostylis pedoglossa Fitzg. = **Crangonorchis pedoglossa** 4a:13  
**Pterostylis pedunculata** R.Br., Prodr. 327 (1810) 4a:14  
Pterostylis plumosa L.Cady = **Plumatichilos plumosum** 4a:24  
Pterostylis praecox sensu Rodway (1903) = **Diplodium alatum** 4a:20

|    |   |        |
|----|---|--------|
|    | Pterostylis pratensis D.L.Jones = <b>Hymenochilus pratensis</b>                                   |        |
|    | Pterostylis rubenachii D.L.Jones = <b>Hymenochilus rubenachii</b>                                 |        |
|    | Pterostylis rufa sensu W.M.Curtis (1979) = <b>Oligochaetochilus squamatus</b>                     | 4a:27  |
|    | Pterostylis sanguinea D.L.Jones & M.A.Clem. = <b>Urochilus sanguineus</b>                         | 4a:28  |
| e  | <b>Pterostylis scabrida</b> Lindley, Gen. Sp. Orchid. Pl. 389 (1840)                              | 4a:19  |
|    | Pterostylis squamata R.Br. = <b>Oligochaetochilus squamatus</b>                                   | 4a:27  |
|    | Pterostylis stenochila D.L.Jones = <b>Bunochilus stenochilus</b>                                  |        |
|    | Pterostylis tasmanica D.L.Jones = <b>Plumatichilos tasmanicum</b>                                 |        |
|    | Pterostylis Xtoveyana Ewart & Sharman = <b>XTaurodium toveyanum</b>                               | 4a:21  |
|    | Pterostylis tunstallii D.L.Jones & M.A.Clem. = <b>Bunochilus tunstallii</b>                       |        |
|    | Pterostylis uliginosa D.L.Jones = <b>Speculantha uliginosa</b>                                    |        |
|    | Pterostylis vittata sensu W.M.Curtis (1979) = <b>Urochilus sanguineus</b>                         | 4a:28  |
|    | Pterostylis wapstrarum D.L.Jones = <b>Hymenochilus wapstrarum</b>                                 |        |
|    | Pterostylis williamsonii D.L.Jones = <b>Bunochilus williamsonii</b>                               |        |
|    | Pterostylis ziegeleri D.L.Jones = <b>Hymenochilus ziegeleri</b>                                   |        |
|    | <b>Pyrorchis nigricans</b> (R.Br.) D.L.Jones & M.A.Clem., Phytologia 77(6): 449 (1994)            | 4a:93  |
|    | <b>Sarcochilus australis</b> (Lindley) Rchb.f. in C.Muell., Ann. Bot. Syst. 6: 501 (1863)         | 4a:131 |
|    | Sarcochilus parviflorus sensu Rodway (1903) = <b>Sarcochilus australis</b>                        | 4a:131 |
| n  | <b>Simpliglottis cornuta</b> (Hook.f.) Szlach., Polish Bot. J. 46(1): 13 (2001)                   | 4a:85  |
| e  | <b>Simpliglottis grammata</b> (G.W.Carr) Jeanes, Muelleria 16: 82 (2002)                          |        |
| ne | <b>Simpliglottis gunnii</b> (Lindley) Szlach., Polish Bot. J. 46(1): 13 (2001)                    | 4a:84  |
| e  | <b>Simpliglottis triceratops</b> (D.L.Jones) Jeanes, Muelleria 16: 82 (2002)                      |        |
| n  | <b>Simpliglottis valida</b> (D.L.Jones) Szlach., Polish Bot. J. 46(1): 14 (2001)                  |        |
| e  | <b>Speculantha aphylla</b> (Lindley) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 82 (2002)     | 4a:24  |
| e  | <b>Speculantha atriola</b> (D.L.Jones) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 82 (2002)   |        |
|    | <b>Speculantha parviflora</b> (R.Br.) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 83 (2002)    | 4a:23  |
|    | <b>Speculantha uliginosa</b> (D.L.Jones) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 83 (2002) |        |
|    | Spiculaea huntiana (F.Muell.) Schltr. = <b>Arthrochilus huntianus</b>                             | 4a:89  |
|    | <b>Spiranthes australis</b> (R.Br.) Lindley, Bot. Reg. 10, subt.823 (1824)                        | 4a:128 |
|    | Spiranthes sinensis subsp. australis (R.Br.) Hara & Kitam. = <b>Spiranthes australis</b>          | 4a:128 |
|    | Stegostyla alpina (R.S.Rogers) D.L.Jones & M.A.Clem. = <b>Caladenia alpina</b>                    | 4a:111 |
|    | Stegostyla angustata (Lindley) D.L.Jones & M.A.Clem. = <b>Caladenia angustata</b>                 | 4a:108 |
|    | Stegostyla atrata (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia atrata</b>                     |        |
|    | Stegostyla congesta (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia congesta</b>                     | 4a:112 |
|    | Stegostyla cracens (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia cracens</b>                   |        |
|    | Stegostyla gracilis (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia gracilis</b>                     | 4a:110 |
|    | Stegostyla transitoria (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia transitoria</b>           |        |
|    | <b>Taurantha concinna</b> (R.Br.) D.L.Jones & M.A.Clem., Austral. Orchid Res. 4: 85 (2002)        | 4a:11  |
|    | <b>XTaurodium toveyanum</b> (Ewart&Sharman) D.L.Jones&M.A.Clem., Austral. Orchid Res.4:87 (2002)  | 4a:21  |
| e  | <b>Thelymitra aggericola</b> D.L.Jones, The Orchadian 12(11): 517 (1999)                          |        |
|    | <b>Thelymitra antennifera</b> (Lindley) Hook.f., Fl. Tasm. 2: 4, t.101A (1858)                    | 4a:51  |
|    | <b>Thelymitra arenaria</b> Lindley, Gen. Sp. Orchid. Pl. 519 (1840)                               |        |
|    | <b>Thelymitra aristata</b> Lindley, Gen. Sp. Orchid. Pl. 521 (1840)                               | 4a:42  |
|    | Thelymitra azurea sensu D.L.Jones et al. (1999) = <b>Thelymitra jonesii</b>                       |        |

|   |  |       |
|---|--|-------|
|   | <b>Thelymitra benthamiana</b> Rchb.f., Beitr. Syst. Pflanzenk. 55 (1871)                     |       |
| n | <b>Thelymitra bracteata</b> J.Z.Weber ex Jeanes, Muelleria 19: 43 (2004)                     |       |
| n | <b>Thelymitra brevifolia</b> Jeanes, Muelleria 19: 30 (2004)                                 |       |
|   | Thelymitra canaliculata sensu W.M.Curtis (1979) = <b>Thelymitra jonesii</b>                  | 4a:46 |
|   | <b>Thelymitra carnea</b> R.Br., Prodr. 314 (1810)  | 4a:50 |
|   | Thelymitra chasmogama sensu W.M.Curtis (1979) = a plant of uncertain hybrid origin           | 4a:47 |
|   | <b>Thelymitra circumsepta</b> Fitzg., Austral. Orchids 1(4): t.1 (1878)                      | 4a:48 |
|   | <b>Thelymitra cyanea</b> (Lindley) Benth., Fl. Austral. 6: 323 (1873)                        | 4a:49 |
|   | Thelymitra decora sensu Census Vasc. Pl. Tasm. edn.2 = <b>Thelymitra simulata</b>            |       |
|   | <b>Thelymitra erosa</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 184 (1998)            |       |
| n | <b>Thelymitra exigua</b> Jeanes, Muelleria 19: 28 (2004)                                     |       |
|   | <b>Thelymitra flexuosa</b> Endl. in Endl. & Fenzl, Nov. Stirp. Dec. 3: 23 (1839)             | 4a:51 |
|   | <b>Thelymitra holmesii</b> Nicholls, Vict. Nat. 49: 263 (1933)                               | 4a:43 |
| e | <b>Thelymitra imbricata</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 186 (1998)        | 4a:43 |
|   | <b>Thelymitra improcera</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 187 (1998)        | 4a:46 |
| n | <b>Thelymitra inflata</b> Jeanes, Muelleria 19: 71 (2004)                                    |       |
|   | <b>Thelymitra Xirregularis</b> Nicholls, Vict. Nat. 63: 126 (1946)                           | 4a:48 |
|   | <b>Thelymitra ixiooides</b> Swartz, Ksvenska Vet. Akad. Handl. 21: 253 (1800)                | 4a:44 |
| e | <b>Thelymitra jonesii</b> Jeanes, Muelleria 15: 81 (2001)                                    | 4a:46 |
|   | <b>Thelymitra juncifolia</b> Lindley, Gen. Sp. Orchid. Pl. 522 (1840)                        |       |
|   | <b>Thelymitra longiloba</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 191 (1998)        | 4a:46 |
| n | <b>Thelymitra lucida</b> Jeanes, Muelleria 19: 70 (2004)                                     |       |
|   | Thelymitra luteocilium sensu W.M.Curtis (1979) = <b>Thelymitra rubra</b>                     | 4a:47 |
|   | Thelymitra Xmacmillanii sensu W.M.Curtis (1979) = a hybrid involving <b>Thelymitra nuda</b>  | 4a:49 |
|   | <b>Thelymitra malvina</b> M.A.Clem., D.L.Jones & Molloy, Austral. Orchid Res. 1: 141 (1989)  |       |
|   | Thelymitra media sensu W.M.Curtis (1979) = <b>Thelymitra improcera</b> & <b>T. longiloba</b> | 4a:46 |
|   | Thelymitra megcalyptra sensu W.M.Curtis (1979) = <b>Thelymitra imbricata</b>                 | 4a:43 |
|   | <b>Thelymitra Xmerraniae</b> Nicholls, Vict. Nat. 46: 139 (1929)                             |       |
|   | <b>Thelymitra mucida</b> Fitzg., Gard. Chron. 17: 495 (1882)                                 | 4a:46 |
|   | <b>Thelymitra nuda</b> R.Br., Prodr. 314 (1810)  | 4a:44 |
|   | <b>Thelymitra pauciflora</b> R.Br., Prodr. 314 (1810)  | 4a:42 |
|   | Thelymitra pauciflora var. holmesii (Nicholls) Nicholls = <b>Thelymitra holmesii</b>         | 4a:43 |
| n | <b>Thelymitra peniculata</b> Jeanes, Muelleria 19: 50 (2004)                                 |       |
| e | <b>Thelymitra polychroma</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 193 (1998)       |       |
|   | Thelymitra pulchella Hook.f., not present in Tasmania, formerly recorded in error            |       |
|   | Thelymitra resecta sensu W.M.Curtis = <b>Thelymitra circumsepta</b>                          | 4a:48 |
|   | <b>Thelymitra rubra</b> Fitzg., Gard. Chron. 17: 495 (1882)                                  | 4a:50 |
| e | <b>Thelymitra silena</b> D.L.Jones, The Orchadian 12(11): 518 (1999)                         |       |
|   | <b>Thelymitra simulata</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 195 (1998)         |       |
| e | <b>Thelymitra spadicea</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 196 (1998)         |       |
| e | <b>Thelymitra sparsa</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 197 (1998)           |       |
|   | <b>Thelymitra Xtruncata</b> R.S.Rogers, Trans. Proc. R. Soc. S.Austral. 41: 343, t.17 (1917) | 4a:45 |
|   | Thelymitra venosa sensu W.M.Curtis (1979) = <b>Thelymitra cyanea</b>                         | 4a:48 |
| e | <b>Thelymitra versicolor</b> Lindley, Gen. Sp. Orchid. Pl. 520 (1840)                        |       |

|    |   |        |
|----|---|--------|
| ne | <b>Thelymitra viridis</b> Jeanes, Muelleria 19: 36 (2004)   |        |
|    | <b>Thynninorchis huntiana</b> (F.Muell.) D.L.Jones & M.A.Clem., Orchadian 13(10): 457 (2002)            | 4a:89  |
| e  | <b>Thynninorchis nothofagicola</b> (D.L.Jones) D.L.Jones & M.A.Clem., Orchadian 13(10): 457 (2002)      |        |
| t  | <b>Townsonia viridis</b> (Hook.f.) Schltr., Feddes Repert. 9: 250 (1911)                                | 4a:118 |
|    | <b>Urochilussanguineus</b> (D.L.Jones&M.A.Clem.)D.L.Jones&M.A.Clem.,Austral.Orchid Res.4:87(2002) 4a:28 |        |

**POACEAE (GRAMINEAE)**

|    |   |        |
|----|---|--------|
| i  | <b>Achnatherum caudatum</b> (Trin.) S.W.L.Jacobs & J.Everett, Telopea 6(4): 582 (1966)              | 4b:192 |
| i  | <b>XAgropogon littoralis</b> (Smith) C.E.Hubb., J. Ecol. 33: 333 (1946)                             | 4b:283 |
|    | Agropyron elongatum (Host) P.Beauv. = <b>Lophopyrum ponticum</b>                                    | 4b:295 |
|    | Agropyron junceiforme (A.& D.Löve) A.& D.Löve = <b>Thinopyrum junceiforme</b>                       | 4b:293 |
|    | Agropyron junceum (L.) P.Beauv. = <b>Thinopyrum junceiforme</b>                                     | 4b:293 |
|    | Agropyron pectinatum (Labill.) P.Beauv. = <b>Australopyrum pectinatum</b>                           | 4b:301 |
|    | Agropyron repens (L.) P.Beauv. = <b>Elytrigia repens</b>  | 4b:295 |
|    | Agropyron scabrum (Labill.) P.Beauv. = <b>Elymus scaber</b>   | 4b:296 |
|    | Agropyron velutinum Nees = <b>Australopyrum velutinum</b>   | 4b:301 |
|    | Agrostis aemula R.Br. = <b>Lachnagrostis aemula</b>   | 4b:258 |
|    | var. setifolia (Hook.f.) Vickery = <b>Lachnagrostis punicea</b> subsp. <b>punicea</b>               | 4b:259 |
|    | Agrostis aequata Nees = <b>Lachnagrostis aequata</b>  | 4b:262 |
|    | <b>Agrostis australiensis</b> Mez, Feddes Repert. 17: 302 (1921)                                    | 4b:257 |
|    | Agrostis sp. aff. australiensis sensu Curtis & Morris (1994) = <b>Agrostis diemenica</b>            | 4b:257 |
|    | Agrostis australiensis sensu Townrow (1969) = <b>Lachnagrostis aequata</b>                          | 4b:262 |
|    | Agrostis avenacea J.Gmelin = <b>Lachnagrostis filiformis</b>  | 4b:259 |
|    | <b>Agrostis bettyae</b> S.W.L.Jacobs, Telopea 9(3): 679 (2001)                                      |        |
|    | Agrostis billardierei R.Br. = <b>Lachnagrostis billardierei</b>                                     | 4b:260 |
|    | var. collicola D.I.Morris = <b>Lachnagrostis collicola</b>  | 4b:261 |
|    | var. filifolia Vickery = <b>Lachnagrostis punicea</b> subsp. <b>filifolia</b>                       | 4b:261 |
|    | var. robusta Vickery = <b>Lachnagrostis robusta</b>   | 4b:261 |
|    | var. tenuiseta D.I.Morris = <b>Lachnagrostis billardierei</b> subsp. <b>tenuiseta</b>               | 4b:261 |
| i  | <b>Agrostis capillaris</b> L., Sp. Pl. 1: 62 (1753)   | 4b:264 |
| i  | var. <b>aristata</b> (Parn.) Druce, Fl. Oxford. edn.2, 474 (1927)                                   |        |
| i  | var. <b>capillaris</b>  |        |
| ne | <b>Agrostis diemenica</b> D.I.Morris, Telopea 10(3): 765 (2004)                                     | 4b:257 |
| i  | <b>Agrostis gigantea</b> Roth, Tent. Fl. Germ. 1: 31 (1788)   | 4b:264 |
|    | Agrostis sp. aff. hiemalis sensu Curtis & Morris (1994) = <b>Agrostis propinqua</b>                 | 4b:258 |
|    | <b>Agrostis joyceae</b> S.W.L.Jacobs, Telopea 9(3): 680 (2001)                                      |        |
|    | Agrostis lacunarum D.I.Morris = <b>Lachnagrostis lacunarum</b>                                      | 4b:260 |
|    | <b>Agrostis muelleriana</b> Vickery, Contr. N.S.W. Natl. Herb. 1: 103 (1941)                        | 4b:256 |
|    | <b>Agrostis parviflora</b> R.Br., Prodr. 170 (1810)   | 4b:256 |
|    | Agrostis sp. aff. parviflora sensu Curtis & Morris (1994) = <b>Agrostis thompsoniae</b>             | 4b:257 |
|    | <b>Agrostis propinqua</b> S.W.L.Jacobs, Telopea 9(3): 681 (2001)                                    |        |
|    | Agrostis rudis sensu Curtis & Morris (1994) = <b>Lachnagrostis</b> sp.                              | 4b:261 |
|    | Agrostis sp. aff. scabra sensu Curtis & Morris (1994) = <b>Agrostis propinqua</b>                   | 4b:258 |
|    | Agrostis scabra sensu Rodway (1903) = <b>Agrostis</b> sp. aff. <b>scabra</b> & <b>A. parviflora</b> |        |

|   |   |        |
|---|---|--------|
| i | <b>Agrostis stolonifera</b> L., Sp. Pl. 1: 62 (1753)  | 4b:262 |
|   | <i>Agrostis semiverticillata</i> (Forssk.) C.Chr. = <b>Polypogon viridis</b>  | 4b:283 |
|   | <i>Agrostis tenuis</i> Sibth. = <b>Agrostis capillaris</b>  | 4b:264 |
|   | <b>Agrostis thompsoniae</b> S.W.L.Jacobs, <i>Telopea</i> 9(3): 682 (2001)   |        |
|   | <b>Agrostis venusta</b> Trin., Mém. Acad. Imp. Sci. St-Pétersbourg, ser. 6 Sci. Math., Pt. 2 Sci. Nat. 4:340 (1841) | 4b:256 |
|   | <i>Agrostis viridis</i> Gouan = <b>Polypogon viridis</b>  | 4b:283 |
|   | <i>Agrostis vulgaris</i> With. = <b>Agrostis capillaris</b>   | 4b:264 |
| i | <b>Aira caryophylla</b> L., Sp. Pl. 1: 66 (1753)  | 4b:246 |
|   | <i>Aira elegans</i> Willd. ex Gaudin = <b>Aira elegantissima</b>  | 4b:247 |
| i | <b>Aira elegantissima</b> Schur, Verh. Mitth. Siebenberg. Vereins Naturwiss. Hermannstadt 4: 85 (1853)              | 4b:247 |
| i | <b>Aira praecox</b> L., Sp. Pl. 1: 65 (1753)  | 4b:246 |
| i | <b>Alopecurus geniculatus</b> L., Sp. Pl. 1: 60 (1753)  | 4b:284 |
| i | <b>Alopecurus myosuroides</b> Huds., Fl. Angl. 23 (1762)  | 4b:285 |
| i | <b>Alopecurus pratensis</b> L., Sp. Pl. 1: 60 (1753)  | 4b:284 |
| i | <b>Ammophila arenaria</b> (L.) Link, Hort. Berol. 1: 105 (1827)   | 4b:274 |
|   | <b>Amphibromus archeri</b> (Hook.f.) P.Morris, Vict. Nat. 51: 146, t.26 figs.2 & 3 (1934)                           | 4b:232 |
|   | <b>Amphibromus fluitans</b> Kirk, Trans. Proc. N.Z. Inst. 16: 374 (1884)  |        |
|   | <b>Amphibromus macrorhinus</b> S.W.L.Jacobs & L.Lapinpuro, <i>Telopea</i> 2(6): 723 (1986)                          | 4b:233 |
|   | <b>Amphibromus neesii</b> Steudel, Syn. Pl. Glum. 1: 328 (1854)   | 4b:234 |
|   | <b>Amphibromus nervosus</b> (Hook.f.) Baill., Hist. Pl. 12: 203 (1894)  |        |
|   | <b>Amphibromus recurvatus</b> Swallen, Amer. J. Bot. 18: 415 (1931)   | 4b:232 |
|   | <b>Amphibromus sinuatus</b> S.W.L.Jacobs & L.Lapinpuro, <i>Telopea</i> 2(6): 727 (1986)                             | 4b:233 |
|   | <i>Anthistiria ciliata</i> sensu Rodway (1903) = <b>Themeda triandra</b>  | 4b:358 |
| i | <b>Anthoxanthum odoratum</b> L., Sp. Pl. 1: 28 (1753)   | 4b:250 |
| i | <b>Aristida benthamii</b> Henrard, Meded. Rijks Herb. Leiden 58A: 246, t.117 (1932)                                 | 4b:323 |
|   | <i>Aristida jerichoensis</i> sensu Census Vasc. Pl. Tasm. edn.1 = <b>Aristida benthamii</b>                         | 4b:323 |
|   | <i>Arrhenatherum avenaceum</i> sensu Rodway (1903) = <b>Arrhenatherum elatius</b>                                   | 4b:234 |
| i | <b>Arrhenatherum elatius</b> (L.) J. & C.Presl, Fl. Cech. 17 (1819)   |        |
| i | var. <b>bulbosum</b> (Willd.) Spenner, Fl. Friburg. 1: 113 (1825)   | 4b:234 |
| e | <b>Australopyrum pectinatum</b> (Labill.) A.Löve, Feddes Repert. 95: 443 (1984)                                     | 4b:301 |
|   | <b>Australopyrum velutinum</b> (Nees) B.Simon, <i>Austrobaileya</i> 2: 241 (1986)                                   | 4b:301 |
|   | <b>Austrodanthonia caespitosa</b> (Gaudich.) H.P.Linder, <i>Telopea</i> 7(3): 271 (1997)                            | 4b:312 |
|   | <b>Austrodanthonia carphoides</b> (Benth.) H.P.Linder, <i>Telopea</i> 7(3): 271 (1997)                              | 4b:306 |
| e | <b>Austrodanthonia diemenica</b> (D.I.Morris) H.P.Linder, <i>Telopea</i> 7(3): 271 (1997)                           | 4b:311 |
|   | <b>Austrodanthonia geniculata</b> (J.M.Black) H.P.Linder, <i>Telopea</i> 7(3): 271 (1997)                           | 4b:308 |
|   | <b>Austrodanthonia laevis</b> (Vickery) H.P.Linder, <i>Telopea</i> 7(3): 272 (1997)                                 | 4b:312 |
|   | <b>Austrodanthonia penicillata</b> (Labill.) H.P.Linder, <i>Telopea</i> 7(3): 272 (1997)                            | 4b:313 |
|   | <b>Austrodanthonia pilosa</b> (R.Br.) H.P.Linder, <i>Telopea</i> 7(3): 272 (1997)                                   | 4b:313 |
| e | <b>Austrodanthonia popinensis</b> (D.I.Morris) H.P.Linder, <i>Telopea</i> 7(3): 273 (1997)                          | 4b:316 |
|   | <b>Austrodanthonia procera</b> (Vickery) S.W.L.Jacobs, <i>Telopea</i> 9(3): 741 (2001)                              | 4b:312 |
|   | <b>Austrodanthonia racemosa</b> (R.Br.) H.P.Linder, <i>Telopea</i> 7(3): 273 (1997)                                 | 4b:314 |
| e | <b>Austrodanthonia remota</b> (D.I.Morris) H.P.Linder, <i>Telopea</i> 7(3): 273 (1997)                              | 4b:316 |
|   | <b>Austrodanthonia setacea</b> (R.Br.) H.P.Linder, <i>Telopea</i> 7(3): 273 (1997)                                  | 4b:311 |
|   | <b>Austrodanthonia tenuior</b> (Steud.) H.P.Linder, <i>Telopea</i> 7(3): 273 (1997)                                 | 4b:310 |

|   |  |        |
|---|--|--------|
|   | <b>Austrofestuca hookeriana</b> (F.Muell. ex Hook.f.) S.W.L.Jacobs, <i>Telopea</i> 3(4): 602 (1990)                  | 4b:223 |
|   | <b>Austrofestuca littoralis</b> (Labill.) E.Alexeev, <i>Byull. Mosk. Obshch. Ispyt. Prir., Biol.</i> 81(5):55 (1976) | 4b:223 |
| e | <b>Austrostipa aphylla</b> (Rodway) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 584 (1996)                        | 4b:192 |
|   | <b>Austrostipa bigeniculata</b> (Hughes) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 584 (1996)                   | 4b:190 |
|   | <b>Austrostipa blackii</b> (C.E.Hubb.) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 584 (1996)                     | 4b:191 |
|   | <b>Austrostipa flavescens</b> (Labill.) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 585 (1996)                    | 4b:190 |
|   | <b>Austrostipa mollis</b> (R.Br.) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 587 (1996)                          | 4b:189 |
|   | <b>Austrostipa nodosa</b> (S.T.Blake) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 587 (1996)                      | 4b:188 |
|   | <b>Austrostipa pubinodis</b> (Trin. & Rupr.) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 588 (1996)               | 4b:187 |
|   | <b>Austrostipa rudis</b> (Sprengel) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 588 (1996)                        | 4b:186 |
|   | subsp. <b>australis</b> (J.Everett & S.W.L.Jacobs) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 588 (1996)         | 4b:186 |
|   | <b>Austrostipa scabra</b> (Lindley) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 588 (1996)                        | 4b:188 |
|   | subsp. <b>falcata</b> (Hughes) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 588 (1996)                             |        |
|   | subsp. <b>scabra</b>   |        |
|   | <b>Austrostipa semibarbata</b> (R.Br.) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 588 (1996)                     | 4b:189 |
|   | <b>Austrostipa stipoides</b> (Hook.f.) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 589 (1996)                     | 4b:186 |
|   | <b>Austrostipa stuposa</b> (Hughes) S.W.L.Jacobs & J.Everett, <i>Telopea</i> 6(4): 589 (1996)                        | 4b:191 |
| i | <b>Avellinia michelii</b> (Savi) Parl., <i>Pl. Nov.</i> 61 (1842)  |        |
|   | <i>Avena alba</i> sensu Townrow (1969) = <b>Avena barbata</b>  | 4b:236 |
| i | <b>Avena barbata</b> Pott ex Link in Schrader, <i>J. für die Bot.</i> 1799(2): 315 (1800)                            | 4b:236 |
| i | <b>Avena fatua</b> L., <i>Sp. Pl.</i> 1: 80 (1753)   | 4b:237 |
| i | <b>Avena sativa</b> L., <i>Sp. Pl.</i> 1: 79 (1753)  | 4b:237 |
| i | <b>Avena sterilis</b> L., <i>Sp. Pl.</i> ed.2 118 (1762)   | 4b:238 |
| i | subsp. <b>ludoviciana</b> (Durieu) Gillet & Magne, <i>Nouvelle Fl. Fr.</i> ed.3 532 (1873)                           | 4b:238 |
| i | <b>Avena strigosa</b> Schreber, <i>Spic. Fl. Lips.</i> 52 (1771)   | 4b:237 |
| i | <b>Bothriochloa macra</b> (Steudel) S.T.Blake, <i>Proc. R. Soc. Qld.</i> 80: 64 (1969)                               | 4b:355 |
| i | <b>Brachypodium distachyon</b> (L.) P.Beauv., <i>Ess. Agrostogr.</i> 101 (1812)                                      | 4b:292 |
| i | <b>Briza maxima</b> L., <i>Sp. Pl.</i> 1: 70 (1753)  | 4b:208 |
| i | <b>Briza minor</b> L., <i>Sp. Pl.</i> 1: 70 (1753)   | 4b:208 |
| i | <b>Bromus alopecuroides</b> Poir., <i>Voy. Barb.</i> 2: 100 (1789)   | 4b:288 |
|   | <i>Bromus arenarius</i> Labill. = <b>Bromus japonicus</b> var. <b>vestitus</b>                                       | 4b:288 |
| i | <b>Bromus brevis</b> Nees in Steudel, <i>Syn. Pl. Gram.</i> 1: 326 (1854)  | 4b:290 |
|   | <i>Bromus catharticus</i> M.Vahl. = <b>Bromus willdenowii</b>  | 4b:290 |
| i | <b>Bromus cecadilla</b> Steudel, <i>Syn. Pl. Gram.</i> 321 (1854)  | 4b:290 |
| i | <b>Bromus diandrus</b> Roth, <i>Bot. Abh. Beobacht.</i> 44 (1787)  | 4b:288 |
|   | <i>Bromus fonkii</i> Philippi = <b>Bromus lithobius</b>  | 4b:290 |
| i | <b>Bromus hordeaceus</b> L., <i>Sp. Pl.</i> 1: 77 (1753)   | 4b:287 |
| i | <b>Bromus japonicus</b> Thunb. ex Murr., <i>Syst. Veg.</i> 119 (1784)  | 4b:288 |
|   | var. <b>vestitus</b> (Schrader) Halacsy, <i>Consp. Fl. Graec.</i> 3: 397 (1904)                                      |        |
| i | <b>Bromus lithobius</b> Trin., <i>Linnaea</i> 10: 303 (1835)   | 4b:290 |
|   | <i>Bromus macrostachys</i> sensu Townrow (1969) = <b>Bromus alopecuroides</b>  | 4b:288 |
| i | <b>Bromus madritensis</b> L., <i>Cent. Pl.</i> 1: 5 (1755)   | 4b:289 |
| i | var. <b>ciliatus</b> Guss., <i>Fl. Sicul. Syn.</i> 1: 78 (1843)  | 4b:289 |
|   | <i>Bromus mollis</i> L. = <b>Bromus hordeaceus</b>   | 4b:287 |

|      |  |        |
|------|--|--------|
| i    | <b>Bromus sterilis</b> L., Sp. Pl. 1: 77 (1753)  | 4b:289 |
|      | <i>Bromus thominii</i> sensu Townrow (1969) = <b>Bromus hordeaceus</b>   | 4b:287 |
|      | <i>Bromus unioloides</i> Kunth = <b>Bromus willdenowii</b>   | 4b:290 |
| i    | <b>Bromus willdenowii</b> Kunth, Rev. Gram. 134 (1829)   | 4b:290 |
| itx# | <b>Calamagrostis epigejos</b> (L.) Roth, Tent. Fl. Germ. 1: 34 (1788)  | 4b:273 |
| i    | <b>Catapodium marinum</b> (L.) C.E.Hubb., Kew Bull. 9: 375 (1954)  | 4b:225 |
| i    | <b>Catapodium rigidum</b> (L.) C.E.Hubb. ex Dony, Fl. Bedfordshire 437 (1953)                                    | 4b:224 |
|      | <i>Ceratochloa unioloides</i> DC. = <b>Bromus willdenowii</b>  | 4b:290 |
| i    | <b>Cortaderia jubata</b> (Lemoine) Stapf in Hook.f., Bot. Mag. 124, t.7607 (1898)                                | 4b:320 |
| it   | <b>Cortaderia richardii</b> (Endl.) Zotov, N.Z. J. Bot. 1: 84 (1963)   | 4b:319 |
| i    | <b>Cortaderia selloana</b> (Schultes & Schultes.f.) Asch. & Graebner, Syn. Mitteleur. Fl. 2: 325 (1900)          | 4b:320 |
|      | <i>Critesion marinum</i> (Huds.) A.Löve = <b>Hordeum marinum</b>   | 4b:300 |
|      | <i>Critesion murinum</i> (L.) A.Löve = <b>Hordeum murinum</b>  | 4b:298 |
| i    | <b>Cynodon dactylon</b> (L.) Pers., Syn. Pl. 1: 85 (1805)  | 4b:330 |
| i    | <b>Cynosurus cristatus</b> L., Sp. Pl. 1: 72 (1753)  | 4b:207 |
| i    | <b>Cynosurus echinatus</b> L., Sp. Pl. 1: 72 (1753)  | 4b:207 |
| i    | <b>Dactylis glomerata</b> L., Sp. Pl. 1: 71 (1753)   | 4b:221 |
|      | <i>Danthonia caespitosa</i> Gaudich. = <b>Austrodanthonia caespitosa</b>   | 4b:312 |
|      | <i>Danthonia carphoides</i> F.Muell. ex Benth. var. <i>angustior</i> Vickery = <b>Austrodanthonia carphoides</b> | 4b:306 |
| i    | <b>Danthonia decumbens</b> (L.) DC. in Lam. & DC., Fl. Fr. ed.3 3: 33 (1805)                                     | 4b:317 |
|      | <i>Danthonia diemenica</i> D.I.Morris = <b>Austrodanthonia diemenica</b>   | 4b:311 |
|      | <i>Danthonia dimidiata</i> Vickery = <b>Rytidosperma dimidiatum</b>  | 4b:317 |
|      | <i>Danthonia erianthe</i> sensu Townrow (1969) = identity uncertain  |        |
|      | <i>Danthonia fortuneae-hibernae</i> Renvoize = <b>Rytidosperma fortuneae-hibernae</b>                            | 4b:310 |
|      | <i>Danthonia geniculata</i> J.Black = <b>Austrodanthonia geniculata</b>  | 4b:308 |
|      | <i>Danthonia gracilis</i> Hook.f. = <b>Notodanthonia gracilis</b>  | 4b:309 |
|      | <i>Danthonia laevis</i> Vickery = <b>Austrodanthonia laevis</b>  | 4b:312 |
|      | <i>Danthonia longifolia</i> sensu Townrow (1969) = <b>Austrodanthonia tenuior</b>                                | 4b:310 |
|      | <i>Danthonia nitens</i> D.I.Morris = <b>Rytidosperma nitens</b>  | 4b:315 |
|      | <i>Danthonia nivicola</i> Vickery = <b>Rytidosperma niviculum</b>  | 4b:315 |
|      | <i>Danthonia nudiflora</i> P.Morris = <b>Rytidosperma nudiflorum</b>   | 4b:314 |
|      | <i>Danthonia pauciflora</i> R.Br. = <b>Rytidosperma pauciflorum</b>  | 4b:309 |
|      | <i>Danthonia penicillata</i> (Labill.) R.Br. ex P.Beauv. = <b>Austrodanthonia penicillata</b>                    | 4b:313 |
|      | <i>Danthonia pilosa</i> R.Br. = <b>Austrodanthonia pilosa</b>  | 4b:313 |
|      | <i>Danthonia popinensis</i> D.I.Morris = <b>Austrodanthonia popinensis</b>                                       | 4b:316 |
|      | <i>Danthonia procera</i> Vickery = <b>Rytidosperma procerum</b>  | 4b:312 |
|      | <i>Danthonia pulvinorum</i> D.I.Morris = <b>Rytidosperma niviculum</b>   | 4b:315 |
|      | <i>Danthonia purpurascens</i> sensu Townrow (1969) = <b>Austrodanthonia tenuior</b>                              | 4b:310 |
|      | <i>Danthonia racemosa</i> R.Br. = <b>Austrodanthonia racemosa</b>  | 4b:314 |
|      | <i>Danthonia remota</i> D.I.Morris = <b>Austrodanthonia remota</b>   | 4b:316 |
|      | <i>Danthonia semiannularis</i> (Labill.) R.Br. = <b>Notodanthonia semiannularis</b>                              | 4b:308 |
|      | <i>Danthonia semiannularis</i> var. <i>gracilis</i> (Hook.f.) Hook.f. = <b>Notodanthonia gracilis</b>            | 4b:309 |
|      | <i>Danthonia setacea</i> R.Br. = <b>Austrodanthonia setacea</b>  | 4b:311 |
|      | <i>Danthonia tenuior</i> (Steud.) Conert = <b>Austrodanthonia tenuior</b>  | 4b:310 |

|    |  |        |
|----|--|--------|
|    | <b>Deschampsia cespitosa</b> (L.) P.Beauv., Ess. Agrostogr. 91: t.18 fig.3 (1812)  | 4b:242 |
| t  | <b>Deschampsia gracillima</b> Kirk, J. Bot. 29: 237 (1891)   | 4b:243 |
|    | Deyeuxia accedens Vickery = <b>Deyeuxia frigida</b>  | 4b:271 |
|    | Deyeuxia aequata (Nees) Benth. = <b>Agrostis rudis</b>   | 4b:261 |
| e  | <b>Deyeuxia apsleyensis</b> D.I.Morris, Muelleria 7(2): 160 (1990)   | 4b:273 |
|    | <b>Deyeuxia benthamiana</b> Vickery, Contr. N.S.W. Natl. Herb. 1: 63 (1940)  | 4b:271 |
|    | Deyeuxia billardierei (R.Br.) Kunth = <b>Agrostis billardierei</b>   | 4b:260 |
|    | <b>Deyeuxia brachyathera</b> (Stapf) Vickery, Contr. N.S.W. Natl. Herb. 1: 68 (1940)   | 4b:272 |
|    | <b>Deyeuxia carinata</b> Vickery, Contr. N.S.W. Natl. Herb. 1: 58 (1940)   | 4b:272 |
|    | <b>Deyeuxia contracta</b> (F.Muell. ex Hook.f.) Vickery, Contr. N.S.W. Natl. Herb. 1: 65 (1940)                                    | 4b:269 |
|    | <b>Deyeuxia decipiens</b> (R.Br.) Vickery, Contr. N.S.W. Natl. Herb. 1: 70 (1940)  | 4b:269 |
|    | <b>Deyeuxia densa</b> Benth., Fl. Austral. 7: 582 (1878)   | 4b:270 |
|    | Deyeuxia forsteri Kunth = <b>Agrostis avenacea</b>   | 4b:259 |
|    | <b>Deyeuxia frigida</b> F.Muell. ex Benth., Fl. Austral. 7: 583 (1878)   | 4b:271 |
|    | <b>Deyeuxia gunniana</b> (Nees) Benth., Fl. Austral. 7: 584 (1878)   | 4b:267 |
|    | <b>Deyeuxia innominata</b> D.I.Morris, Muelleria 7(2): 164 (1990)  | 4b:268 |
| ex | <b>Deyeuxia lawrencei</b> Vickery, Contr. N.S.W. Natl. Herb. 1: 48 (1940)  | 4b:266 |
|    | <b>Deyeuxia minor</b> F.Muell. ex Benth., Fl. Austral. 7: 582 (1878)   | 4b:270 |
|    | Deyeuxia montana Benth. = <b>Deyeuxia monticola</b>  | 4b:268 |
|    | <b>Deyeuxia monticola</b> (Roemer & Schultes) Vickery, Contr. N.S.W. Natl. Herb. 1: 56 (1940)                                      | 4b:268 |
|    | Deyeuxia parviseta sensu Townrow (1969) = <b>Deyeuxia innominata</b>   | 4b:268 |
|    | <b>Deyeuxia quadriseta</b> (Labill.) Benth., Fl. Austral. 7: 581 (1878)  | 4b:267 |
|    | <b>Deyeuxia rodwayi</b> Vickery, Contr. N.S.W. Natl. Herb. 1: 60 (1940)  | 4b:271 |
|    | <b>Deyeuxia scaberula</b> Vickery, Contr. N.S.W. Natl. Herb. 1: 64 (1940)  | 4b:270 |
|    | Deyeuxia scabra sensu Rodway (1903) = <b>Deyeuxia contracta</b> & <b>D. scaberula</b>  |        |
| i# | <b>Dichanthium sericeum</b> (R.Br.) A.Camus, Bull. Mus. Hist. Nat. Paris 27: 549 (1921)  | 4b:355 |
|    | <b>Dichelachne crinita</b> (L.f.) Hook.f., Fl. Nov.-Zel. 1: 293 (1853)   | 4b:277 |
|    | <b>Dichelachne inaequiglumis</b> (Hackel ex Cheeseman) Edgar & Connor, N.Z. J. Bot. 20: 307 (1982)                                 | 4b:278 |
|    | <b>Dichelachne micrantha</b> (Cav.) Domin, Biblioth. Bot. 20, Heft 85: 353 (1915)  | 4b:278 |
|    | <b>Dichelachne rara</b> (R.Br.) Vickery, Contr. N.S.W. Natl. Herb. 1: 337 (1951)   | 4b:279 |
|    | Dichelachne sciurea sensu Rodway (1903) = <b>Dichelachne</b> spp. excluding <i>D. crinita</i>                                      |        |
|    | <b>Dichelachne sieberiana</b> Trin. & Rupr., Mém. Acad. Imp. Sci. St.-Pétersbourg,<br>ser.6 Sci. Math., Pt.2 Sci. Nat. 5: 2 (1842) |        |
| i# | <b>Digitaria ciliaris</b> (Retz.) Koeler, Descr. Gram. 27 (1802)   | 4b:347 |
| i  | <b>Digitaria sanguinalis</b> (L.) Scop., Fl. Carniol. ed.2 1: 52 (1772)  | 4b:346 |
| i# | <b>Digitaria ternata</b> (A.Rich.) Stapf, Fl. Capensis 7: 736 (1898)   | 4b:346 |
|    | <b>Distichlis distichophylla</b> (Labill.) Fassett, Rhodora 27: 71 (1925)  | 4b:324 |
|    | Distichlis maritima sensu Rodway (1903) = <b>Distichlis distichophylla</b>   | 4b:324 |
|    | <b>Dryopoa dives</b> (F.Muell.) Vickery, Contr. N.S.W. Natl. Herb. 3(4): 196 (1963)  | 4b:199 |
| i  | <b>Echinochloa crus-galli</b> (L.) P.Beauv., Ess. Agrostogr. 53 (1812)   | 4b:336 |
|    | var. frumentacea (Link) W.F.Wright = <b>Echinochloa frumentacea</b>  | 4b:338 |
| i  | <b>Echinochloa esculenta</b> (A.Braun) H.Scholz, Taxon 41(3): 523 (1992)   | 4b:336 |
| i  | <b>Echinochloa frumentacea</b> Link, Hort. Bot. Berol. 1: 204 (1827)   | 4b:338 |
| i# | <b>Echinochloa oryzoides</b> (Ard.) Fritsch, Verh. Zool.-Bot. Ges. Wien. 41: 742 (1891)  |        |



|    |   |        |
|----|---|--------|
|    | Echinochloa utilis Ohwi & Yabuno = <b>Echinochloa esculenta</b>   | 4b:336 |
|    | <b>Echinopogon ovatus</b> (Forst.f.) P.Beauv., Ess. Agrostogr. 42, t.9 fig.5 (1812)                         | 4b:275 |
|    | <b>Ehrharta acuminata</b> (R.Br.) Sprengel, Syst. Veg. 2: 114 (1825)  | 4b:182 |
| i  | <b>Ehrharta calycina</b> Smith, Pl. Icon. Ined. 2, t.33 (1790)  | 4b:178 |
|    | <b>Ehrharta distichophylla</b> Labill., Nov. Holl. Pl. 1: 90, t.117 (1806)                                  | 4b:183 |
| i  | <b>Ehrharta erecta</b> Lam., Encycl. 2: 347 (1786)  | 4b:179 |
|    | <b>Ehrharta juncea</b> (R.Br.) Sprengel, Syst. Veg. 2: 114 (1825)   | 4b:183 |
| i  | <b>Ehrharta longiflora</b> Smith, Pl. Icon. Ined. 2, t.32 (1790)  | 4b:179 |
| e  | <b>Ehrharta oreophila</b> (D.I.Morris) Willemse, Blumea 28: 193 (1982)                                      | 4b:182 |
| e  | var. <b>minor</b> (D.I.Morris) Willemse, Blumea 28: 193 (1982)  | 4b:182 |
| e  | var. <b>oreophila</b>   | 4b:182 |
|    | <b>Ehrharta stipoides</b> Labill., Nov. Holl. Pl. 1: 91, t.118 (1805)                                       | 4b:181 |
| e  | <b>Ehrharta tasmanica</b> (Hook.f.) Willemse, Blumea 28: 193 (1982)   | 4b:181 |
| e  | var. <b>subalpina</b> (F.Muell. ex Benth.) Willemse, Blumea 28: 193 (1982)                                  | 4b:182 |
| e  | var. <b>tasmanica</b>   | 4b:181 |
| i  | <b>Ehrharta villosa</b> (L.f.) Schultes f. ex Schultes & Schultes f., Syst. Veg. 7: 1374 (1829)             | 4b:184 |
| i# | <b>Eleusine indica</b> (L.) Gaertn., Fruct. Sem. Pl. 1: 8 (1788)  |        |
|    | Elymus elongatus (Host) Runemark = <b>Lophopyrum ponticum</b>   | 4b:295 |
|    | Elymus farctus subsp. boreali-atlanticus (Simonet & Guinochet) Melderis = <b>Thinopyrum junceiforme</b>     |        |
|    | Elymus repens (L.) Gould = <b>Elytrigia repens</b>  | 4b:295 |
|    | <b>Elymus scaber</b> (R.Br.) A.Löve, Feddes Repert. 95: 468 (1984)  | 4b:296 |
| i  | <b>Elytrigia repens</b> (L.) Nevski, Trudy Bot. Inst. Akad. Nauk SSSR, ser.1, Fl.Sist.Vyss.Rast.1:14 (1933) | 4b:295 |
|    | <b>Eragrostis brownii</b> (Kunth) Nees ex Steudel, Nom. Bot. 1: 562 (1840)                                  | 4b:326 |
| i  | <b>Eragrostis cilianensis</b> (All.) Link ex Vign., Malpighia 18: 386 (1904)                                | 4b:326 |
| i# | <b>Eragrostis curvula</b> (Schrader) Nees, Fl. Afr. Austral. 1: 397 (1841)                                  | 4b:327 |
|    | Eragrostis molybdea Vickery = <b>Eragrostis brownii</b>   | 4b:326 |
| i# | <b>Eragrostis tenuifolia</b> (A.Rich.) Hochst. ex Steudel, Syn. Pl. Glum. 1: 268 (1854)                     | 4b:327 |
|    | Erythranthera australis (Petrie) Zotov = <b>Rytidosperma australe</b>                                       | 4b:318 |
| ex | <b>Festuca archeri</b> Alexeev, Nov. Syst. Pl. Vasc. 24: 8 (1987)   | 4b:428 |
| i  | <b>Festuca arundinacea</b> Schreber, Spic. Fl. Lips. 57 (1771)  | 4b:197 |
|    | Festuca asperula sensu Townrow (1969) = <b>Festuca plebeia</b>  | 4b:196 |
|    | Festuca bromoides L. = <b>Vulpia bromoides</b>  | 4b:203 |
|    | Festuca dives F.Muell. = <b>Dryopoa dives</b>   | 4b:199 |
|    | Festuca elatior L. subsp. arundinacea (Schreber) Hackel = <b>Festuca arundinacea</b>                        | 4b:197 |
|    | Festuca hookeriana F.Muell. ex Hook.f. = <b>Austrofestuca hookeriana</b>                                    | 4b:223 |
|    | Festuca littoralis Labill. = <b>Austrofestuca littoralis</b>  | 4b:223 |
| i  | <b>Festuca nigrescens</b> Lam., Encycl. Méth. Bot. 2: 460 (1788)  | 4b:197 |
|    | Festuca ovina sensu Rodway (1903) = <b>Festuca plebeia</b>  | 4b:196 |
| e  | <b>Festuca plebeia</b> R.Br., Prodr. 178 (1810)   | 4b:196 |
| i  | <b>Festuca rubra</b> L., Sp. Pl. 1: 74 (1753)   | 4b:197 |
|    | Gastridium lendigerum sensu Rodway (1903) = <b>Gastridium ventricosum</b>                                   | 4b:280 |
| it | <b>Gastridium ventricosum</b> (Gouan) Schinz & Thell., Viert. Naturf. Ges. Zurich 58: 39 (1913)             | 4b:280 |
| i  | <b>Gaudinia fragilis</b> (L.) P.Beauv., Ess. Agrostogr. 95 (1812)   | 4b:238 |
|    | <b>Glyceria australis</b> C.E.Hubb., Bull. Misc. Inform. 10: 450 (1934)                                     | 4b:228 |

|    |  |        |
|----|--|--------|
| i  | <b>Glyceria declinata</b> Bréb., Fl. Normandie ed.3: 354 (1859)                                | 4b:229 |
| i  | <b>Glyceria fluitans</b> (L.) R.Br., Prodr. 179 (1810)   | 4b:229 |
|    | <i>Glyceria fluitans</i> sensu Rodway (1903) = <b>Glyceria australis</b>                       | 4b:228 |
| i  | <b>Glyceria maxima</b> (Hartm.) Holmb., Bot. Not. 1919: 97 (1919)                              | 4b:229 |
| i# | <b>Glyceria notata</b> Chevall, Fl. Gen. Env. Paris 2: 174 (1827)                              | 4b:229 |
|    | <i>Glyceria plicata</i> (Fries) Fries = <b>Glyceria notata</b>                                 | 4b:229 |
|    | <i>Glyceria stricta</i> Hook.f. = <b>Puccinellia stricta</b>                                   | 4b:205 |
| i  | <b>Hainardia cylindrica</b> (Willd.) Greuter, Boissiera 13: 177 (1967)                         | 4b:227 |
|    | <b>Hemarthria uncinata</b> R.Br., Prodr. 207 (1810)  | 4b:359 |
| e  | <b>Hierochloe fraseri</b> Hook.f., Fl. Antarct. 1: 93 (1845)                                   | 4b:248 |
|    | <b>Hierochloe rariflora</b> Hook.f., Fl. Antarct. 1: 93 (1845)                                 | 4b:249 |
|    | <b>Hierochloe redolens</b> (Vahl) Roemer & Schultes, Syst. Veg. 2: 514 (1817)                  | 4b:247 |
| i  | <b>Holcus lanatus</b> L., Sp. Pl. 2: 1048 (1753)   | 4b:245 |
| i  | <b>Holcus mollis</b> L., Syst. Nat. ed.10 2: 1305 (1759)                                       | 4b:245 |
| i  | <b>Hordeum distichon</b> L., Sp. Pl. 1: 85 (1753)  | 4b:297 |
|    | <i>Hordeum hystrix</i> sensu Townrow (1969) = <b>Hordeum marinum</b> subsp. <b>gussonianum</b> | 4b:300 |
|    | <i>Hordeum leporinum</i> Link. = <b>Hordeum murinum</b> subsp. <b>leporinum</b>                | 4b:298 |
| i  | <b>Hordeum marinum</b> Huds., Fl. Angl. ed.2 1: 57 (1778)                                      | 4b:300 |
| i  | subsp. <b>gussoneanum</b> (Parl.) Thell., Viert. Natl. Ges. Zurich 52: 441 (1908)              | 4b:300 |
| i  | subsp. <b>marinum</b>  | 4b:300 |
| i  | <b>Hordeum murinum</b> L., Sp. Pl. 1: 85 (1753)  | 4b:298 |
| i  | subsp. <b>glaucum</b> (Steudel) Tzvelev, Novit. Syst. Pl. Vasc. (Leningrad) 8: 67 (1971)       | 4b:298 |
| i  | subsp. <b>leporinum</b> (Link) Arcang., Comp. Fl. Ital. 805 (1882)                             | 4b:298 |
| i  | subsp. <b>murinum</b>  | 4b:298 |
| i  | <b>Hordeum vulgare</b> L., Sp. Pl. 1: 84 (1753)  | 4b:297 |
|    | <i>Imperata arundinacea</i> sensu Rodway (1903) = <b>Imperata cylindrica</b>                   | 4b:353 |
|    | <b>Imperata cylindrica</b> (L.) Raeuschel, Nomencl. Bot. ed.3, 10 (1797)                       | 4b:353 |
|    | var. <b>major</b> (Nees) C.E.Hubb., Grasses of Mauritius & Rodriguez 96 (1940)                 | 4b:353 |
|    | <b>Isachne globosa</b> (Thunb.) Kuntze, Rev. Gen. Pl. 2: 778 (1891)                            |        |
|    | <i>Koeleria cristata</i> Pers. = <b>Rostraria cristata</b>                                     | 4b:241 |
|    | <i>Koeleria cristata</i> sensu Rodway (1903) = <b>Koeleria macrantha</b>                       | 4b:240 |
| ix | <b>Koeleria macrantha</b> (Ledeb.) Schult., Mantissa 2: 345 (1824)                             | 4b:240 |
|    | <i>Koeleria phleoides</i> sensu Rodway (1903) = <b>Rostraria cristata</b>                      | 4b:241 |
|    | <b>Lachnagrostis aemula</b> (R.Br.) Trin., Fund. Agrost. 128 (1820)                            | 4b:258 |
|    | <b>Lachnagrostis aequata</b> (Nees) S.W.L.Jacobs, Telopea 9(3): 445 (2001)                     | 4b:262 |
|    | <b>Lachnagrostis billardierei</b> (R.Br.) Trin., Fund. Agrost. 128 (1820)                      | 4b:260 |
|    | subsp. <b>billardierei</b>   | 4b:260 |
| e  | subsp. <b>tenuiseta</b> (D.I.Morris) S.W.L.Jacobs, Telopea 9(3): 445 (2001)                    | 4b:261 |
| e  | <b>Lachnagrostis collicola</b> (D.I.Morris) S.W.L.Jacobs, Telopea 9(3): 445 (2001)             | 4b:261 |
|    | <b>Lachnagrostis filiformis</b> (Forst.) Trin., Fund. Agrost. 128 (1820)                       | 4b:259 |
| e  | <b>Lachnagrostis lacunarum</b> (D.I.Morris) S.W.L.Jacobs, Telopea 9(3): 446 (2001)             | 4b:260 |
|    | <b>Lachnagrostis punicea</b> (A.J.Brown & N.G.Walsh) S.W.L.Jacobs, Telopea 9(4): 837 (2002)    | 4b:259 |
|    | subsp. <b>filifolia</b> (Vickery) S.W.L.Jacobs, Telopea 9(3): 446 (2001)                       | 4b:261 |
|    | subsp. <b>punicea</b>  | 4b:259 |

|    |  |        |
|----|--|--------|
|    | <b>Lachnagrostis robusta</b> (Vickery) S.W.L.Jacobs, Telopea 9(3): 447 (2001)                        | 4b:261 |
|    | Lachnagrostis rudis (Roem. & Schult.) Trin. = <b>Lachnagrostis scabra</b>                            | 4b:261 |
|    | <b>Lachnagrostis scabra</b> (Beauv.) Nees ex Steudel, Nomen. Bot. edn.2, 1: 250 (1840)               | 4b:261 |
| i  | <b>Lagurus ovatus</b> L., Sp. Pl. 1: 81 (1753)   | 4b:281 |
|    | Lepturus cylindricus (Willd.) Trin. = <b>Hainardia cylindrica</b>                                    | 4b:227 |
|    | Lepturus incurvatus Trin. = <b>Parapholis incurva</b>  | 4b:226 |
| it | <b>Leymus arenarius</b> (L.) Hochst., Flora (Regensb.) 31: 118 (1848)                                |        |
| i  | <b>Lolium loliaceum</b> (Bory & Chaub.) Hand.-Mazz., Ann. Naturh. Mus. Hoffmus. Wien 28: 32 (1914)   | 4b:202 |
| i  | <b>Lolium multiflorum</b> Lam., Fl. Franc. 3: 621 (1779)   | 4b:201 |
| i  | <b>Lolium perenne</b> L., Sp. Pl. 1: 83 (1753)   | 4b:201 |
|    | Lolium perenne var. italicum (A.Braun) Rodway = <b>Lolium multiflorum</b>                            |        |
| i  | <b>Lolium rigidum</b> Gaudin, Agrost. Helv. 1: 334 (1811)  | 4b:201 |
| i  | <b>Lolium temulentum</b> L., Sp. Pl. 1: 83 (1753)  | 4b:200 |
| i  | forma <b>arvense</b> (With.) Junge, Jahrb. Hamb. Wiss. Anst. Beih. 3, 30: 314 (1913)                 | 4b:200 |
|    | Lophochloa cristata (L.) Hylander = <b>Rostraria cristata</b>  | 4b:241 |
|    | Lophopyrum elongatum sensu Census (1999) = <b>Lophopyrum ponticum</b>                                | 4b:295 |
| i  | <b>Lophopyrum ponticum</b> (Podp.) A.Löve, Feddes Repert. 95(7-8): 489 (1984)                        | 4b:295 |
|    | Microlaena stipoides (Labill.) R.Br. = <b>Ehrharta stipoides</b>                                     | 4b:181 |
|    | Microlaena tasmanica (Hook.f.) Hook.f. ex Benth. = <b>Ehrharta tasmanica</b>                         | 4b:181 |
|    | var. subalpina F.Muell. ex Benth. = <b>Ehrharta tasmanica</b> var. <b>subalpina</b>                  | 4b:182 |
|    | var. tasmanica = <b>Ehrharta tasmanica</b> var. <b>tasmanica</b>                                     | 4b:181 |
| i# | <b>Molineriella minuta</b> (L.) Rouy, Fl. France 14: 102 (1913)                                      |        |
|    | Monerma cylindrica (Willd.) Cosson & Durieu = <b>Hainardia cylindrica</b>                            | 4b:227 |
| it | <b>Nardus stricta</b> L., Sp. Pl. 1: 53 (1753)   | 4b:184 |
| i  | <b>Nassella neesiana</b> (Trin. & Rupr.) Barkworth, Taxon 39: 611 (1990)                             |        |
| i  | <b>Nassella trichotoma</b> (Nees) Hackel ex Arechav., Anales Mus. Nac. Montevideo 1: 366 t.19 (1896) | 4b:193 |
|    | Notodanthonia caespitosa (Gaudich.) Zotov = <b>Austrodanthonia caespitosa</b>                        | 4b:312 |
|    | Notodanthonia carphoides (F.Muell. ex Benth.) Zotov = <b>Austrodanthonia carphoides</b>              | 4b:306 |
|    | Notodanthonia diemenica (D.I.Morris) H.P.Linder = <b>Austrodanthonia diemenica</b>                   | 4b:311 |
|    | Notodanthonia geniculata (J.Black) Zotov = <b>Austrodanthonia geniculata</b>                         | 4b:308 |
| t  | <b>Notodanthonia gracilis</b> (Hook.f.) Zotov, N.Z. J. Bot 1(1): 123 (1963)                          | 4b:309 |
|    | Notodanthonia laevis (Vickery) Zotov = <b>Austrodanthonia laevis</b>                                 | 4b:312 |
|    | Notodanthonia penicillata (Labill.) Zotov = <b>Austrodanthonia penicillata</b>                       | 4b:313 |
|    | Notodanthonia pilosa (R.Br.) Zotov = <b>Austrodanthonia pilosa</b>                                   | 4b:313 |
|    | Notodanthonia popinensis (D.I.Morris) H.P.Linder = <b>Austrodanthonia popinensis</b>                 | 4b:316 |
|    | Notodanthonia racemosa (R.Br.) Zotov = <b>Austrodanthonia racemosa</b>                               | 4b:314 |
|    | Notodanthonia remota (D.I.Morris) H.P.Linder = <b>Austrodanthonia remota</b>                         | 4b:316 |
|    | <b>Notodanthonia semiannularis</b> (Labill.) Zotov, N.Z. J. Bot 1(1): 116 (1963)                     | 4b:308 |
|    | Notodanthonia setacea (R.Br.) Veldk. = <b>Austrodanthonia setacea</b>                                | 4b:311 |
|    | Notodanthonia tenuior (Steud.) Conert = <b>Austrodanthonia tenuior</b>                               | 4b:310 |
|    | Oryzopsis miliacea (L.) Benth. & Hook.f. ex Asch. & Schweinf. = <b>Piptatherum miliaceum</b>         | 4b:194 |
| i# | <b>Panicum capillare</b> L., Sp. Pl. 1: 58 (1753)  | 4b:335 |
| i# | var. <b>capillare</b>  | 4b:335 |
| i# | var. <b>occidentale</b> Rydb., Contr. U.S. Natl. Herb. 3: 186 (1895)                                 | 4b:335 |

|    |  |        |
|----|--|--------|
| i# | <b>Panicum gilvum</b> Launert, Mit. Bot. Staatssamml. Munchen 8: 153 (1970)                            | 4b:335 |
| i  | <b>Panicum hillmanii</b> Chase, J. Wash. Acad. Sci. 14: 345, fig.1. (1934)                             | 4b:334 |
| i  | <b>Panicum miliaceum</b> L., Sp. Pl. 1: 58 (1753)  | 4b:333 |
|    | <b>Panicum sanguinale</b> L. = <b>Digitaria sanguinalis</b>  | 4b:346 |
|    | <b>Panicum schinzii</b> sensu Curtis & Morris (1995) = <b>Panicum gilvum</b>                           | 4b:335 |
| i  | <b>Parapholis incurva</b> (L.) C.E.Hubb., Blumea suppl. 3: 14 (1946)                                   | 4b:226 |
| i  | <b>Parapholis strigosa</b> (Dumort.) C.E.Hubb., Blumea suppl. 3: 14 (1946)                             | 4b:226 |
|    | <b>Paspalidium gracile</b> sensu Townrow (1969) = identity uncertain                                   |        |
| i  | <b>Paspalum dilatatum</b> Poiret in Lam., Encycl. 5: 35 (1804)   | 4b:340 |
| i  | <b>Paspalum distichum</b> L., Syst. Nat. ed.10, 2: 855 (1759)  | 4b:340 |
| i  | <b>Paspalum urvillei</b> Steudel, Syn. Pl. Glum. 1: 24 (1853)  | 4b:340 |
| i# | <b>Pennisetum alopecuroides</b> (L.) Sprengel, one record from Launceston but probably not naturalised |        |
| i  | <b>Pennisetum clandestinum</b> Hochst. ex Chiov., Ann. Inst. Bot. Roma 8: 41 (1903)                    | 4b:348 |
| i  | <b>Pennisetum macrourum</b> Trin., Gram. Pan. 64 (1826)  | 4b:351 |
| i  | <b>Pennisetum villosum</b> R.Br. ex Fresen., Mus. Senckenb. Abh. 2: 134 (1837)                         | 4b:349 |
|    | <b>Pentapogon billardierei</b> sensu Rodway (1903) = <b>Pentapogon quadrifidus</b>                     | 4b:275 |
|    | <b>Pentapogon quadrifidus</b> (Labill.) Baill., Hist. Pl. 12: 280 (1893)                               | 4b:275 |
| e  | var. <b>parviflorus</b> (Benth.) D.I.Morris, Muelleria 7(2): 167 (1990)                                | 4b:276 |
|    | var. <b>quadrifidus</b>  | 4b:275 |
| i  | <b>Phalaris aquatica</b> L., Cent. Pl. 1: 4 (1755)   | 4b:252 |
| i  | <b>Phalaris arundinacea</b> L., Sp. Pl. 1: 55 (1753)   | 4b:251 |
| i  | <b>Phalaris canariensis</b> L., Sp. Pl. 1: 54 (1753)   | 4b:251 |
| i  | <b>Phalaris minor</b> Retz., Obs. Bot. 3: 8 (1783)   | 4b:252 |
| i  | <b>Phalaris paradoxa</b> L., Sp. Pl. ed.2, 2: 1665 (1763)  | 4b:251 |
|    | <b>Phalaris tuberosa</b> L. = <b>Phalaris aquatica</b>   | 4b:252 |
| i  | <b>Phleum pratense</b> L., Sp. Pl. 1: 59 (1753)  | 4b:286 |
|    | <b>Phragmites australis</b> (Cav.) Trin. ex Steudel, Nomencl. Bot. ed.2 2: 324 (1841)                  | 4b:322 |
|    | <b>Phragmites communis</b> sensu Rodway (1903) = <b>Phragmites australis</b>                           | 4b:322 |
| i  | <b>Piptatherum miliaceum</b> (L.) Cosson, Notes Pl. Crit. ser.2, fasc.3: 129 (1851)                    | 4b:194 |
| i  | <b>Poa annua</b> L., Sp. Pl. 1: 68 (1753)  | 4b:211 |
|    | <b>Poa billardierei</b> sensu Rodway (1903) = <b>Poa poiformis</b>                                     | 4b:220 |
| i  | <b>Poa bulbosa</b> L., Sp. Pl. 1: 70 (1753)  | 4b:213 |
|    | <b>Poa caespitosa</b> sensu Rodway (1903) = <b>P. gunnii</b> , <b>Poa rodwayi</b> & <b>P. tenera</b>   |        |
|    | var. <b>alpina</b> F.Muell. ex Benth. = <b>Poa gunnii</b>  | 4b:218 |
|    | var. <b>australis</b> (R.Br.) Benth. = <b>Poa rodwayi</b>  | 4b:215 |
|    | var. <b>tenera</b> (F.Muell.) Benth. = <b>Poa tenera</b>   | 4b:217 |
|    | <b>Poa clelandii</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 193 (1970)                               | 4b:221 |
|    | <b>Poa clivicola</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 213 (1970)                               |        |
| i  | <b>Poa compressa</b> L., Sp. Pl. 1: 69 (1753)  | 4b:214 |
|    | <b>Poa costiniana</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 214 (1970)                              | 4b:218 |
|    | <b>Poa fawcettiae</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 232 (1970)                              | 4b:217 |
|    | <b>Poa gunnii</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 217 (1970)                                  | 4b:218 |
|    | <b>Poa halmaturina</b> J.Black, Trans. R. Soc. S.Austral. 66: 248 (1943)                               | 4b:220 |
|    | <b>Poa hiemata</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 230 (1970)                                 | 4b:217 |

|   |   |        |
|---|---|--------|
|   | <b>Poa hookeri</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 222 (1970)                                  | 4b:216 |
| i | <b>Poa infirma</b> Kunth in Humboldt, Bonpland & Kunth, Nov. Gen. Sp. 1: 158 (1816)                     | 4b:213 |
| e | <b>Poa jugicola</b> D.I.Morris, Muellera 7(2): 167 (1990)   | 4b:219 |
|   | <b>Poa labillardierei</b> Steudel, Syn. Pl. Glumac. 1: 262 (1854)                                       | 4b:219 |
|   | var. <b>acris</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 205 (1970)                                   | 4b:220 |
|   | var. <b>labillardierei</b>  | 4b:219 |
| e | <b>Poa mollis</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 241 (1970)                                   | 4b:215 |
|   | <b>Poa poiformis</b> (Labill.) Druce, Bot. Soc. Exch. Club Brit. Isles 1916: 640 (1917)                 | 4b:220 |
|   | var. <b>poiformis</b>   | 4b:220 |
|   | var. <b>ramifer</b> D.I.Morris, Muellera 7(2): 169 (1990)   | 4b:220 |
| i | <b>Poa pratensis</b> L., Sp. Pl. 1: 67 (1753)   | 4b:214 |
|   | <i>Poa rigida</i> L. = <b>Catapodium rigidum</b>  | 4b:224 |
|   | <b>Poa rodwayi</b> Vickery, Contr. N.S.W. Natl. Herb. 4(4): 235 (1970)                                  | 4b:215 |
|   | <b>Poa saxicola</b> R.Br., Prodr. 180 (1810)  | 4b:214 |
|   | <b>Poa sieberiana</b> Sprengel, Syst. Veg. Cur. Post. 35 (1827)   | 4b:216 |
|   | <b>Poa tenera</b> F.Muell. ex Hook.f., Fl. Tasm. 2: 124, t.164a (1858)                                  | 4b:217 |
| i | <b>Poa trivialis</b> L., Sp. Pl. 1: 67 (1753)   | 4b:213 |
|   | <i>Polypogon littoralis</i> Smith = <b>XAgropogon littoralis</b>  | 4b:283 |
| i | <b>Polypogon maritimus</b> Willd., Ges. Naturf. Freunde Berlin Neue Schriften 3: 442 (1801)             | 4b:282 |
| i | <b>Polypogon monspeliensis</b> (L.) Desf., Fl. Atlant. 1: 67 (1798)                                     | 4b:282 |
| i | <b>Polypogon viridis</b> (Gouan) Breistr., Bull. Soc. Bot. Fr. 110: 56 (1966)                           | 4b:283 |
|   | <i>Psamma arenaria</i> (L.) P.Beauv. = <b>Ammophila arenaria</b>  | 4b:274 |
| i | <b>Puccinellia distans</b> (L.) Parl., Fl. Ital. 1: 366 (1848)  | 4b:205 |
|   | <b>Puccinellia stricta</b> (Hook.f.) Blom, Acta Hort. Gothob. 5: 89 (1930)                              | 4b:205 |
|   | var. <b>perlaxa</b> Stapf ex N.G.Walsh, Muellera 7: 382 (1991)  | 4b:205 |
|   | var. <b>stricta</b>   | 4b:205 |
| i | <b>Rostraria cristata</b> (L.) Tzvelev, Novosti Sist. Vyssh. Rast. 7: 47 (1971)                         | 4b:241 |
|   | <b>Rytidosperma australe</b> (Petrie) Clayton & Renvoize ex Connor & Edgar, N.Z. J. Bot. 25: 166 (1987) | 4b:318 |
|   | <b>Rytidosperma dimidiatum</b> (Vickery) Connor & Edgar, N.Z. J. Bot. 17: 332 (1979)                    | 4b:317 |
| e | <b>Rytidosperma fortuneae-hibernae</b> (Renvoize) Connor & Edgar, N.Z. J. Bot. 17: 332 (1979)           | 4b:310 |
| e | <b>Rytidosperma nitens</b> (D.I.Morris) H.P.Linder, Telopea 6(4): 614 (1996)                            | 4b:315 |
|   | <b>Rytidosperma niviculum</b> (Vickery) Connor & Edgar, N.Z. J. Bot. 17: 332 (1979)                     | 4b:315 |
|   | <b>Rytidosperma nudiflorum</b> (P.Morris) Connor & Edgar, N.Z. J. Bot. 17: 332 (1979)                   | 4b:314 |
| e | <b>Rytidosperma pauciflorum</b> (R.Br.) Connor & Edgar, N.Z. J. Bot. 17: 332 (1979)                     | 4b:309 |
|   | <i>Rytidosperma procerum</i> (Vickery) Connor & Edgar = <b>Austrodanthonia procera</b>                  | 4b:312 |
|   | <i>Schedonorus hookerianus</i> (F.Muell. ex Hook.f.) P.Beauv. = <b>Austrofestuca hookeriana</b>         | 4b:223 |
|   | <i>Schedonorus littoralis</i> (Labill.) P.Beauv. = <b>Austrofestuca littoralis</b>                      | 4b:223 |
| i | <b>Sclerochloa dura</b> (L.) P.Beauv., Ess. Agrostogr. 98, t.19 fig.4 (1812)                            | 4b:225 |
| i | <b>Secale cereale</b> L., Sp. Pl. 1: 85 (1753)  | 4b:302 |
|   | <i>Setaria geniculata</i> (Lam.) P.Beauv. = <b>Setaria gracilis</b>                                     | 4b:344 |
|   | var. <b>pauciseta</b> Desv. = <b>Setaria gracilis</b> var. <b>pauciseta</b>                             | 4b:344 |
| i | <b>Setaria gracilis</b> Kunth in Humboldt, Bonpland & Kunth, Nov. Gen. Sp. 1: 109 (1816)                | 4b:344 |
| i | var. <b>pauciseta</b> (Desv.) B.Simon, Austrobaileya 2(1): 22 (1984)                                    | 4b:344 |
| i | <b>Setaria italica</b> (L.) P.Beauv., Ess. Agrostogr. 51, 170, 178 (1812)                               | 4b:343 |

|   |  |        |
|---|--|--------|
| i | <b>Setaria verticillata</b> (L.) P.Beauv., Ess. Agrostogr. 51, 171, 178 (1812)                               | 4b:342 |
| i | <b>Setaria viridis</b> (L.) P.Beauv., Ess. Agrostogr. 51, 171, 178 (1812)                                    | 4b:343 |
|   | Sieglingia decumbens (L.) Bernh. = <b>Danthonia decumbens</b>  | 4b:317 |
| i | <b>Spartina anglica</b> C.E.Hubb. in Heywood, Bot. J. Linn. Soc. 76: 364 (1978)                              | 4b:330 |
|   | Spartina townsendii sensu Townrow (1969) = <b>Spartina anglica</b>   | 4b:330 |
|   | Spinifex hirsutus sensu Rodway (1903) = <b>Spinifex sericeus</b>   | 4b:351 |
|   | <b>Spinifex sericeus</b> R.Br., Prodr. 198 (1810)  | 4b:351 |
| i | <b>Sporobolus africanus</b> (Poirlet) Robyns & Tournay, Bull. Jard. Bot. Brux. 25: 242 (1955)                | 4b:328 |
|   | Sporobolus capensis Kunth = <b>Sporobolus africanus</b>  | 4b:328 |
|   | Sporobolus indicus (L.) R.Br. = <b>Sporobolus africanus</b>  | 4b:328 |
|   | <b>Sporobolus virginicus</b> (L.) Kunth, Revis. Gram. 67 (1829)  | 4b:328 |
|   | Stenotaphrum americanum Schrank = <b>Stenotaphrum secundatum</b>   | 4b:344 |
| i | <b>Stenotaphrum secundatum</b> (Walter) Kuntze, Revis. Gen. Pl. 2: 794 (1891)                                | 4b:344 |
|   | Stipa aphanoneura Hughes = <b>Austrostipa flavescens</b>   | 4b:190 |
|   | Stipa aphylla (Rodway) J.Townrow = <b>Austrostipa aphylla</b>  | 4b:192 |
|   | Stipa bigeniculata Hughes = <b>Austrostipa bigeniculata</b>  | 4b:190 |
|   | Stipa blackii C.E.Hubb. = <b>Austrostipa blackii</b>   | 4b:191 |
|   | Stipa caudata Trin. = <b>Achnatherum caudatum</b> 4b:192   |        |
|   | Stipa compacta Hughes = <b>Austrostipa flavescens</b>  | 4b:190 |
|   | Stipa elatior (Benth.) Hughes = <b>Austrostipa flavescens</b>  | 4b:190 |
|   | Stipa eremophila sensu Townrow (1969) = identity uncertain   |        |
|   | Stipa falcata sensu Townrow (1978) = <b>Austrostipa scabra</b> p.p.maj.                                      | 4b:188 |
|   | Stipa flavescens Labill. = <b>Austrostipa flavescens</b>   | 4b:190 |
|   | Stipa mollis R.Br. = <b>Austrostipa mollis</b>   | 4b:189 |
|   | Stipa neesiana Trin. & Rupr. = <b>Nassella neesiana</b>  |        |
|   | Stipa nervosa Vickery var. neutralis sensu Townrow (1978) = <b>Austrostipa rudis</b> subsp. <b>australis</b> | 4b:186 |
|   | Stipa nodosa S.T.Blake = <b>Austrostipa nodosa</b>   | 4b:188 |
|   | Stipa pubescens sensu Rodway (1903) = <b>Austrostipa pubinodis</b>   | 4b:187 |
|   | Stipa pubescens var. aphylla Rodway = <b>Austrostipa aphylla</b>   | 4b:192 |
|   | Stipa pubinodis Trin. & Rupr. = <b>Austrostipa pubinodis</b>   | 4b:187 |
|   | Stipa rudis Sprengel = <b>Austrostipa rudis</b>  | 4b:186 |
|   | Stipa scabra Lindley = <b>Austrostipa scabra</b>   | 4b:188 |
|   | Stipa semibarbata R.Br. = <b>Austrostipa semibarbata</b>   | 4b:189 |
|   | Stipa setacea sensu Rodway (1903) = <b>Austrostipa stiposa</b>   | 4b:191 |
|   | Stipa stipoides (Hook.f.) Veldkamp = <b>Austrostipa stipoides</b>  | 4b:186 |
|   | Stipa stiposa Hughes = <b>Austrostipa stiposa</b>  | 4b:191 |
|   | Stipa teretifolia Steudel = <b>Austrostipa stipoides</b>   | 4b:186 |
|   | Stipa variabilis sensu Townrow (1969) = <b>Austrostipa nodosa</b>  | 4b:188 |
|   | Tetrarrhena acuminata R.Br. = <b>Ehrharta acuminata</b>  | 4b:182 |
|   | Tetrarrhena distichophylla (Labill.) R.Br. = <b>Ehrharta distichophylla</b>                                  | 4b:183 |
|   | Tetrarrhena juncea R.Br. = <b>Ehrharta juncea</b>  | 4b:183 |
|   | Tetrarrhena oreophila D.I.Morris = <b>Ehrharta oreophila</b>   | 4b:182 |
|   | var. minor D.I.Morris = <b>Ehrharta oreophila</b> var. <b>minor</b>  | 4b:182 |
|   | var. oreophila = <b>Ehrharta oreophila</b> var. <b>oreophila</b>   | 4b:182 |

|    |  |        |
|----|--|--------|
|    | <i>Themeda australis</i> (R.Br.) Stapf = <b>Themeda triandra</b>                             | 4b:358 |
|    | <b>Themeda triandra</b> Forsskal, Fl. Aegypt.-Arab. 178 (1775)                               | 4b:358 |
| i  | <b>Thinopyrum junceiforme</b> (A.& D.Löve) A.Löve, Taxon 29: 351 (1980)                      | 4b:293 |
|    | <i>Thonandia gracilis</i> (Hook.f.) H.P.Linder = <b>Notodanthonia gracilis</b>               | 4b:309 |
|    | <i>Thonandia semiannularis</i> (Labill.) H.P.Linder = <b>Notodanthonia semiannularis</b>     | 4b:308 |
|    | <b>Trisetum spicatum</b> (L.) Richter, Pl. Eur. 1: 59 (1890)                                 | 4b:239 |
|    | subsp. <b>australiense</b> Hultén, Svensk. Bot. Tidskr. 53: 220 (1959)                       | 4b:239 |
|    | <i>Trisetum subspicatum</i> sensu Rodway (1903) = <b>Trisetum spicatum</b>                   | 4b:239 |
| i  | <b>Triticum aestivum</b> L., Sp. Pl. 1: 85 (1753)  | 4b:303 |
| i  | <b>Vulpia bromoides</b> (L.) Gray, Nat. Arr. Brit. Pl. 2: 124 (1821)                         | 4b:203 |
| i# | <b>Vulpia fasciculata</b> (Forsskal) Samp., Lista Esp. Herb. Port. 24 (1913)                 | 4b:204 |
|    | <i>Vulpia megalura</i> (Nutt.) Rydb. = <b>Vulpia myuros</b> forma <b>megalura</b>            | 4b:204 |
| i  | <b>Vulpia muralis</b> (Kunth) Nees, Linnaea 19: 694 (1847)                                   |        |
| i  | <b>Vulpia myuros</b> (L.) C.Gmelin, Fl. Bad. 1: 9 (1805)                                     | 4b:203 |
| i  | forma <b>megaleura</b> (Nutt.) Stace & R.Cotton, Watsonia 11: 72 (1976)                      | 4b:204 |
| i  | forma <b>myuros</b>  |        |
|    | <b>Zoysia</b> = <b>Zoysia</b>  |        |
|    | <b>Zoysia macrantha</b> Desv., Opusc. Sci. Phys. Nat. 54 (1831)                              | 4b:332 |
|    | <i>Zoysia pungens</i> sensu Rodway (1903) = <b>Zoysia macrantha</b>                          | 4b:332 |
|    | <b>POSIDONIACEAE</b>   |        |
|    | <b>Posidonia australis</b> Hook.f., Fl. Tasm. 2: 43 (1858)                                   | 4b:25  |
|    | <b>POTAMOGETONACEAE</b>  |        |
|    | <b>Potamogeton australiensis</b> A.Benn., J. Bot. Lond. 48: 149 (1910)                       | 4b:19  |
|    | <i>Potamogeton cheesemanii</i> sensu J.H.Willis (1973) = <b>Potamogeton tricarinatus</b>     | 4b:19  |
|    | <b>Potamogeton crispus</b> L., Sp. Pl. 1: 126 (1753)   | 4b:18  |
|    | <i>Potamogeton natans</i> sensu Rodway (1903) = <b>Potamogeton tricarinatus</b>              | 4b:19  |
|    | <i>Potamogeton obtusifolius</i> sensu Rodway (1903) = <b>Potamogeton crispus</b>             | 4b:18  |
|    | <b>Potamogeton ochreatus</b> Raoul, Ann. Sci. Nat. Bot. ser.3, 2: 117 (1844)                 | 4b:18  |
|    | <b>Potamogeton pectinatus</b> L., Sp. Pl. 1: 127 (1753)                                      | 4b:18  |
|    | <b>Potamogeton perfoliatus</b> L., Sp. Pl. 1: 126 (1753)                                     | 4b:18  |
|    | <i>Potamogeton praelongus</i> sensu Rodway (1903) = <b>Potamogeton ochreatus</b>             | 4b:18  |
|    | <b>Potamogeton tricarinatus</b> F.Muell. & A.Benn. ex A.Benn., J. Bot. Lond. 30: 229 (1892)  | 4b:19  |
|    | <b>RESTIONACEAE</b>  |        |
|    | <i>Acion hookeri</i> (D.I.Morris) B.G.Briggs & L.A.S.Johnson = <b>Chordifex hookeri</b>      | 4b:46  |
|    | <i>Acion monocephalum</i> (R.Br.) B.G.Briggs & L.A.S.Johnson = <b>Chordifex monocephalum</b> | 4b:47  |
|    | <b>Apodasmia brownii</b> (Hook.f.) B.G.Briggs & L.A.S.Johnson, Telopea 7(4): 371 (1998)      | 4b:39  |
|    | <b>Baloskion australe</b> (R.Br.) B.G.Briggs & L.A.S.Johnson, Telopea 8(1): 23 (1998)        | 4b:46  |
|    | <b>Baloskion tetraphyllum</b> (Labill.) B.G.Briggs & L.A.S.Johnson, Telopea 8(1): 23 (1998)  | 4b:45  |
|    | subsp. <b>tetraphyllum</b>   |        |
|    | <i>Calorophus ater</i> L.A.S.Johnson & B.G.Briggs = <b>Calorophus erostris</b>               | 4b:42  |
|    | <b>Calorophus elongatus</b> Labill., Nov. Holl. Pl. 2: 78, t.228 (1806)                      | 4b:41  |
| e  | <b>Calorophus erostris</b> (C.B.Clarke) L.A.S.Johnson & B.G.Briggs in Curtis & Morris,       |        |

|                             |   |        |
|-----------------------------|---|--------|
|                             | Stud. Fl. Tasm. 4b: 425 (1994)  | 4b:42  |
|                             | <i>Calorophus lateriflorus</i> (R.Br.) F.Muell. = <b>Empodisma minus</b>                            | 4b:43  |
| ne                          | <b>Chordifex hookeri</b> (D.I.Morris) B.G.Briggs, <i>Telopea</i> 10(3): 685 (2004)                  | 4b:46  |
| ne                          | <b>Chordifex monocephalus</b> (R.Br.) B.G.Briggs, <i>Telopea</i> 10(3): 685 (2004)                  | 4b:47  |
|                             | <b>Empodisma minus</b> (Hook.f.) L.A.S.Johnson & D.F.Cutler, <i>Kew Bull.</i> 28: 383 (1974)        | 4b:43  |
|                             | <b>Eurychorda complanata</b> (R.Br.) B.G.Briggs & L.A.S.Johnson, <i>Telopea</i> 7(4): 359 (1998)    | 4b:46  |
|                             | <b>Hypolaena fastigiata</b> R.Br., <i>Prodr.</i> 251 (1810)   | 4b:40  |
|                             | <i>Hypolaena lateriflora</i> (R.Br.) Benth. = <b>Empodisma minus</b>                                | 4b:43  |
|                             | <i>Hypolaena longissima</i> sensu Rodway (1903) = <b>Calorophus elongatus</b>                       | 4b:41  |
|                             | <i>Leptocarpus brownii</i> Hook.f. = <b>Apodasmia brownii</b>                                       | 4b:39  |
|                             | <b>Leptocarpus tenax</b> (Labill.) R.Br., <i>Prodr.</i> 250 (1810)                                  | 4b:39  |
|                             | <b>Lepyrodia muelleri</b> Benth., <i>Fl. Austral.</i> 7: 215 (1878)                                 | 4b:43  |
|                             | <i>Lepyrodia tasmanica</i> Hook.f. = <b>Sporadanthus tasmanicus</b>                                 | 4b:44  |
|                             | <i>Restio australis</i> R.Br. = <b>Baloskion australe</b>   | 4b:46  |
|                             | <i>Restio complanatus</i> R.Br. = <b>Eurychorda complanata</b>                                      | 4b:46  |
|                             | <i>Restio glaber</i> (Rodway) L.A.S.Johnson & B.G.Briggs = <b>Chordifex monocephalus</b>            | 4b:47  |
|                             | <i>Restio gracilis</i> sensu Rodway (1903) = <b>Baloskion australe</b>                              | 4b:46  |
|                             | <i>Restio hookeri</i> D.I.Morris = <b>Chordifex hookeri</b>   | 4b:46  |
|                             | <i>Restio monocephalus</i> R.Br. = <b>Chordifex monocephalus</b>                                    | 4b:47  |
|                             | <i>Restio oligocephalus</i> F.Muell. = <b>Chordifex hookeri</b>                                     | 4b:46  |
|                             | <i>Restio tetraphyllum</i> Labill. = <b>Baloskion tetraphyllum</b>                                  | 4b:45  |
|                             | <b>Sporadanthus tasmanicus</b> (Hook.f.) B.G.Briggs & L.A.S.Johnson, <i>Telopea</i> 8(1): 31 (1998) | 4b:44  |
| e                           | <b>Winifredia sola</b> L.A.S.Johnson & B.G.Briggs, <i>Telopea</i> 2(6): 738 (1986)                  | 4b:47  |
| <br><b>RUPPIACEAE</b>       |   |        |
|                             | <i>Ruppia maritima</i> sensu Rodway (1903) = <b>Ruppia polycarpa</b>                                | 4b:21  |
|                             | <b>Ruppia megacarpa</b> R.Mason, <i>N.Z. J. Bot.</i> 5: 525 (1967)                                  | 4b:21  |
|                             | <b>Ruppia polycarpa</b> R.Mason, <i>N.Z. J. Bot.</i> 5: 524 (1967)                                  | 4b:21  |
|                             | <b>Ruppia tuberosa</b> J.S.Davis & Toml., <i>J. Arnold Arbor.</i> 55: 60 (1974)                     | 4b:21  |
| <br><b>TYPHACEAE</b>        |   |        |
|                             | <i>Typha angustifolia</i> sensu Rodway (1903) = <b>Typha domingensis &amp; T. orientalis</b>        |        |
|                             | <b>Typha domingensis</b> Pers., <i>Syn. Pl.</i> 2: 532 (1807)                                       | 4b:364 |
| i                           | <b>Typha latifolia</b> L., <i>Sp. Pl.</i> 2: 971 (1753)   | 4b:363 |
|                             | <b>Typha orientalis</b> C.Presl, <i>Abh. Bohm. Ges. Wiss. ser.5</i> 6: 599 (1851)                   | 4b:363 |
| <br><b>XANTHORRHOEACEAE</b> |   |        |
|                             | <i>Lomandra glauca</i> sensu J.H.Willis (1973) = <b>Lomandra nana</b>                               | 4b:422 |
|                             | <b>Lomandra longifolia</b> Labill., <i>Nov. Holl. Pl.</i> 1: 92, t.119 (1805)                       | 4b:422 |
|                             | <b>Lomandra nana</b> (A.Lee) A.Lee, <i>Fl. Australia</i> 46: 223 (1986)                             | 4b:422 |
| e                           | <b>Xanthorrhoea arenaria</b> Bedford, <i>Fl. Australia</i> 46: 225 (1986)                           | 4b:421 |
|                             | <b>Xanthorrhoea australis</b> R.Br., <i>Prodr.</i> 288 (1810)                                       | 4b:419 |
| e                           | <b>Xanthorrhoea bracteata</b> R.Br., <i>Prodr.</i> 288 (1810)                                       | 4b:421 |
|                             | <i>Xanthorrhoea minor</i> sensu Rodway (1903) = <b>Xanthorrhoea arenaria &amp; X. bracteata</b>     | 4b:421 |



|  |  |        |
|--|--|--------|
|  | <i>Xerotes glauca</i> sensu Rodway (1903) = <b>Lomandra nana</b>       | 4b:422 |
|  | <i>Xerotes longifolia</i> (Labill.) R.Br. = <b>Lomandra longifolia</b> | 4b:422 |

#### **XYRIDACEAE**

|   |  |       |
|---|--|-------|
|   | <i>Xyris gracilis</i> subsp. <i>tasmanica</i> D.I.Morris = <b>Xyris tasmanica</b>            | 4b:35 |
| e | <b>Xyris marginata</b> Rendle, J. Bot. 37: 503 (1899)  | 4b:35 |
| e | <b>Xyris muelleri</b> Malme, Svensk. Bot. Tidskr. 21: 381 (1928)                             | 4b:35 |
|   | <b>Xyris operculata</b> Labill., Nov. Holl. Pl. 1: 14, t.10 (1805)                           | 4b:34 |
| e | <b>Xyris tasmanica</b> (D.I.Morris) A.N.Doust & B.J.Conn, Aust. Syst. Bot. 10(2): 212 (1997) | 4b:35 |

#### **ZANNICHELLIACEAE**

|  |  |       |
|--|--|-------|
|  | <b>Lepilaena australis</b> Drumm. ex Harvey in Hooker's J. Bot. Kew Gard. Misc. 7: 58 (1855)     | 4b:22 |
|  | <b>Lepilaena bilocularis</b> Kirk, Trans. Proc. N.Z. Inst. 28: 500 (1896)                        | 4b:24 |
|  | <b>Lepilaena cylindrocarpa</b> (Koernicke in Walp.) Benth., Fl. Austral. 7: 180 (1878)           | 4b:22 |
|  | <b>Lepilaena marina</b> E.Robertson in Womersley, Marine Benthic Fl. Sthn. Austral. 1: 80 (1984) | 4b:23 |
|  | <b>Lepilaena patentifolia</b> E.Robertson, Fl. S.Austral. 4: 1736 (1986)                         | 4b:23 |
|  | <b>Lepilaena preissii</b> (Lehm.) F.Muell., Fragm. 8: 217 (1874)                                 | 4b:24 |

#### **ZOSTERACEAE**

|  |  |       |
|--|--|-------|
|  | <b>Heterozostera tasmanica</b> (Martens ex Asch.) Hartog, Seagrasses of the World 116 (1970) | 4b:28 |
|  | <b>Nanozostera muelleri</b> (Irmisch ex Asch.) Toml. & Posl., Taxon 50: 433 (2001)           | 4b:29 |
|  | <i>Zostera muelleri</i> Irmisch ex Asch. = <b>Nanozostera muelleri</b>                       | 4b:29 |
|  | <i>Zostera nana</i> sensu Rodway (1903) = <b>Nanozostera muelleri</b>                        | 4b:29 |
|  | <i>Zostera tasmanica</i> Martens ex Asch. = <b>Heterozostera tasmanica</b>                   | 4b:28 |

## **Gymnospermae**

#### **CUPRESSACEAE**

|   |  |     |
|---|--|-----|
| e | <b>Athrotaxis cupressoides</b> D.Don, Ann. Nat. Hist. 1: 234 (1838)                        | 1:6 |
| e | <b>Athrotaxis Xlaxifolia</b> Hook., Icon. Pl. 6: t.573 (1843)                              | 1:6 |
| e | <b>Athrotaxis selaginoides</b> D.Don, Ann. Nat. Hist. 1: 234 (1838)                        | 1:7 |
|   | <b>Callitris oblonga</b> Rich., Comm. Bot. Conif. Cycad. 47, t.18 fig.1 (1826)             | 1:5 |
| e | subsp. <b>oblonga</b>  |     |
|   | <b>Callitris rhomboidea</b> R.Br. ex Rich., Comm. Bot. Conif. Cycad. 47, t.18 fig.1 (1826) | 1:5 |
| e | <b>Diselma archeri</b> Hook.f., Fl. Tasm. 1: 353 t.98 (1857)                               | 1:5 |
|   | <i>Fitzroya archeri</i> Benth. & Hook.f. = <b>Diselma archeri</b>                          | 1:5 |

#### **PINACEAE**

|   |  |     |
|---|--|-----|
| i | <b>Pinus radiata</b> D.Don, Trans. Linn. Soc. Lond. 17: 442 (1837) | 1:8 |
|---|--|-----|

#### **PODOCARPACEAE**

|   |  |     |
|---|--|-----|
|   | <i>Dacrydium franklinii</i> Hook.f. = <b>Lagarostrobos franklinii</b>            | 1:3 |
| e | <b>Lagarostrobos franklinii</b> (Hook.f.) Quinn, Austral. J. Bot. 30: 316 (1982) | 1:3 |

|   |   |     |
|---|---|-----|
| e | <b>Microcachrys tetragona</b> (Hook.) Hook.f., Lond. J. Bot. 4: 150 (1845)                        | 1:3 |
| e | <b>Microstrobos niphophilus</b> J.Garden & L.A.S.Johnson, Contr. N.S.W. Natl. Herb. 1: 316 (1951) | 1:2 |
|   | <i>Pherosphaera hookeriana</i> sensu Rodway (1903) = <b>Microstrobos niphophilus</b>              | 1:2 |
| e | <b>Phyllocladus aspleniifolius</b> (Labill.) Rich. ex Hook.f., Lond. J. Bot. 4: 151 (1845)        | 1:2 |
|   | <i>Phyllocladus rhomboidalis</i> A. & L.Rich. = <b>Phyllocladus aspleniifolius</b>                | 1:2 |
|   | <i>Podocarpus alpinus</i> Hook.f. = <b>Podocarpus lawrencei</b>                                   | 1:4 |
|   | <b>Podocarpus lawrencei</b> Hook.f., Lond. J. Bot. 4: 151 (1845)                                  | 1:4 |

## Pteridophyta

### ADIANTACEAE

**Adiantum aethiopicum** L., Syst. Nat. ed.10, 2: 1329 (1759)

**Anogramma leptophylla** (L.) Link, Fil. Spec. 137 (1841)

**Cheilanthes austrotenuifolia** Quirk & Chambers, Austral. J. Bot. 31: 510 (1983)

**Cheilanthes distans** (R.Br.) Mett., Abh. Senckenberg. Naturf. Ges. 3: 69 (1859)

**Cheilanthes sieberi** Kunze in Lehm., Pl. Preiss. 2: 112 (1846)

subsp. **sieberi**

*Cheilanthes tenuifolia* sensu Rodway (1903) = **Cheilanthes austrotenuifolia**

**Pellaea calidirupium** Brownsey & Lovis, N.Z. J. Bot. 28: 197 (1990)

**Pellaea falcata** (R.Br.) Fée, Mem. Foug. 5: 129 (1850)

*Platyloma falcata* Smith = **Pellaea falcata**

### ASPLENIACEAE

**Asplenium appendiculatum** (Labill.) C.Presl, Tent. Pterid. 106 (1836)

subsp. **appendiculatum**

**Asplenium bulbiferum** Forst.f., Fl. Ins. Austral. 80 (1786)

subsp. **gracillimum** (Colenso) Brownsey, N.Z. J. Bot. 15: 60 (1977)

**Asplenium flabellifolium** Cav., Descr. Pl. 257 (1801)

**Asplenium flaccidum** Forst.f., Fl. Ins. Austral. 80 (1786)

subsp. **flaccidum**

**Asplenium hookerianum** Colenso, Tasm. J. Nat. Sci. 2: 169 (1844)

**Asplenium obtusatum** Forst.f., Fl. Ins. Austral. 80 (1786)

subsp. **northlandicum** Brownsey, N.Z. J. Bot. 15: 49, t.6 (1977)

*Asplenium terrestre* Brownsey = **Asplenium appendiculatum**

**Asplenium trichomanes** L., Sp. Pl. 2: 1080 (1753)

subsp. **quadrivalens** D.Meyer emend Lovis, Brit. Fern Gaz. 9: 152 (1964)

subsp. **trichomanes**

**Pleurosorus rutifolius** (R.Br.) Fée, Mem. Foug. 5: 179, t.16c (1852)

### ATHYRIACEAE

*Allantodia australis* R.Br. = **Diplazium australe**

*Athyrium australe* C.Presl = **Diplazium australe**

*Cystopteris fragilis* var. *tasmanica* (Hook.) Hook.f. = **Cystopteris tasmanica**

**Cystopteris tasmanica** Hook., Sp. Fil. 1: 199 (1846)

**Diplazium australe** (R.Br.) N.A.Wakef., Vict. Nat. 58: 142 t.3 (1942)

#### AZOLLACEAE

**Azolla filiculoides** Lam., Encycl. Méth. Bot. 1: 343 (1783)

#### BLECHNACEAE

*Blechnum aggregatum* sensu J.H.Willis (1973) = **Blechnum chambersii**

**Blechnum cartilagineum** Swartz, Syn. Fil. 114, 312 (1806)

**Blechnum chambersii** Tind. in Beadle et al., Fl. Sydney Region ed.2, 86 (1972)

**Blechnum fluviatile** (R.Br.) E.J.Löwe ex Salomon, Nomencl. Gefässkrypt. 115 (1883)

**Blechnum minus** (R.Br.) Ettingsh., Denkschr. Kaiserl. Akad. Wiss. Math.-Naturwiss. Kl. 23: 63, t.8 (1864)

**Blechnum nudum** (Labill.) Mett. ex Luer., Fl. Allgem. Bot. Zeit. 59: 292 (1876)

**Blechnum patersonii** (R.Br.) Mett., Fil. Hort. Bot. Lips. 64, t.4 figs.4-10 (1856)

subsp. **patersonii**

**Blechnum penna-marina** (Poiret) Kuhn, Fil. Afr. 92 (1868)

subsp. **alpina** (R.Br.) T.C.Chambers & P.A.Farrant, Fern Gaz. 15: 96 (1996)

*Blechnum procerum* sensu J.H.Willis (1973) = **Blechnum wattsii**

**Blechnum vulcanicum** (Blume) Kuhn in Miquel, Ann. Mus. Bot. Lugduno-Batavi 4: 284 (1869)

**Blechnum wattsii** Tind., Contr. N.S.W. Natl. Herb. 3: 247 (1963)

**Doodia australis** (Parris) Parris, Fl. Australia 48: 710 (1998)

**Doodia caudata** (Cav.) R.Br., Prodr. 151 (1810)

*Doodia media* (Tasmanian plants) = **Doodia australis**

*Lomaria alpina* Sprengel = **Blechnum penna-marina** subsp. **alpina**

*Lomaria discolor* sensu Rodway (1903) = **Blechnum nudum**

*Lomaria fluviatilis* Sprengel = **Blechnum fluviatile**

*Lomaria lanceolata* Sprengel = **Blechnum chambersii**

*Lomaria patersonii* Sprengel = **Blechnum patersonii**

*Lomaria procera* sensu Rodway (1903) = **Blechnum wattsii**

*Lomaria procera* var. *paludosa* Rodway = **Blechnum minus**

*Lomaria vulcanica* Blume = **Blechnum vulcanicum**

#### CULCITACEAE

**Calochlaena dubia** (R.Br.) M.D.Turner & R.A.White, Amer. Fern J. 78: 92 (1988)

*Culcita dubia* (R.Br.) Maxon = **Calochlaena dubia**

*Davallia dubia* R.Br. = **Calochlaena dubia**

#### CYATHEACEAE

*Alsophila australis* R.Br. = **Cyathea australis**

**Cyathea australis** (R.Br.) Domin, Pteridophyta 262 (1929)

**Cyathea cunninghamii** Hook.f., Icon. Pl. 10, t.985 (1854)

**Cyathea Xmarcescens** N.A.Wakef., Vict. Nat. 59: 33, figs.1-5 (1942)

#### DAVALLIACEAE

*Davallia dubia* R.Br. = **Calochlaena dubia** (Culcitaceae)

## DENNSTAEDTIACEAE

**Histiopteris incisa** (Thunb.) J.Smith, Hist. Fil. 295 (1875)

**Hypolepis amaurobachis** (Kunze) Hook., Sp. Fil. 2: 62 (1852)

*Hypolepis australis* N.A.Wakef. = **Hypolepis amaurobachis**

t **Hypolepis distans** Hook., Sp. Fil. 2: 70, t.95c (1852)

**Hypolepis glandulifera** Brownsey & Chinnock, J. Adelaide Bot. Gard. 10: 16, fig.8 (1987)

**Hypolepis muelleri** N.A.Wakef., Vict. Nat. 60: 42 (1943)

*Hypolepis punctata* sensu J.H.Willis (1973) = **Hypolepis glandulifera**

**Hypolepis rugosula** (Labill.) J.Smith, Comp. Bot. Mag., n.s. 2: 8 (1846)

*Hypolepis tenuifolia* sensu Rodway (1903) = **Hypolepis amaurobachis** & **H. rugosula**

*Polypodium punctatum* sensu Rodway (1903) = **Hypolepis rugosula** & **H. amaurobachis**

**Pteridium esculentum** (Forst.f.) Cockayne, Rep. Bot. Survey Tongariro Natl. Park 34 (1908)

## DICKSONIACEAE

**Dicksonia antarctica** Labill., Nov. Holl. Pl. 2: 100, t.249 (1807)

## DRYOPTERIDACEAE

**Lastreopsis acuminata** (Houlston) Morton, Contr. U.S. Natl. Herb. 38: 245 (1973)

**Lastreopsis hispida** (Swartz) Tind., Vict. Nat. 73: 183 (1957)

*Lastreopsis shepherdii* (Kuntze ex Mett.) Tind. = **Lastreopsis acuminata**

*Nephrodium decompositum* sensu Rodway (1903) = **Lastreopsis acuminata**

*Nephrodium hispidum* (Swartz) Hook. = **Lastreopsis hispida**

*Polystichum coriaceum* Schott = **Rumohra adiantiformis**

**Polystichum proliferum** (R.Br.) C.Presl, Tent. Pterid. 83 (1836)

*Polystichum vestitum* sensu Rodway (1903) = **Polystichum proliferum**

**Rumohra adiantiformis** (Forst.f.) Ching, Sinensia 5: 70 (1934)

## GLEICHENIACEAE

e **Gleichenia abscida** Rodway, Tasm. Fl. 289 (1903)

t **Gleichenia alpina** R.Br., Prodr. 161 (1810)

*Gleichenia circinata* sensu Rodway (1903) = **Gleichenia microphylla**

**Gleichenia dicarpa** R.Br., Prodr. 161 (1810)

*Gleichenia flabellata* sensu Rodway (1903) = **Sticherus tener** & **S. urceolatus**

**Gleichenia microphylla** R.Br., Prodr. 161 (1810)

**Sticherus lobatus** N.A.Wakef., Vict. Nat. 60: 110 (1943)

**Sticherus tener** (R.Br.) Ching, Sunyatsenia 5: 283 (1940)

**Sticherus urceolatus** M.Garrett & Kantvilas in M.Garrett, G.Kantvilas & H.Laws, Muelleria 11: 103 (1998)

## GRAMMITIDACEAE

**Ctenopteris heterophylla** (Labill.) Tind., Amer. Fern J. 41: 100, t.6 (1951)

*Grammitis armstrongii* Tind. = **Grammitis poeppigiana**

*Grammitis australis* R.Br. = **Grammitis billardierei**

**Grammitis billardierei** Willd., Sp. Pl. ed.4 5(1): 139 (1810)

e **Grammitis garrettii** Parris, Fl. Australia 48: 713 (1998)

- t **Grammitis gunnii** Parris, Fl. Australia 48: 713 (1998)  
**Grammitis magellanica** Desv., Berl. Mag. 5: 313 (1811)  
 subsp. **nothofageti** Parris in Parris & Given, N.Z. J. Bot. 14: 100 (1976)  
 Grammitis meridionalis Parris = **Grammitis magellanica** subsp. **nothofageti** p.p.maj.  
**Grammitis poeppigiana** (Mett.) Pichi-Serm., Webbia 32: 461 (1978)
- t **Grammitis pseudociliata** Parris in Parris & Given, N.Z. J. Bot. 14: 106 (1976)  
 Polypodium grammitidis R.Br. = **Ctenopteris heterophylla**

#### HEMIONITIDIACEAE

- Gymnogramme leptophylla (L.) Desv. = **Anogramma leptophylla** (Adiantaceae)  
 Gymnogramme rutaefolia Hook. = **Pleurosorus rutifolius** (Aspleniaceae)

#### HYMENOPHYLLACEAE

- Apteropteris applanata A.M.Gray & R.G.Williams = **Sphaerocionium applanatum**  
**Crepidomanes venosum** (R.Br.) Bostock, Fl. Australia 48: 706 (1998)  
**Hymenophyllum australe** Willd., Sp. Pl. ed.5, 5: 527 (1810)  
**Hymenophyllum cupressiforme** Labill., Nov. Holl. Pl. 2: 102, t.250 fig.2 (1806)  
**Hymenophyllum flabellatum** Labill., Nov. Holl. Pl. 2: 101, t.250 fig.1 (1806)  
 Hymenophyllum javanicum sensu Rodway (1903) = **Hymenophyllum australe**  
 Hymenophyllum malingii sensu Rodway (1903) = **Sphaerocionium applanatum**  
**Hymenophyllum marginatum** Hook. & Grev., Icon. Fil. 1: t.34 (1827)  
**Hymenophyllum peltatum** (Poiret) Desv., Prodr. 333 (1827)  
**Hymenophyllum rarum** R.Br., Prodr. 159 (1810)  
 Hymenophyllum tunbridgense sensu Rodway (1903) = **Hymenophyllum cupressiforme**  
 Hymenophyllum wilsoni sensu Rodway (1903) = **Hymenophyllum peltatum**  
 Mecodium australe (Willd.) Copel. = **Hymenophyllum australe**  
 Mecodium flabellatum (Labill.) Copel. = **Hymenophyllum flabellatum**  
 Mecodium rarum (R.Br.) Copel. = **Hymenophyllum rarum**  
 Polyphlebium venosum (R.Br.) Copel. = **Crepidomanes venosum**
- e **Sphaerocionium applanatum** (A.M.Gray & R.G.Williams) K.Iwats., J. Fac. Sci. Univ. Tokyo, sect.3, Bot. 13: 214 (1982)  
 Trichomanes venosum R.Br. = **Crepidomanes venosum**

#### ISOETACEAE

- Isoetes drummondii** A.Braun, Monatsber. K. Akad. Wiss. Berlin 593 (1868)  
 subsp. **drummondii**
- e **Isoetes elatior** F.Muell. ex A.Braun, Linnaea 25: 722 (1853)  
 e **Isoetes gunnii** A.Braun, Monatsber. K. Akad. Wiss. Berlin 535 (1868)  
 e **Isoetes humilior** F.Muell. ex A.Braun, Linnaea 25: 722 (1853)  
 Isoetes lacustris sensu Rodway (1903) = **Isoetes elatior**, **I. humilior** & **I. gunnii**  
**Isoetes muelleri** A.Braun, Monatsber. K. Akad. Wiss. Berlin 541 (1868)

#### LINDSAEACEAE

- Lindsaea cuneata (Forst.f.) Chr. = **Lindsaea trichomanoides**  
**Lindsaea linearis** Swartz, J. Bot. (Schrader) 1800(2): 78 (1801)

**Lindsaea trichomanoides** Dryand., Trans. Linn. Soc. London 3: 43, t.11 (1797)

Lindsaya = **Lindsaea**

#### LYCOPODIACEAE

**Huperzia australiana** (Herter) Holub, Folia Geobot. Phytotax. 20: 70 (1985)

**Huperzia varia** (R.Br.) Trev. Atti Soc. Ital. Sci. Nat. 17: 247 (1874)

t **Lycopodiella diffusa** (R.Br.) B.Øllg., Opera Bot. 92: 176 (1987)

**Lycopodiella lateralis** (R.Br.) B.Øllg., Opera Bot. 92: 176 (1987)

**Lycopodiella serpentina** (Kunze) B.Øllg., Opera Bot. 92: 176 (1987)

*Lycopodium australianum* Herter = **Huperzia australiana**

*Lycopodium clavatum* L.

var. *fastigiatum* (R.Br.) Benth. = **Lycopodium fastigiatum**

*Lycopodium densum* Labill. = **Lycopodium deuterodensum**

**Lycopodium deuterodensum** Herter, Estud. Bot. (Montevideo) 20: 15 (1949)

**Lycopodium fastigiatum** R.Br., Prodr. 165 (1810)

*Lycopodium laterale* R.Br. = **Lycopodiella lateralis**

*Lycopodium myrtifolium* auct. non Forst.f. = **Huperzia varia**

**Lycopodium scariosum** Forst.f., Fl. Ins. Austr. 87 (1786)

*Lycopodium selago* sensu Rodway (1903) = **Huperzia australiana**

*Lycopodium serpentinum* Kunze = **Lycopodiella serpentina**

*Lycopodium varium* R.Br. = **Huperzia varia**

**Phylloglossum drummondii** Kunze, Bot. Zeitung (Berlin) 1: 721 (1843)

#### MARSILEACEAE

i **Marsilea mutica** Mett., Ann. Sci. Nat. Bot. ser.4, 15: 88 (1861)

*Pilularia globulifera* sensu Rodway (1903) = **Pilularia novae-hollandiae**

**Pilularia novae-hollandiae** A.Braun, Monatsber. Konigl. Preuss. Akad. Wiss. Berlin 1863: 435 (1864)

#### OPHIOGLOSSACEAE

x **Botrychium australe** R.Br., Prodr. 164 (1810)

**Botrychium lunaria** (L.) Swartz, J. Bot. (Schrader) 1800(2): 110 (1801)

*Botrychium ternatum* sensu Rodway (1903) = **Botrychium australe**

**Ophioglossum lusitanicum** L., Sp. Pl. 2: 1063 (1753)

subsp. **coriaceum** (A.Cunn.) R.T.Clausen, Mem. Torrey Bot. Club 19(2): 161 (1938)

#### OSMUNDACEAE

**Todea barbara** (L.) T.Moore, Ind. Fil. 119 (1857)

#### POLYPODIACEAE

*Microsorium diversifolium* (Willd.) Copel. = **Microsorium pustulatum**

**Microsorium pustulatum** (Forst.f.) Copel., Univ. Calif. Publ. Bot. 16: 114 (1929)

subsp. **pustulatum**

*Phymatodes billardierei* C.Presl = **Microsorium pustulatum**

*Phymatosorus pustulatus* (Forst.f.) Large = **Microsorium pustulatum**

*Polypodium grammitidis* R.Br. = **Ctenopteris heterophylla** (Grammitidaceae)

*Polypodium punctatum* sensu Rodway (1903) = **Hypolepis rugosula** p.p.maj. (Dennstaedtiaceae)

#### **PSILOTACEAE**

*Tmesipteris billardi* Endl. = **Tmesipteris obliqua**

**Tmesipteris elongata** Dangeard, *Botaniste* ser.2: 213 (1890-91)

**Tmesipteris obliqua** Chinnock, *Muelleria* 8: 60 (1993)

**Tmesipteris parva** N.A.Wakef., *Vict. Nat.* 60: 143 (1944)

*Tmesipteris tannensis* sensu Rodway (1903) = **Tmesipteris parva**, **T. elongata** & **T. obliqua**

*Tmesipteris tungana* H.N.Barber = **Tmesipteris elongata**

#### **PTERIDACEAE**

*Litobrochia comans* C.Presl = **Pteris comans**

*Litobrochia incisa* C.Presl = **Histiopteris incisa** (Dennstaedtiaceae)

*Pteris aquilina* sensu Rodway (1903) = **Pteridium esculentum** (Dennstaedtiaceae)

**Pteris comans** Forst.f., *Fl. Ins. Austral.* 79 (1786)

**Pteris tremula** R.Br., *Prodr.* 154 (1810)

#### **SCHIZAEACEAE**

**Schizaea asperula** N.A.Wakef., *Vict. Nat.* 59: 89 (1942)

**Schizaea bifida** Willd., *Abh. Kurfurstl.-Mainz. Akad. Nützl. Wiss. Erfurt* 5: 30, t.3 fig.3 (1802)

**Schizaea fistulosa** Labill., *Nov. Holl. Pl.* 2: 103, t.250 fig.3 (1807)

#### **SELAGINELLACEAE**

**Selaginella gracillima** (Kunze) Spring ex Salomon, *Nomencl.* 353 (1883)

*Selaginella preissiana* sensu Rodway (1903) = **Selaginella gracillima**

**Selaginella uliginosa** (Labill.) Spring, *Bull. Acad. Roy. Sci. Bruxelles* 10: 136 (1843)

#### **THELYPTERIDACEAE**

*Cyclosorus penniger* (Forst.f.) Ching = **Pneumatopteris pennigera**

**Pneumatopteris pennigera** (Forst.f.) Holtt., *Blumea* 21: 305 (1973)

## **The Vascular Flora of Macquarie Island**

### **Dicotyledonae**

#### **APIACEAE**

**Azorella macquariensis** Orch., *Muelleria* 7: 16 (1989)

**Hydrocotyle novae-zeelandiae** DC., *Prodr.* 4: 67 (1830)

**Stilbocarpa polaris** (Hombr. & Jacquinot ex Hook.f.) A.Gray, Bot. U.S. Explor. Exped. Bot. Phan. 7: 4 (1854)

#### ASTERACEAE

*Cotula plumosa* (Hook.f.) Hook.f. = **Leptinella plumosa**

**Leptinella plumosa** Hook.f., Fl. Antarct. 1: 26 (1844)

**Pleurophyllum hookeri** Buchan., Trans. N.Z. Inst. 16: 395 (1884)

#### BRASSICACEAE

**Cardamine corymbosa** Hook.f., Fl. Antarct. 1: 6 (1844)

#### CALLITRICHACEAE

**Callitriche antarctica** Engelm. ex Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 20 (1867)

#### CARYOPHYLLACEAE

i **Cerastium fontanum** Baumg., Enum. Stirp. Transs. 1: 425 (1816)

    subsp. **fontanum**

**Colobanthus affinis** (Hook.) Hook.f., Fl. Tasm. 1: 45 (1855) 1:74

**Colobanthus apetalus** (Labill.) Druce, Bot. Exch. Club Soc. Brit. Isles 1916: 616 (1917) 1:74

    var. **alpinus** (Kirk) L.B.Moore in H.H.Allan, Fl. New Zealand 1: 214 (1961)

**Colobanthus muscoides** Hook.f., Fl. Antarct. 1: 14 (1844)

i **Stellaria media** (L.) Cirillo, Essent. Pl. Char. Comment. 36 (1784) 1:71

**Stellaria parviflora** Banks & Sol. ex Hook.f., Fl. Nov.-Zel. 1: 25 (1852)

#### CRASSULACEAE

t **Crassula moschata** Forst.f., Comm. Gotting. 9: 26 (1789)

#### HALORAGACEAE

**Myriophyllum triphyllum** Orch., Brunonia 2: 259 (1980)

#### ONAGRACEAE

**Epilobium brunnescens** (Cockayne) Raven & Engelhorn, N.Z. J. Bot. 9: 350 (1971)

    subsp. **brunnescens**

**Epilobium pedunculare** A.Cunn., Ann. Nat. Hist. 3: 31 (1839)

#### POLYGONACEAE

i **Rumex crispus** L., Sp. Pl. 1: 335 (1753) 3:584

#### PORTULACACEAE

?i **Montia fontana** L., Sp. Pl. 1: 87 (1753) 1:79

    subsp. **fontana**

#### RANUNCULACEAE

**Ranunculus crassipes** Hook.f., Fl. Antarct. 1: 224 (1845)



## ROSACEAE

*Acaena magellanica* (Lam.) Vahl, Enum. Pl. 1: 297 (1804)

*Acaena minor* (Hook.f.) Allan, Fl. New Zealand 1: 363 (1961)

## RUBIACEAE

*Coprosma perpusilla* Colenso, Trans. N.Z. Inst. 22: 466 (1890)

subsp. *subantarctica* Orch., Brunonia 9: 133 (1987)

*Galium antarcticum* Hook.f., Fl. Antarct. 1: 303 (1846)

## Monocotyledonae

### CYPERACEAE

*Carex trifida* Cav., Icon. 5: 41, t.465 (1799)

*Isolepis aucklandica* Hook.f., Fl. Antarct. 1: 88 (1844), t.50 (1845)

4b:102

*Uncinia divaricata* Boott in Hook.f., Fl. Nov.-Zel. 1: 286 (1853)

*Uncinia hookeri* Boott in Hook.f., Fl. Antarct. 1: 91 (1844); t.51 (1845)

### JUNCACEAE

*Juncus scheuchzerioides* Gaudich., Ann. Sci. Nat. (Paris) 5: 100 (1825)

*Luzula crinita* Hook.f., Fl. Antarct. 1: 84, t.48 (1844)

### ORCHIDACEAE

*Corybas dienemus* D.L.Jones = ***Nematoceras dienema***

*Corybas macranthus* sensu W.M.Curtis (1979) = ***Nematoceras dienema***

4a:121

***Nematoceras dienema*** (D.L.Jones) D.L.Jones, M.A.Clem. & Molloy, Orchadian 13(10): 449 (2002)

4a:121

### POACEAE

*Agrostis magellanica* Lam., Tabl. Encycl. 1: 160 (1791)

i *Anthoxanthum odoratum* L., Sp. Pl. 1: 28 (1753)

4b:250

*Deschampsia cespitosa* (L.) P.Beauv., Ess. Agrostogr. 91: t.18 fig.3 (1812)

4b:242

*Deschampsia chapmanii* Petrie, Trans. & Proc. N.Z. Inst. 23: 401 (1891)

*Festuca contracta* Kirk, Trans. & Proc. N.Z. Inst. 27: 353 (1895)

i *Poa annua* L., Sp. Pl. 1: 68 (1753)

4b:211

*Poa cookii* (Hook.f.) Hook.f., Philos. Trans. 168: 22 (1879)

*Poa foliosa* (Hook.f.) Hook.f., Handb. N.Z. Fl. 338 (1864)

*Poa hamiltonii* Kirk = ***Poa cookii***

*Poa litorosa* Cheeseman, Man. N.Z. Fl. 1156 (1906)

*Puccinellia macquariensis* (Cheeseman) Allan & Jansen, Trans. & Proc. Roy. Soc. N.Z. 69: 268 (1939)

## Pteridophyta

### BLECHNACEAE

**Blechnum penna-marina** (Poiret) Kuhn, Fil. Afr. 92 (1868)

subsp. **alpina** (R.Br.) T.C.Chambers & P.A.Farrant, Fern Gaz. 15: 96 (1996)

**DRYOPTERIDACEAE**

**Polystichum vestitum** (Forst.f.) C.Presl, Tent. Pterid. 83 (1836)

**GRAMMITIDACEAE**

**Grammitis poeppigiana** (Mett.) Pichi-Serm., Webbia 32: 461 (1978)

**HYMENOPHYLLACEAE**

**Hymenophyllum falklandicum** Baker in Hook. & J.G.Baker, Syn. Fil. edn.2 68 (1874)

**LYCOPODIACEAE**

**Huperzia australiana** (Herter) Holub, Folia Geobot. Phytotax. 20: 70 (1985)