

# A CENSUS OF THE VASCULAR PLANTS OF TASMANIA, INCLUDING MACQUARIE ISLAND

MF de Salas & ML Baker

2017 edition

Tasmanian Vascular Plant Census 2017

A Census of the Vascular Plants of Tasmania, including Macquarie Island.

2017 edition

MF de Salas and ML Baker

Postal address:

Tasmanian Herbarium  
PO Box 5058  
UTAS LPO  
Sandy Bay, Tasmania 7005  
Australia

Street address:

College Road  
Sandy Bay, Tasmania 7005  
Australia

© Tasmanian Herbarium, Tasmanian Museum and Art Gallery

Published by the Tasmanian Herbarium, Tasmanian Museum and Art Gallery

GPO Box 1164  
Hobart, Tasmania 7001  
Australia  
[www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)

Cite as: de Salas, M.F. and Baker, M.L. (2017) *A Census of the Vascular Plants of Tasmania, including Macquarie Island*.  
(Tasmanian Herbarium, Tasmanian Museum and Art Gallery, Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)

ISBN 978-1-921599-84-2 (PDF)

## Introduction

The classification systems used in this Census largely follow Cronquist (1981) for flowering plants (Angiosperms) and McCarthy (1998) for conifers, ferns and their allies. The same systems are used to arrange the botanical collections of the Tasmanian Herbarium and by the *Flora of Australia* series published by the Australian Biological Resources Study (ABRS). For a more up-to-date classification of the flora, refer to *The Flora of Tasmania Online* (Duretto 2009+) which currently follows APG II (2003). To determine the families in which genera are placed, refer to Appendix 2 at the end of this document.

This census also serves as an index to *The Student's Flora of Tasmania* (Curtis 1963, 1967, 1979; Curtis & Morris 1975, 1994). Species accounts can be found in *The Student's Flora of Tasmania* by referring to the volume and page number reference that is given in the far right column (e.g. 3:539). Families with accounts completed in the *Flora of Tasmania Online* (Duretto 2009+) are highlighted at the family entry (e.g. FTO 1).

The *Census of the Vascular Plants of Tasmania* incorporates every name (including synonyms) used to refer to Tasmanian plants in the major taxonomic publications about Tasmania, and tries to account for every species name attributed to Tasmania, whether erroneously or correctly. Names from the following publications have been included: Jacques-Julien Houtou de Labillardière's *Novae Hollandiae Plantarum Specimen* (1804-1807), Robert Brown's *Prodromus Florae Novae Hollandiae et Insulae van-Diemen* (1810), Joseph Dalton Hooker's *The Botany of the Antarctic Voyage of H. M. Discovery Ships Erebus and Terror in the Years 1839 - 1843 Under the Command of Captain Sir James Clark Ross. Part III: Flora Tasmaniae* (1855-1859, complete work published in 1860), all volumes of George Bentham's *Flora Australiensis* (1863-1878) and Leonard Rodway's *The Tasmanian Flora* (1903). In addition, all names used in Cheeseman's (1919) *The Vascular Flora of Macquarie Island* have been accounted for in the Macquarie Island section.

According to this *Census*, the Tasmanian flora contains 2719 vascular plants, of which 1918 (70%) are considered native and 801 (30%) have naturalised from elsewhere. Among the native taxa, 530 (28%) are endemic to the State. Thirty-seven of the State's exotic taxa, are considered sparingly naturalised, and are known only from a small number of populations. Twenty-four native taxa are recognised as extinct, whereas 8 naturalised taxa are considered to have either not persisted in Tasmania or have been eradicated. The sub-antarctic Macquarie Island, considered part of Tasmania, supports 49 species of vascular plants, of which 42 are considered native and 7 naturalised. For some basic statistics on the Tasmanian flora see Tables 1–3.

Five new native taxa are recognised in the 2017 edition, three at the rank of subspecies and two at the rank of species (Appendix 1a). The names of several taxa have changed since the previous edition (Appendix 1b), including the recently described *Pimelea leiophylla*, a new endemic species recognised for the State. Several taxa have had their status changed since the previous edition (Appendix 1c). Twenty-one taxa were previously considered sparingly naturalised and are now considered to be fully naturalised. *Cardamine tryssa* was previously considered to be extinct in Tasmania, and has been recently re-discovered. Compared to the 2016 *Census*, this edition contains 79 fewer exotic taxa (Appendix 1d). These were listed in previous editions as *sparingly naturalised*, but a recent review by Baker *et al.* (in prep.) has found no conclusive evidence to demonstrate that they have, in fact, ever become naturalised. This review has led to numerous taxa having status changes in the 2017 edition and to the inclusion of a new status of *doubtfully naturalised*. Finally, the species, *Leptecophylla juniperina*, is no longer considered to occur in Tasmania.

## Acknowledgments

This checklist was developed using BRAHMS (Botanical Research And Herbarium Management System) database management system. The authors would like to thank the BRAHMS team for making possible (and infinitely easier) the publication of an annual up-to-date census. In addition the authors would like to thank the APC (Australian Plant Census) group, whose work brings to our attention a myriad of taxonomic updates we might otherwise miss. We would like to thank Lynette Cave of the Tasmanian Herbarium for her assistance in checking the validity and publication details of numerous taxa. Mark Wapstra (ECOtas) and Wendy Potts (DPIPWE) provided useful comments on drafts of this publication and for these we are grateful.

**Table 1: Some basic statistics on the status of Tasmanian vascular plants**

		Native		Naturalised
		non-endemic	endemic	
<b>Dicotyledons</b>	<b>1739</b>	799	363	577
<b>Monocotyledons</b>	<b>860</b>	492	150	218
<b>Gymnosperms</b>	<b>15</b>	2	9	4
<b>Pteridophytes</b>	<b>105</b>	95	8	2
<b>TOTAL</b>	<b>2719</b>	<b>1388</b>	<b>530</b>	<b>801</b>

**Table 2: Some basic statistics on the status of Macquarie Island vascular plants**

		Native		Naturalised
		non-endemic	endemic	
<b>Dicotyledons</b>	<b>24</b>	20	1	3
<b>Monocotyledons</b>	<b>20</b>	13	3	4
<b>Gymnosperms</b>	<b>0</b>	0	0	0
<b>Pteridophytes</b>	<b>5</b>	5	0	0
<b>Total</b>	<b>49</b>	<b>38</b>	<b>4</b>	<b>7</b>

**Table 3: Statistics of Tasmanian and Macquarie Island vascular plant families and genera**

	Tasmanian families	Tasmanian genera	Macquarie Island families	Macquarie Island genera
<b>Dicotyledons</b>	109	551	13	19
<b>Monocotyledons</b>	29	241	4	12
<b>Gymnosperms</b>	3	10	0	0
<b>Pteridophytes</b>	26	43	5	5
<b>Total</b>	<b>167</b>	<b>845</b>	<b>22</b>	<b>36</b>

**Symbols used in text:**

- e endemic in Tasmania
- t within Australia, occurs only in Tasmania
- x considered by the Tasmanian Herbarium to be extinct or eradicated in Tasmania or Macquarie Island
- i introduced and naturalised in Tasmania
- # sparingly naturalised or known from only one or two populations or collections
- \* doubtfully naturalised: collections exist but no evidence exists to establish if a taxon is naturalised in Tasmania
- n a change since the 2016 edition (see Appendix I for summary)

no symbol indicates that the taxon is native in Tasmania and the Australian mainland

NOTE: The 2017 Census no longer includes information on the listing status of species listed on the Tasmanian *Threatened Species Protection Act 1995*, the *Commonwealth Environment Protection and Biodiversity Conservation Act 1999*, or the *Tasmanian Weed Management Act 1999*. For this information please refer to current legislation.

# Tasmanian Vascular Plant Census

## DICOTYLEDONEAE

	<b>ACANTHACEAE</b>	FTO 120
i	<b>Acanthus mollis</b> L., Sp. Pl. 2: 639 (1753)	3:539
	<b>ACERACEAE</b>	FTO 86
i	<b>Acer pseudoplatanus</b> L., Sp. Pl. 2: 1054 (1753)	
	<b>AIZOACEAE</b>	FTO 101
	<i>Aptenia cordifolia</i> (L.f.) Schwantes = <b>Mesembryanthemum cordifolium</b>	
i #	<b>Carpobrotus aequilaterus</b> (Haw.) N.E.Br., J. Bot. 66: 324 (1928)	
i	<b>Carpobrotus edulis</b> (L.) N.E.Br., Gen. S. Afr. Fl. Pl. 249 (1926)	2:238
	<b>Carpobrotus rossii</b> (Haw.) Schwantes, Gartenflora 77: 68 (1928)	2:238
	<i>Disphyma australe</i> (Aiton) N.E.Br. sensu Curtis (1963) = <b>Disphyma crassifolium</b> subsp. <b>clavellatum</b> (misapplied in Tasmania)	2:239
	<b>Disphyma crassifolium</b> (L.) L.Bolus subsp. <b>clavellatum</b> (Haw.) Chinnock, Fl. S. Austral., ed. 4, 1: 194 (1986)	2:239
i	<b>Drosanthemum candens</b> (Haw.) Schwantes, Z. Sukkulantenk. 3: 29 (1927)	
i	<b>Galenia pubescens</b> (Eckl. & Zeyh.) Druce var. <b>pubescens</b> , Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 624 (1917)	2:241
	<i>Galenia secunda</i> (L.f.) Sond. sensu Curtis (1963) = <b>Galenia pubescens</b> var. <b>pubescens</b> (misapplied in Tasmania)	2:241
i	<b>Lampranthus glaucus</b> (L.) N.E.Br., Gard. Chron. 87: 212 (1930)	2:240
	<i>Mesembryanthemum aequilaterale</i> Haw. sensu Rodway (1903) = <b>Carpobrotus rossii</b> (misapplied in Tasmania)	2:238
	<i>Mesembryanthemum australe</i> Sol. sensu Rodway (1903) = <b>Disphyma crassifolium</b> subsp. <b>clavellatum</b> (misapplied in Tasmania)	2:239
i #	<b>Mesembryanthemum cordifolium</b> L.f., Suppl. Pl. 260 (1782)	
i	<b>Mesembryanthemum crystallinum</b> L., Sp. Pl. 1: 480 (1753)	
	<i>Tetragonia expansa</i> Murray = <b>Tetragonia tetragonoides</b>	2:240
	<b>Tetragonia implexicoma</b> (Miq.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 148 (1856)	2:241
	<b>Tetragonia tetragonoides</b> (Pall.) Kuntze, Revis. Gen. Pl. 1: 264 (1891)	2:240
	<b>AMARANTHACEAE</b>	FTO 98
	<b>Alternanthera denticulata</b> R.Br., Prodr. Fl. Nov. Holland. 417 (1810)	3:568
	<i>Alternanthera sessilis</i> (L.) R.Br. ex DC. sensu Hooker (1860) = <b>Alternanthera denticulata</b> (misapplied in Tasmania)	
i	<b>Amaranthus albus</b> L., Syst. Nat., ed. 10: 1268 (1759)	
i	<b>Amaranthus deflexus</b> L., Mant. Pl. 2: 295 (1771)	3:567
n	<i>Amaranthus graecizans</i> L. subsp. <i>silvestris</i> (Vill.) Brenan – previously listed as naturalised but insufficient evidence exists to support this	
	<i>Amaranthus hybridus</i> L. sensu Curtis (1967) = <b>Amaranthus powellii</b> (misapplied in Tasmania)	3:566
i	<b>Amaranthus powellii</b> S.Watson, Proc. Amer. Acad. Arts. 10: 347 (1875)	3:566

	Amaranthus retroflexus L. sensu Curtis (1967) = <b>Amaranthus powellii</b> (misapplied in Tasmania)	3:566
n	Amaranthus spinosus L. – previously listed as naturalised but insufficient evidence exists to support this	
	<b>Hemichroa pentandra</b> R.Br., Prodr. Fl. Nov. Holland. 409 (1810)	3:576
	<b>Ptilotus spathulatus</b> (R.Br.) Poir., Encycl. (Lamarck) Suppl. 4: 620 (1816)	3:567
	Trichinium spathulatum R.Br. = <b>Ptilotus spathulatus</b>	
<b>APIACEAE (UMBELLIFERAE)</b>		
	Aciphylla procumbens (F.Muell.) Benth. = <b>Anisotome procumbens</b>	2:261
	<b>Actinotus bellidioides</b> (Hook.f.) Benth., Fl. Austral. 3: 369 (1867)	2:253
e	<b>Actinotus moorei</b> F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania 1895: 65 (1896)	2:254
e	<b>Actinotus suffocatus</b> (Hook.f.) Rodway, Pap. & Proc. Roy. Soc. Tasmania 1893: 180 (1894)	2:254
n	Aegopodium podagraria L. – previously listed as naturalised but insufficient evidence exists to support this	
i	<b>Ammi majus</b> L., Sp. Pl. 1: 243 (1753)	
e	<b>Anisotome procumbens</b> (F.Muell.) C.J.Webb, Fl. & Fauna Alpine Australasia: 395 (1986)	2:261
i	<b>Anthriscus caucalis</b> M.Bieb., Fl. Taur.-Caucas. 1: 230 (1808)	
	<b>Apium annum</b> P.S.Short, J. Adelaide Bot. Gard. 1: 230 (1979)	
	Apium australe Thouars sensu Bentham (1867) misapplied to <b>A. annum</b> , <b>A. insulare</b> & <b>A. prostratum</b>	
	<b>Apium insulare</b> P.S.Short, J. Adelaide Bot. Gard. 1: 228 (1979)	
	<b>Apium prostratum</b> Labill. ex Vent. subsp. <b>prostratum</b> var. <b>filiforme</b> (A.Rich.) Kirk, Stud. Fl. New Zealand: 196 (1899)	2:255
	<b>Apium prostratum</b> Labill. ex Vent. subsp. <b>prostratum</b> var. <b>prostratum</b> , Jard. Malmaison 2: t.81 (1805)	2:255
	Azorella dichopetala Benth. nom. illeg. = <b>Dichosciadium ranunculaceum</b>	
	Azorella saxifraga (Hook.f.) Benth. = <b>Oschatzia saxifraga</b>	2:251
	Caldasia brachycarpa Hook.f. = <b>Oreomyrrhis eriopoda</b>	
	Caucalis infesta Curtis = Torilis arvensis	
	<b>Centella cordifolia</b> (Hook.f.) Nannf., Svensk Bot. Tidskr. 18: 418 (1924)	2:247
i	<b>Conium maculatum</b> L., Sp. Pl. 1: 243 (1753)	2:254
	Crantzia lineata Nutt. sensu Bentham (1867) = <b>Lilaeopsis polyantha</b> (misapplied in Tasmania)	
i	<b>Cyclosporum leptophyllum</b> (Pers.) Sprague ex Britton & P.Wilson, Bot. Porto Rico 6: 52 (1925)	
	Daucus brachiatus Sieber ex DC. = <b>Daucus glochidiatus</b>	
i	<b>Daucus carota</b> L., Sp. Pl. 1: 242 (1753)	2:262
	<b>Daucus glochidiatus</b> (Labill.) Fisch., C.A.Mey. & Avé-Lall., Index Seminum [St. Petersburg] 9, Suppl. 11 (1844)	2:262
	Daucus pusillus Michx. sensu Hooker (1860) = <b>Daucus glochidiatus</b> (misapplied in Tasmania)	
e	<b>Dichosciadium ranunculaceum</b> (F.Muell.) Domin var. <b>tasmanicum</b> (Hook.f.) Domin, Repert. Spec. Nov. Regni Veg. 5: 105 (1908)	2:250
	Didiscus humilis Hook.f. f. breviscapis Domin = <b>Trachymene humilis</b> subsp. <b>breviscapa</b>	
	Didiscus humilis Hook.f. f. humilis = <b>Trachymene humilis</b> subsp. <b>humilis</b>	
	Didiscus pilosus Hook.f. nom. illeg. = <b>Trachymene composita</b>	
e	<b>Diplaspis cordifolia</b> (Hook.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 157 (1856)	2:252
e	<b>Diplaspis hydrocotyle</b> Hook.f., London J. Bot. 6: 469 bis (1847)	2:251

	<b>Eryngium ovinum</b> A.Cunn., Geogr. Mem. New South Wales [Field]: 358 (1825)	2:252
	<i>Eryngium rostratum</i> Cav. sensu Curtis (1963) = <b>Eryngium ovinum</b> (misapplied in Tasmania)	2:252
	<b>Eryngium vesiculosum</b> Labill., Nov. Holl. Pl. 1: 73 t.98 (1805)	2:252
i	<b>Foeniculum vulgare</b> Mill., Gard. Dict., ed. 8, no.1 (1768)	2:259
	<i>Hydrocotyle asiatica</i> L. sensu Rodway (1903) = <b>Centella cordifolia</b> (misapplied in Tasmania)	
	<b>Hydrocotyle callicarpa</b> Bunge, Pl. Preiss. [J.G.C.Lehman] 1: 283 (1845)	2:246
	<b>Hydrocotyle capillaris</b> F.Muell. ex Klatt, Linnaea 29: 707 (1859)	2:247
	<b>Hydrocotyle comocarpa</b> F.Muell., Vict. Naturalist 3: 127 (1887)	
	<b>Hydrocotyle foveolata</b> H.Eichler, Suppl. Black's Fl. S. Austral., ed. 2: 248 (1965)	
	<b>Hydrocotyle hirta</b> R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 204 (1820)	2:245
	<i>Hydrocotyle javanica</i> Thunb. sensu Curtis (1963) = <b>Hydrocotyle hirta</b> (misapplied in Tasmania)	2:245
	<b>Hydrocotyle laxiflora</b> DC., Prodr. [A. P. de Candolle] 4: 61 (1830)	2:246
	<b>Hydrocotyle muscosa</b> R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 208 (1820)	2:246
	<i>Hydrocotyle peduncularis</i> R.Br. ex A.Rich. = <b>Hydrocotyle sibthorpioides</b>	
	<b>Hydrocotyle pterocarpa</b> F.Muell., Defin. Austral. Pl. 46 (1855)	2:246
	<b>Hydrocotyle sibthorpioides</b> Lam., Encycl. (Lamarck) 3: 153 (1789)	2:245
	<i>Hydrocotyle tasmanica</i> Hook.f. – a name of uncertain application	
	<b>Hydrocotyle tripartita</b> R.Br. ex A.Rich., Ann. Gen. Sci. Phys. 4: 209 (1820)	
	<i>Hydrocotyle vagans</i> Hook.f. – a name of uncertain application	
	<i>Lilaeopsis</i> sp. West Coast (A.Moscal 5655) Tas Herbarium = <b>Lilaeopsis novae-zelandiae</b>	
	<i>Lilaeopsis brownii</i> A.W.Hill = <b>Lilaeopsis polyantha</b>	2:260
	<i>Lilaeopsis gunnii</i> A.W.Hill = <b>Lilaeopsis polyantha</b>	
t	<b>Lilaeopsis novae-zelandiae</b> (Gand.) A.W.Hill, J. Linn. Soc. Bot.47: 549 (1927)	
	<b>Lilaeopsis polyantha</b> (Gand.) H.Eichler, Taxon 12: 296 (1963)	2:260
	<i>Oreomyrrhis andicola</i> (Kunth) Endl. ex Hook.f. sensu Bentham (1867) misapplied to <b>O. ciliata</b> , <b>O. eriopoda</b> & <b>O. sessiliflora</b>	
	<b>Oreomyrrhis argentea</b> (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 162 (1856)	2:257
	<i>Oreomyrrhis brachycarpa</i> (Hook.f.) Hook.f. = <b>Oreomyrrhis eriopoda</b>	
	<b>Oreomyrrhis ciliata</b> Hook.f., London J. Bot. 6: 471 bis (1847)	2:258
	<b>Oreomyrrhis eriopoda</b> (DC.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 162 (1856)	2:257
e	<b>Oreomyrrhis gunnii</b> Mathias & Constance, Univ. Calif. Publ. Bot. 27: 395 (1955)	2:258
e	<b>Oreomyrrhis sessiliflora</b> Hook.f., London J. Bot. 6: 471 bis (1847)	2:258
e	<b>Oschatzia saxifraga</b> (Hook.f.) Walp., Ann. Bot. Syst. (Walpers) 1: 340 (1848)	2:251
i	<b>Pastinaca sativa</b> L., Sp. Pl. 1: 262 (1753)	2:261
i	<b>Petroselinum crispum</b> (Mill.) Fuss, Fl. Transsilv. 254 (1866)	2:256
	<i>Scandix glochidiata</i> Labill. = <b>Daucus glochidiatus</b>	
i	<b>Scandix pecten-veneris</b> L., Sp. Pl. 1: 256 (1753)	2:259
i	<i>Torilis arvensis</i> (Huds.) Link possibly recorded in error	
i	<b>Torilis nodosa</b> (L.) Gaertn., Fruct. Sem. Pl. 1: 82, t.20 (1788)	2:262
	<i>Trachymene anisocarpa</i> (Turcz.) B.L.Burtt sensu Curtis (1963) = <b>Trachymene composita</b> (misapplied in Tasmania)	2:248
	<i>Trachymene australis</i> Benth. nom. illeg. = <b>Trachymene composita</b>	
	<b>Trachymene composita</b> (Domin) B.L.Burtt var. <b>composita</b> , J. Bot. 79: 45 (1941)	2:248
	<b>Trachymene composita</b> (Domin) B.L.Burtt var. <b>robertsonii</b> (Domin) J.M.Hart, Austral. Syst. Bot. 19: 40 (2006)	



	<b>Trachymene humilis</b> (Hook.f.) Benth. subsp. <b>breviscapa</b> (Domin) P.S.Short, Muelleria 6: 166 (1986)	2:248
	<b>Trachymene humilis</b> (Hook.f.) Benth. subsp. <b>humilis</b> , Fl. Austral. 3: 351 (1867)	2:248
	<b>Xanthosia dissecta</b> Hook.f., Icon. Pl. 4: t.302 (1841)	2:250
	Xanthosia montana Sieber ex DC. = <b>Xanthosia pilosa</b>	
	<b>Xanthosia pilosa</b> Rudge, Trans. Linn. Soc. London 10: 301 (1811)	2:249
	Xanthosia pusilla Bunge sensu Curtis (1963) = <b>Xanthosia tasmanica</b> (misapplied in Tasmania)	2:250
	<b>Xanthosia tasmanica</b> Domin, Repert. Spec. Nov. Regni Veg. 4: 298 (1907)	2:250
	<b>Xanthosia ternifolia</b> J.M.Hart & Henwood, Austral. Syst. Bot. 13: 256 (2000)	
	<b>Xanthosia tridentata</b> DC., Prodr. [A. P. de Candolle] 4: 75 (1830)	2:249
<b>APOCYNACEAE</b>		
	<b>Alyxia buxifolia</b> R.Br., Prodr. Fl. Nov. Holland. 470 (1810)	3:472
	Lyonsia straminea R.Br. sensu Rodway (1903) = <b>Parsonsia brownii</b> (misapplied in Tasmania)	
	<b>Parsonsia brownii</b> (Britten) Pichon, Notul. Syst. (Paris) 14: 10 (1950)	3:473
	Parsonsia straminea (R.Br.) F.Muell. sensu Curtis (1967) = <b>Parsonsia brownii</b> (misapplied in Tasmania)	2:473
i	<b>Vinca major</b> L., Sp. Pl. 1: 209 (1753)	3:472
<b>AQUIFOLIACEAE</b>		
i	<b>Ilex aquifolium</b> L., Sp. Pl. 1: 125 (1753)	
<b>ARALIACEAE</b>		
i	<b>Hedera helix</b> L., Sp. Pl. 1: 202 (1753)	
	Nothopanax gunnii (Hook.f.) Seem. = <b>Pseudopanax gunnii</b>	2:264
	Panax gunnii Hook.f. = <b>Pseudopanax gunnii</b>	
	Panax sambucifolius Sieber ex DC. sensu Bentham (1867) = <b>Polyscias sp. Douglas-Denison (R.Schahinger HO526133) Tas Herbarium</b>	
	Panax sambucifolius Sieber ex DC. = <b>Polyscias sambucifolia</b>	
	<b>Polyscias sp. Douglas-Denison (R.Schahinger HO526133) Tas Herbarium</b>	
	Polyscias aff. sambucifolia 'Douglas-Denison' (R.B.Schahinger HO526133) = <b>Polyscias sp. Douglas-Denison (R.Schahinger HO526133) Tas Herbarium</b>	
i	<b>Polyscias sambucifolia</b> (Sieber ex DC.) Harms, Nat. Pflanzenfam. [Engler & Prantl] III 8: 45 (1894)	
e	<b>Pseudopanax gunnii</b> (Hook.f.) K.Koch, Wochenschr. Gärtnerei Pflanzenk. 2: 366 (1859)	2:264
<b>ASCLEPIADACEAE</b>		
i	<b>Gomphocarpus fruticosus</b> (L.) W.T.Aiton subsp. <b>fruticosus</b> , Hortus Kew. (W.T.Aiton), ed. 2, 2: 80 (1811)	
<b>ASTERACEAE (COMPOSITAE)</b>		
e	<b>Abrotanella forsteroides</b> (Hook.f.) Benth., Fl. Austral. 3: 554 (1867)	2:359
e	<b>Abrotanella scapigera</b> (F.Muell.) Benth., Fl. Austral. 3: 554 (1867)	2:359
i	<b>Achillea distans</b> Waldst. & Kit. ex Willd., Sp. Pl., ed. 4 [Willdenow] 3: 2207 (1800)	2:351
i	<b>Achillea millefolium</b> L., Sp. Pl. 2: 899 (1753)	2:351
	Achillea tanacetifolia All. = <b>Achillea distans</b>	2:351
	<b>Actites megalocarpus</b> (Hook.f.) Lander, Telopea 1: 130 (1976)	2:390
	<b>Allittia cardiocarpa</b> (F.Muell. ex Benth.) P.S.Short, Muelleria 20: 55 (2004)	2:296
i	<b>Ammobium alatum</b> R.Br., Bot. Mag. 51: t.2459 (1824)	2:343
	Ammobium calyceroides (Cass.) Anderb. = <b>Nablonium calyceroides</b>	2:349

	Angianthus eriocephalus (A.Gray) Benth. = <b>Angianthus preissianus</b>	2:344
	<b>Angianthus preissianus</b> (Steetz) Benth., Fl. Austral. 3: 566 (1867)	2:344
i	<b>Anthemis arvensis</b> L., Sp. Pl. 2: 894 (1753)	2:352
i	<b>Anthemis cotula</b> L., Sp. Pl. 2: 894 (1753)	2:352
	Anthemis nobilis L. = <b>Chamaemelum nobile</b>	2:352
	Anthemis tinctoria L. = Cota tinctoria	2:352
	Apalochlamys billardierei DC. = <b>Apalochlamys spectabilis</b>	
	<b>Apalochlamys spectabilis</b> (Labill.) Steud., Nomencl. Bot. [Steudel], ed. 2, 1: 255 (1840)	2:340
i	<b>Arctium minus</b> (Hill) Bernh., Syst. Verz. (Bernhardi): 154 (1800)	2:375
i	<b>Arctotheca calendula</b> (L.) K.Lewin, J. S. African Bot. 8: 284 (1942)	2:374
i	<b>Arctotheca populifolia</b> (P.J.Bergius) Norl., Aquilo, Ser. Bot. 6: 84 (1967)	
i	<b>Arctotis stoechadifolia</b> P.J.Bergius, Descr. Pl. Cap. 324 (1767)	2:374
	<b>Argentipallium dealbatum</b> (Labill.) Paul G.Wilson, Nuytsia 8: 458 (1992)	2:330
	<b>Argentipallium obtusifolium</b> (Sond.) Paul G.Wilson, Nuytsia 8: 458 (1992)	2:329
e	<b>Argentipallium × spiceri</b> (F.Muell.) Paul G.Wilson, Nuytsia 8: 458 (1992)	2:329
	<b>Argyrotegium fordianum</b> (M.Gray) J.M.Ward & Breitw., New Zealand J. Bot. 41: 609 (2003)	
	<b>Argyrotegium mackayi</b> (Buchanan) J.M.Ward & Breitw., New Zealand J. Bot. 41: 609 (2003)	2:320
	<b>Argyrotegium nitidulum</b> (Hook.f.) J.M.Ward & Breitw., New Zealand J. Bot. 41: 609 (2003)	
	<b>Argyrotegium poliochlorum</b> (N.G.Walsh) J.M.Ward & Breitw., New Zealand J. Bot. 41: 609 (2003)	2:320
	Aster aculeatus Labill. = <b>Olearia ramulosa</b>	
	Aster argophyllus Labill. = <b>Olearia argophylla</b>	
	Aster microphyllus Labill. nom. illeg. = <b>Olearia lepidophylla</b>	
	Aster myrsinoides Labill. = <b>Olearia myrsinoides</b>	
	Aster phlogopappus Labill. = <b>Olearia phlogopappa</b>	
	Aster ramulosus Labill. = <b>Olearia ramulosa</b>	
	Aster stellulatus Labill. = <b>Olearia stellulata</b>	
	Aster subulatus Michx. = <b>Symphotrichum subulatum</b>	2:299
	Aster viscosus Labill. = <b>Olearia viscosa</b>	
	<b>Bedfordia arborescens</b> Hochr., Candollea 5: 332 (1934)	
e	<b>Bedfordia linearis</b> (Labill.) DC. subsp. <b>linearis</b> , Prodr. [A. P. de Candolle] 6: 441 (1838)	2:371
e	<b>Bedfordia linearis</b> (Labill.) DC. subsp. <b>oblongifolia</b> Orchard var. <b>curvifolia</b> Orchard, Muellera 19: 93 (2004)	2:371
e	<b>Bedfordia linearis</b> (Labill.) DC. subsp. <b>oblongifolia</b> Orchard var. <b>oblongifolia</b> , Muellera 19: 90 (2004)	2:371
e	<b>Bedfordia linearis</b> (Labill.) DC. × <b>B. salicina</b> (Labill.) DC.	
e	<b>Bedfordia salicina</b> (Labill.) DC., Prodr. [A. P. de Candolle] 6: 441 (1838)	2:371
	Bellis aculeata Labill. = <b>Brachyscome aculeata</b>	
	Bellis graminea Labill. = <b>Brachyscome graminea</b>	
i	<b>Bellis perennis</b> L., Sp. Pl. 2: 886 (1753)	2:299
	Bellis stipitata Labill. = <b>Lagenophora stipitata</b>	
i	<b>Berkheya rigida</b> (Thunb.) Bolus & Wolley-Dod, Trans. S.African. Philos. Soc. 14: 289 (1904)	
	Brachyglottis brunonis (Hook.f.) B.Nord. = <b>Centropappus brunonis</b>	2:365
	<b>Brachyscome aculeata</b> (Labill.) Less., Syn. Gen. Compos. 192 (1832)	2:296
	Brachyscome angustifolia A.Cunn. ex DC. = <b>Brachyscome graminea</b>	2:295

	Brachyscome cardiocarpa F.Muell. ex Benth. = <b>Allittia cardiocarpa</b>	2:296
	<b>Brachyscome ciliaris</b> (Labill.) Less., Syn. gen. Compos. 192 (1832)	2:298
	Brachyscome ciliaris (Labill.) Less. var. robusta Benth. = <b>Brachyscome rigidula</b>	
	<b>Brachyscome decipiens</b> Hook.f., London J. Bot. 6: 114 (1847)	2:295
	<b>Brachyscome diversifolia</b> (Graham ex Hook.) Fisch. & C.A.Mey., Index Seminum [St. Petersburg (Petropolitanus)] 2: 31 (1835)	2:297
	Brachyscome diversifolia (Graham ex Hook.) Fisch. & C.A.Mey. var. maritima Benth. = <b>Brachyscome diversifolia</b>	2:297
	<b>Brachyscome graminea</b> (Labill.) F.Muell., Fragm. (Mueller) 1(3): 49 (1858)	2:295
	Brachyscome linearifolia DC. sensu Hooker (1860) = <b>Allittia cardiocarpa</b> (misapplied in Tasmania)	
	Brachyscome nivalis F.Muell. var. alpina sensu Curtis (1963) = <b>Brachyscome radicans</b> (misapplied in Tasmania)	2:297
	<b>Brachyscome parvula</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 185 (1856)	2:298
	<b>Brachyscome perpusilla</b> (Steetz) J.M.Black, Fl. S. Austral. [J.M. Black] 4: 587 (1929)	
	Brachyscome pumila Walp. sensu Hooker (1860) – a name of uncertain application	
	<b>Brachyscome radicans</b> Steetz, Pl. Preiss. [J.G.C.Lehman] 1: 429 (1845)	2:295
t	<b>Brachyscome radicata</b> Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 127 (1852)	2:298
	<b>Brachyscome rigidula</b> (DC.) G.L.Davis, Proc. Linn. Soc. New South Wales 73: 219 (1948)	2:298
	Brachyscome scapiformis DC. = <b>Brachyscome spathulata</b>	2:296
	Brachyscome scapiformis DC. var. tenuiscapa (Hook.f.) Benth. = <b>Brachyscome tenuiscapa</b>	
	Brachyscome sieberi DC. var. gunnii DC. = <b>Brachyscome aculeata</b>	
	<b>Brachyscome spathulata</b> Gaudich., Voy. Uranie, Bot. 468 (1830)	
	Brachyscome spathulata Gaudich. subsp. glabra (DC.) H.M.Stace = <b>Brachyscome spathulata</b>	2:296
	Brachyscome stricta DC. = <b>Brachyscome aculeata</b>	2:296
	Brachyscome tadgellii Tovey & P.Morris sensu Buchanan 1995 previously recorded in error	
e	<b>Brachyscome tasmanica</b> P.S.Short, J. Adelaide Bot. Gard. 28: 160 (2014)	
e	<b>Brachyscome tenuiscapa</b> Hook.f., London J. Bot. 6: 114 (1847)	2:294
	Brachyscome tenuiscapa Hook.f. var. pubescens (Benth.) G.L.Davis sensu Buchanan (1989) = <b>Brachyscome staceae</b> (N.S.W., previously recorded in error)	2:294
	Bracteantha bicolor (Lindl.) Anderb. & Haegi = <b>Xerochrysum bicolor</b>	2:331
	Bracteantha bracteata (Vent.) Anderb. & Haegi = <b>Xerochrysum bracteatum</b>	
	Bracteantha palustris Flann = <b>Xerochrysum palustre</b>	
	Bracteantha papillosa (Labill.) Anderb. & Haegi = <b>Xerochrysum papillosum</b>	2:331
	Bracteantha subundulata (Sch.Bip.) Paul G.Wilson = <b>Xerochrysum subundulatum</b>	2:330
	Cacalia linearis Labill. = <b>Bedfordia linearis</b>	
	Cacalia salicina Labill. = <b>Bedfordia salicina</b>	
	Calea aculeata Labill. = <b>Cassinia aculeata</b>	
	Calea spectabilis Labill. = <b>Apalochlamys spectabilis</b>	
i	<b>Calendula arvensis</b> L., Sp. Pl., ed. 2, 2: 1303 (1763)	2:372
i	<b>Calendula officinalis</b> L., Sp. Pl. 2: 921 (1753)	2:373
	Calocephalus brownii (Cass.) F.Muell. = <b>Leucophyta brownii</b>	2:345
	<b>Calocephalus citreus</b> Less., Syn. Gen. Compos. 271 (1832)	2:346
	<b>Calocephalus lacteus</b> Less., Syn. Gen. Compos. 271 (1832)	2:345
	Carduus arvensis (L.) Robson = <b>Cirsium arvense</b>	

	Carduus lanceolatus L. = <b>Cirsium vulgare</b>	
	Carduus marianus L. = <b>Silybum marianum</b>	
i	<b>Carduus nutans</b> L., Sp. Pl. 2: 821 (1753)	
	Carduus pratensis Huds. sensu Rodway (1903) = <b>Cirsium vulgare</b> (misapplied in Tasmania)	
i	<b>Carduus pycnocephalus</b> L., Sp. Pl., ed. 2, 2: 1151 (1763)	
i	<b>Carduus tenuiflorus</b> Curtis, Fl. Londin. (Curtis) 2(6): t.169 (1793)	2:376
i	<b>Carthamus lanatus</b> L., Sp. Pl. 2: 830 (1753)	2:380
i	<b>Carthamus tinctorius</b> L., Sp. Pl. 2: 830 (1753)	
	<b>Cassinia aculeata</b> (Labill.) R.Br. subsp. <b>aculeata</b> , Trans. Linn. Soc. London 12: 127 (1817)	2:339
	Cassinia longifolia R.Br. sensu Curtis (1963) = <b>Cassinia trinerva</b>	2:340
	<b>Cassinia rugata</b> N.G.Walsh, Muelleria 7: 141 (1990)	
	Cassinia spectabilis (Labill.) R.Br. = <b>Apalochlamys spectabilis</b>	2:340
	<b>Cassinia trinerva</b> N.A.Wakef., Vict. Naturalist 68: 69 (1951)	2:340
e	<b>Celmisia asteliifolia</b> Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 35 (1844)	2:310
	Celmisia longifolia Cass. sensu Curtis (1963) = <b>Celmisia asteliifolia</b>	2:310
	Celmisia longifolia Cass. var. saxifraga Benth. = <b>Celmisia saxifraga</b>	
e	<b>Celmisia saxifraga</b> (Benth.) W.M.Curtis, Taxon 17: 467 (1968)	2:310
n i *	Centaurea calcitrapa L., Sp. Pl. 2: 917 (1753)	2:379
n i *	Centaurea cyanus L., Sp. Pl. 2: 911 (1753)	
	Centaurea jacea L. sensu Buchanan (2007) = <b>Centaurea × moncktonii</b>	2:378
i	<b>Centaurea melitensis</b> L., Sp. Pl. 2: 917 (1753)	2:380
i	<b>Centaurea × moncktonii</b> C.E.Britton, Rep. Bot. Soc. Exch. Club Brit. Isles 1920, 6: 172 (1921)	2:378
	Centaurea nigra L. sensu Buchanan (2007) = <b>Centaurea × moncktonii</b>	2:379
n	Centaurea solstitialis L. – previously listed as naturalised but insufficient evidence exists to support this	2:379
	<b>Centipeda cunninghamii</b> (DC.) A.Braun & Asch., Index Seminum Hort. Bot. Berol. App. 6 (1867)	
	<b>Centipeda elatinoides</b> (Less.) Benth. & Hook.f. ex O.Hoffm., Nat. Pflanzenfam. [Engler & Prantl] 4(5): 280 (1892)	2:358
	Centipeda minima (L.) A.Braun & Asch. sensu Curtis (1963) = <b>Centipeda elatinoides</b>	2:358
e	<b>Centropappus brunonis</b> Hook.f., London J. Bot. 6: 124 (1847)	2:365
i	<b>Chamaemelum nobile</b> (L.) All., Fl. Pedem. 1: 185 (1785)	2:352
	Chondrilla juncea L. sensu Curtis (1963) anecdotal, possibly recorded in error	2:388
i	<b>Chrysanthemoides monilifera</b> (L.) Norl. subsp. <b>monilifera</b> , Stud. Calendulae 1: 374 (1943)	2:373
	Chrysanthemum leucanthemum L. = <b>Leucanthemum vulgare</b>	2:353
	Chrysanthemum parthenium (L.) Bernh. = <b>Tanacetum parthenium</b>	2:353
n	<b>Chrysocephalum apiculatum</b> (Labill.) Steetz subsp. <b>apiculatum</b> , Pl. Preiss. [J.G.C.Lehman] 1: 474 (1845)	2:332
	<b>Chrysocephalum baxteri</b> (A.Cunn. ex DC.) Anderb., Compositae Newslett. 19: 22 (1991)	2:328
n	<b>Chrysocephalum semipapposum</b> (Labill.) Steetz subsp. <b>asperum</b> (Steetz) Paul G.Wilson, Nuytsia 27: 70 (2016)	
n	<b>Chrysocephalum semipapposum</b> (Labill.) Steetz subsp. <b>lineare</b> Paul G.Wilson, Nuytsia 27: 67 (2016)	
n	<b>Chrysocephalum semipapposum</b> (Labill.) Steetz subsp. <b>semipapposum</b> , Pl. Preiss. [J.G.C.Lehman] 1: 474 (1845)	2:332
	Chrysocoma cinerea Labill. = <b>Ozothamnus turbinatus</b>	

	Chrysocoma reticulata Labill. = <b>Ozothamnus reticulatus</b>	
	Chrysocoma squamata Labill. = <b>Leptorhynchus squamatus</b>	
i	<b>Cichorium intybus</b> L., Sp. Pl. 2: 813 (1753)	2:380
i	<b>Cirsium arvense</b> (L.) Scop. var. <b>arvense</b> , Fl. Carniol., ed. 2, 2: 126 (1772)	2:377
i	<b>Cirsium vulgare</b> (Savi) Ten., Fl. Napol. 5: 209 (1836)	2:376
	Cnicus arvensis (L.) Roth = <b>Cirsium arvense</b>	2:377
	Conyza albida Willd. ex Spreng. = <b>Conyza sumatrensis</b>	2:315
i	<b>Conyza bonariensis</b> (L.) Cronquist, Bull. Torrey Bot. Club 70: 632 (1943)	2:315
i	<b>Conyza canadensis</b> (L.) Cronquist, Bull. Torrey Bot. Club 70: 632 (1943)	2:314
	Conyza chilensis Spreng. = <b>Conyza primulifolia</b>	2:314
	Conyza floribunda Kunth sensu Curtis (1963) = <b>Conyza sumatrensis</b>	2:315
i x	<b>Conyza primulifolia</b> (Lam.) Cuatrec. & Lourteig, Phytologia 58: 475 (1985)	2:314
	Conyza scabiosifolia J.Rémy = <b>Conyza primulifolia</b>	2:314
i	<b>Conyza sumatrensis</b> (Retz.) E.Walker, J. Jap. Bot. 46(3): 72 (1971)	2:315
	Coronidium sp. Lowland Swamps (V.Stajsic 4226) Vic Herbarium = <b>Coronidium gunnianum</b>	
	<b>Coronidium gunnianum</b> (Hook.) N.G.Walsh, Muelleria 32: 20 (2014)	
	<b>Coronidium monticola</b> N.G.Walsh, Muelleria 32: 21 (2014)	
	<b>Coronidium scorpioides</b> (Labill.) Paul G.Wilson, Nuytsia 18: 326 (2008)	2:328
	Cota tinctoria (L.) J.Gay – formerly considered naturalised though no evidence exists to support this	2:352
	<b>Cotula alpina</b> (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 192 t.51A (1856)	2:356
	<b>Cotula australis</b> (Sieber ex Spreng.) Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) I: 128 (1852)	2:356
i	<b>Cotula coronopifolia</b> L., Sp. Pl. 2: 892 (1753)	2:356
	Cotula coronopifolia L. var. integrifolia (Hook.f.) Rodway = <b>Cotula coronopifolia</b>	
	Cotula filicula (Hook.f.) Benth. = <b>Leptinella filicula</b>	2:357
	Cotula filifolia Thunb. sensu Rodway (1903) = <b>Cotula vulgaris</b> var. <b>australasica</b> (misapplied in Tasmania)	
	Cotula integrifolia Hook.f. nom. illeg. = <b>Cotula coronopifolia</b>	
	Cotula longipes (Hook.f.) W.M.Curtis = <b>Leptinella longipes</b>	2:357
	Cotula reptans (Benth.) Benth. = <b>Leptinella reptans</b>	2:357
	Cotula reptans (Benth.) Benth. var. major Benth. = <b>Leptinella longipes</b>	
	<b>Cotula vulgaris</b> Levyns var. <b>australasica</b> J.H.Willis, Vict. Naturalist 73: 201 (1957)	2:355
	Craspedia alpina Backh. ex Hook.f. = <b>Craspedia macrocephala</b>	2:347
	Craspedia coolaminica J.Everett & Joy Thomps. = <b>Craspedia gracilis</b>	
e	<b>Craspedia cynurica</b> Rozefelds & A.M.Buchanan, Kanunnah 4: 106 (2011)	
e	<b>Craspedia glabrata</b> (Hook.f.) Rozefelds, Telopea 9: 814 (2002)	2:347
e	<b>Craspedia glauca</b> (Labill.) Spreng., Syst. Veg. (ed. 16) [Sprengel] 3: 441 (1826)	2:347
	Craspedia glauca (Labill.) Spreng. var. glabrata Hook.f. ex W.M.Curtis nom. inval. =	
	<b>Craspedia glabrata</b>	2:347
	Craspedia glauca (Labill.) Spreng. var. gracilis W.M.Curtis nom. inval. = <b>Craspedia gracilis</b>	2:347
	Craspedia glauca (Labill.) Spreng. var. macrocephala W.M.Curtis nom. inval. =	
	<b>Craspedia macrocephala</b>	2:347
	<b>Craspedia gracilis</b> Hook.f., London J. Bot. 6: 118 (1847)	
e	<b>Craspedia macrocephala</b> Hook., Bot. Mag. 62: t. 3415 (1835)	2:347
	<b>Craspedia paludicola</b> J.Everett & Doust, Telopea 5: 35 (1992)	

e	<b>Craspedia preminghana</b> Rozefelds, <i>Telopea</i> 9: 816 (2002)	
	<i>Craspedia richia</i> Cass. nom. illeg. = <b>Craspedia glauca</b>	
	<i>Craspedia richia</i> Cass. var. <i>alpina</i> (Backh. ex Hook.f.) Benth. = <b>Craspedia macrocephala</b>	
	<i>Craspedia richia</i> Cass. var. <i>glabrata</i> Hook.f. = <b>Craspedia glabrata</b>	
n	<i>Craspedia richia</i> Cass. var. <i>gracilis</i> (Hook.f.) Hook.f. = <b>Craspedia gracilis</b>	
n	<i>Craspedia richia</i> Cass. var. <i>linearis</i> Hook.f. = <b>Craspedia gracilis</b>	
	<i>Craspedia richia</i> Cass. var. <i>macrocephala</i> (Hook.) Benth. = <b>Craspedia macrocephala</b>	
e	<b>Craspedia rosulata</b> Rozefelds & A.M.Buchanan, <i>Kanunnah</i> 4: 111 (2011)	
i	<b>Crepis capillaris</b> (L.) Wallr., <i>Erst. Beitr. Fl. Hercyn.</i> 287 (1840)	2:384
	<i>Crepis nicaeensis</i> Baldinger & Pers. sensu Curtis (1963) recorded in error	2:384
i	<b>Crepis setosa</b> Haller f., <i>Arch. Bot. (Leipzig)</i> 1(2): 1 (1797)	2:384
	<i>Cryptostemma calendula</i> (L.) R.Br. = <b>Arctotheca calendula</b>	
	<i>Cymbonotus lawsonianus</i> Gaudich. sensu Curtis (1963) = <b>Cymbonotus preissianus</b>	2:373
	<b>Cymbonotus preissianus</b> Steetz, <i>Pl. Preiss</i> 1: 486 (1845)	2:373
n i x	<b>Cynara cardunculus</b> L. subsp. <b>flavescens</b> Wiklund, <i>Bot. J. Linn. Soc.</i> 109: 120 (1992)	
i	<b>Delairea odorata</b> Lem., <i>Ann. Sci. Nat., Bot., sér. 3, 1:</i> 380 (1844)	2:367
i	<b>Dimorphotheca fruticosa</b> (L.) DC., <i>Prodr. [A. P. de Candolle]</i> 6: 71 (1838)	
i	<b>Dittrichia graveolens</b> (L.) Greuter, <i>Exsicc. Genav. Conserv. Bot. Distrib. Fasc. 4:</i> 71 (1973)	2:348
	<i>Erechtites argutus</i> (A.Rich.) DC. = <b>Senecio glomeratus</b> subsp. <b>glomeratus</b>	
	<i>Erechtites gunnii</i> Hook.f. = <b>Senecio gunnii</b>	
	<i>Erechtites hispidula</i> (A.Rich.) DC. = <b>Senecio hispidulus</b>	
	<i>Erechtites prenathoides</i> (A.Rich.) DC. = <b>Senecio prenanthoides</b>	
	<i>Erechtites quadridentata</i> (Labill.) DC. = <b>Senecio quadridentatus</b>	
	<i>Erechtites quadridentata</i> (Labill.) DC. var. <i>glabrescens</i> (DC.) Benth. sensu Bentham (1867) = <b>Senecio quadridentatus</b> (misapplied in Tasmania)	
	<i>Erechtites quadridentata</i> (Labill.) DC. var. <i>gunnii</i> (Hook.f.) Benth. = <b>Senecio gunnii</b>	
	<i>Erigeron bellidioides</i> (Hook.f.) S.J.Forbes & D.I.Morris = <b>Pappochroma bellidioides</b>	
	<i>Erigeron gunnii</i> (Hook.f.) F.Muell. ex Hook.f. = <b>Pappochroma gunnii</b>	
	<i>Erigeron gunnii</i> (Hook.f.) F.Muell. ex Hook.f. var. <i>bellidioides</i> (Hook.f.) Hook.f. = <b>Pappochroma bellidioides</b>	
i	<b>Erigeron karvinskianus</b> DC., <i>Prodr. [A. P. de Candolle]</i> 5: 285 (1836)	2:312
	<i>Erigeron linifolius</i> Willd. = <b>Conyza bonariensis</b>	
	<i>Erigeron mucronatus</i> DC. = <b>Erigeron karvinskianus</b>	2:312
	<i>Erigeron pappocromus</i> Labill. = <b>Pappochroma pappocromum</b>	2:311
	<i>Erigeron pappocromus</i> Labill. var. <i>billardieri</i> Benth. = <b>Pappochroma pappocromum</b>	2:311
	<i>Erigeron pappocromus</i> Labill. var. <i>gunnii</i> Benth. = <b>Pappochroma gunnii</b>	2:312
	<i>Erigeron pappocromus</i> Labill. var. <i>oblongatus</i> Benth. = <b>Pappochroma tasmanicum</b>	
	<i>Erigeron pappocromus</i> Labill. var. <i>stellatus</i> (Hook.f.) Benth. = <b>Pappochroma stellatum</b>	
	<i>Erigeron stellatus</i> (Hook.f.) W.M.Curtis = <b>Pappochroma stellatum</b>	2:312
	<i>Erigeron tasmanicus</i> (Hook.f.) Hook.f. = <b>Pappochroma tasmanicum</b>	2:312
	<i>Erigeron trigonus</i> S.J.Forbes & D.I.Morris = <b>Pappochroma trigonum</b>	
	<i>Euchiton argentifolius</i> (N.A.Wakef.) Anderb. = <b>Argyrotegium mackayi</b>	2:320
	<i>Euchiton collinus</i> Cass. = <b>Euchiton japonicus</b>	2:319
	<i>Euchiton fordianus</i> (M.Gray) P.S.Short = <b>Argyrotegium fordianum</b>	
	<b>Euchiton involucratus</b> (G.Forst.) Holub, <i>Folia Geobot. Phytotax.</i> 9: 271 (1974)	2:318

	<b>Euchiton japonicus</b> (Thunb.) Holub, Folia Geobot. Phytotax. 9: 271 (1974)	2:319
t	<b>Euchiton lateralis</b> (C.J.Webb) Breiwt. & J.M.Ward, New Zealand J. Bot. 36: 303 (1998)	
	<b>Euchiton limosus</b> (D.G.Drury) Holub, Folia Geobot. Phytotax. 9: 271 (1974)	
e	<b>Euchiton litticola</b> A.M.Buchanan, Pap. & Proc. Roy. Soc. Tasmania 133: 115 (1999)	
	<i>Euchiton poliochlorus</i> N.G.Walsh = <b>Argyrotegium poliochlorum</b>	2:320
	<b>Euchiton sphaericus</b> (Willd.) Holub, Folia Geobot. Phytotax. 9: 271 (1974)	
	<b>Euchiton traversii</b> (Hook.f.) Holub, Folia Geobot. Phytotax. 9: 271 (1974)	2:320
	<b>Euchiton umbricola</b> (J.H.Willis) Anderb., Opera Bot. 104: 167 (1991)	2:319
	<i>Eupatorium ferrugineum</i> Labill. = <b>Ozothamnus ferrugineus</b>	
	<i>Eupatorium rosmarinifolium</i> Labill. = <b>Ozothamnus rosmarinifolius</b>	
	<i>Eurybia alpina</i> Hook.f. = <b>Olearia tasmanica</b>	
	<i>Eurybia argophylla</i> (Labill.) Cass. = <b>Olearia argophylla</b>	
	<i>Eurybia ciliata</i> Benth. = <b>Olearia ciliata</b>	
	<i>Eurybia ericoides</i> Steetz = <b>Olearia ericoides</b>	
	<i>Eurybia erubescens</i> Sieber ex DC. = <b>Olearia erubescens</b>	
	<i>Eurybia floribunda</i> Hook.f. = <b>Olearia floribunda</b>	
	<i>Eurybia fulvida</i> Cass. nom. illeg. = <b>Olearia stellulata</b>	
	<i>Eurybia glandulosa</i> (Labill.) DC. = <b>Olearia glandulosa</b>	
	<i>Eurybia gunniana</i> DC. = <b>Olearia phlogopappa</b> subsp. <b>gunniana</b>	
	<i>Eurybia ledifolia</i> DC. = <b>Olearia ledifolia</b>	
	<i>Eurybia linearifolia</i> DC. = <b>Olearia axillaris</b>	
	<i>Eurybia linifolia</i> Hook.f. = <b>Olearia glutinosa</b>	
	<i>Eurybia lirata</i> (Sims) DC. = <b>Olearia lirata</b>	
	<i>Eurybia myrsinoides</i> (Labill.) Nees = <b>Olearia myrsinoides</b>	
	<i>Eurybia obcordata</i> Hook.f. = <b>Olearia obcordata</b>	
	<i>Eurybia persoonioides</i> DC. = <b>Olearia persoonioides</b>	
	<i>Eurybia pinifolia</i> Hook.f. = <b>Olearia pinifolia</b>	
	<i>Eurybia ramulosa</i> (Labill.) DC. = <b>Olearia ramulosa</b>	
	<i>Eurybia viscosa</i> (Labill.) Cass. = <b>Olearia viscosa</b>	
i	<b>Euryops abrotanifolius</b> (L.) DC., Prodr. [A. P. de Candolle] 6: 443 (1838)	2:372
e	<b>Ewartia catipes</b> (DC.) Beauverd, Bull. Soc. Bot. Genève, Ser. 2, 2: 238 (1910)	2:322
e	<b>Ewartia meredithiae</b> (F.Muell.) Beauverd, Bull. Soc. Bot. Genève, Ser. 2, 2: 240 (1910)	2:322
e	<b>Ewartia planchonii</b> (Hook.f.) Beauverd, Bull. Soc. Bot. Genève, Ser. 2, 3: 253 (1911)	2:321
	<i>Felicia erigeroides</i> DC. sensu Raphael (1955) = <b>Erigeron karvinskianus</b>	
	<i>Filago gallica</i> L. = <b>Logfia gallica</b>	2:315
i	<b>Galinsoga parviflora</b> Cav., Icon. [Cavanilles] 3: 41 t.281 (1795)	2:349
i	<b>Gamochaeta calviceps</b> (Fernald) Cabrera, Bol. Soc. Argent. Bot. 9: 368 (1961)	
i	<b>Gamochaeta purpurea</b> (L.) Cabrera, Bol. Soc. Argent. Bot. 9: 377 (1961)	2:321
i	<b>Gazania linearis</b> (Thunb.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 624 (1917)	
	<i>Gnaphalium alpigenum</i> F.Muell. ex Hook.f. nom. illeg. = <b>Euchiton umbricola</b>	
	<i>Gnaphalium apiculatum</i> Labill. = <b>Chrysocephalum apiculatum</b>	
	<i>Gnaphalium argentifolium</i> N.A.Wakef. = <b>Argyrotegium mackayi</b>	2:320
	<i>Gnaphalium candidissimum</i> Lam. = <b>Vellereophyton dealbatum</b>	2:318
	<i>Gnaphalium collinum</i> Labill. var. <i>collinum</i> = <b>Euchiton japonicus</b>	2:319
	<i>Gnaphalium collinum</i> Labill. var. <i>monocephalum</i> Hook.f. = <b>Euchiton lateralis</b>	2:319

Gnaphalium fordianum M.Gray = <b>Argyrotegium fordianum</b>	
Gnaphalium indicum L. recorded in error	
<b>Gnaphalium indutum</b> Hook.f. subsp. <b>indutum</b> , London J. Bot. 6: 121 (1847)	2:320
Gnaphalium involucratum G.Forst. = <b>Euchiton involucratus</b>	2:318
Gnaphalium japonicum Thunb. = <b>Euchiton japonicus</b>	
Gnaphalium japonicum Thunb. var. <b>radicans</b> = <b>Argyrotegium mackayi</b>	
Gnaphalium luteo-album L. = <b>Helichrysum luteoalbum</b>	2:318
Gnaphalium planchonii Hook.f. = <b>Ewartia planchonii</b>	
Gnaphalium polycaulon Pers. sensu Buchanan (2009) recorded in error	
Gnaphalium purpureum L. = <b>Gamochoeta purpurea</b>	2:321
Gnaphalium semipapposum Labill. = <b>Chrysocephalum semipapposum</b>	
Gnaphalium supinum L. sensu Curtis (1963) = <b>Argyrotegium poliochlorum</b>	2:320
Gnaphalium traversii Hook.f. = <b>Euchiton traversii</b>	2:320
Gnaphalium umbricola J.H.Willis = <b>Euchiton umbricola</b>	2:319
Hedypnois cretica (L.) Dum.Cours. = <b>Leontodon rhagadioloides</b>	2:382
Hedypnois rhagadioloides (L.) F.W.Schmidt = <b>Leontodon rhagadioloides</b>	2:382
Helichrysum acuminatum DC. = <b>Xerochrysum subundulatum</b>	2:330
Helichrysum adenophorum F.Muell. recorded in error	
Helichrysum antennarium (DC.) F.Muell. ex Benth. = <b>Ozothamnus antennaria</b>	2:336
Helichrysum apiculatum (Labill.) D.Don = <b>Chrysocephalum apiculatum</b>	2:332
Helichrysum argophyllum (A.Cunn. ex DC.) N.A.Wakef. = <b>Ozothamnus argophyllum</b>	2:335
Helichrysum baccharioides F.Muell. nom. illeg. = <b>Ozothamnus hookeri</b>	
Helichrysum backhousei (Hook.f.) F.Muell. ex Benth. = <b>Ozothamnus rodwayi</b>	2:337
Helichrysum backhousei (Hook.f.) F.Muell. ex Benth. var. <b>backhousii</b> = <b>Ozothamnus rodwayi</b> var. <b>rodwayi</b>	2:337
Helichrysum backhousei (Hook.f.) F.Muell. ex Benth. var. <b>kingii</b> W.M.Curtis = <b>Ozothamnus rodwayi</b> var. <b>kingii</b>	2:337
Helichrysum backhousei (Hook.f.) F.Muell. ex Benth. var. <b>oreophilum</b> W.M.Curtis = <b>Ozothamnus rodwayi</b> var. <b>oreophilus</b>	2:337
Helichrysum baxteri A.Cunn. ex DC. = <b>Chrysocephalum baxteri</b>	2:328
Helichrysum bicolor Lindl. = <b>Xerochrysum bicolor</b>	2:331
Helichrysum bracteatum (Vent.) Andrews = <b>Xerochrysum bracteatum</b>	
Helichrysum bracteatum (Vent.) Andrews var. <b>albidum</b> DC. = <b>Xerochrysum papillosum</b> (Tasmanian material)	
Helichrysum bracteolatum (Hook.f.) Benth. = <b>Ozothamnus bracteolatus</b>	
Helichrysum cinereum (Labill.) Benth. = <b>Ozothamnus turbinatus</b>	
Helichrysum costatifructum R.V.Sm. = <b>Ozothamnus costatifructus</b>	2:332
Helichrysum dealbatum Labill. = <b>Argentipallium dealbatum</b>	2:330
Helichrysum dendroideum N.A.Wakef. = <b>Ozothamnus ferrugineus</b>	2:335
Helichrysum ericeteum W.M.Curtis = <b>Ozothamnus ericifolius</b>	2:334
Helichrysum expansifolium (P.Morris & J.H.Willis) N.T.Burb. = <b>Ozothamnus expansifolius</b>	2:337
Helichrysum gunnii (Hook.f.) Benth. = <b>Ozothamnus bracteolatus</b>	2:333
Helichrysum hookeri (Sond.) Druce = <b>Ozothamnus hookeri</b>	2:337
Helichrysum ledifolium (A.Cunn. ex DC.) Benth. = <b>Ozothamnus ledifolius</b>	2:334
<b>Helichrysum leucopsideum</b> DC., Prodr. [A. P. de Candolle] 6: 193 (1838)	2:331



	<b>Helichrysum luteoalbum</b> (L.) Rchb., Handb. Gewächsk., ed. 2, 2: 1460 (1829)	2:318
	Helichrysum lycopodioides (Hook.f.) Benth. = <b>Ozothamnus lycopodioides</b>	2:338
n	Helichrysum milliganii Hook.f. = <b>Xerochrysum milliganii</b>	2:330
	Helichrysum obcordatum (DC.) Benth. = <b>Ozothamnus obcordatus</b>	2:336
	Helichrysum obtusifolium F.Muell. & Sond. ex Sond. = <b>Argentipallium obtusifolium</b>	2:329
	Helichrysum papillosum Labill. = <b>Xerochrysum papillosum</b>	2:331
	Helichrysum paraliium (N.T.Burb.) W.M.Curtis = <b>Ozothamnus turbinatus</b>	2:333
	Helichrysum pleurandroides F.Muell. sensu Buchanan et al. (1989) = <b>Ozothamnus ericifolius</b>	2:334
e	<b>Helichrysum pumilum</b> Hook.f. var. <b>pumilum</b> , Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 213 (1856)	2:330
e	<b>Helichrysum pumilum</b> Hook.f. var. <b>spathulatum</b> A.M.Buchanan, Pap. & Proc. Roy. Soc. Tasmania 121: 54 (1987)	2:330
	Helichrysum purpurascens (DC.) W.M.Curtis = <b>Ozothamnus purpurascens</b>	2:334
	Helichrysum reticulatum (Labill.) Benth. = <b>Ozothamnus reticulatus</b>	2:332
	Helichrysum rosmarinifolium (Labill.) Benth. = <b>Ozothamnus rosmarinifolius</b>	2:335
	Helichrysum rutidolepis DC. sensu Buchanan et al (1989) = <b>Coronidium gunnium</b> & <b>C. monticola</b>	
	Helichrysum scorpioides Labill. = <b>Coronidium scorpioides</b>	2:328
	Helichrysum scutellifolium (Hook.f.) F.Muell. = <b>Ozothamnus scutellifolius</b>	2:338
	Helichrysum selaginoides (Sond. & F.Muell.) Benth. = <b>Ozothamnus selaginoides</b>	2:339
	Helichrysum semipapposum (Labill.) DC. = <b>Chrysocephalum semipapposum</b>	2:332
	Helichrysum spiceri F.Muell. = <b>Argentipallium spiceri</b>	2:329
	Helichrysum thyrsoideum (DC.) P.Morris & J.H.Willis = <b>Ozothamnus thyrsoideus</b>	2:335
	Helipterum albicans (A.Cunn.) DC. var. <b>incanum</b> (Hook.) Paul G.Wilson = <b>Leucochrysum albicans</b> var. <b>tricolor</b>	2:324
	Helipterum anthemoides (Sieber ex Spreng.) DC. = <b>Rhodanthe anthemoides</b>	2:324
	Helipterum australe (A.Gray) Druce = <b>Triptilodiscus pygmaeus</b>	
	Helipterum demissum (A.Gray) Druce = <b>Hyalosperma demissum</b>	2:325
	Helipterum exiguum F.Muell. = <b>Hyalosperma demissum</b>	
	Helipterum incanum DC. nom. illeg., nom. superfl. = <b>Leucochrysum albicans</b> var. <b>tricolor</b>	
	Helminthia echioides (L.) Gaertn. = <b>Helminthotheca echioides</b>	
i	<b>Helminthotheca echioides</b> (L.) Holub, Folia Geobot. Phytotax. 8: 176 (1973)	2:383
	Hieracium aurantiacum L. subsp. <b>carpathicola</b> Nägeli & Peter = <b>Pilosella aurantiaca</b> subsp. <b>aurantiaca</b>	2:385
	Hieracium brunneocroceum Pugsley = <b>Pilosella aurantiaca</b> subsp. <b>aurantiaca</b>	2:385
	Hieracium pilosella L. = <b>Pilosella officinarum</b> subsp. <b>officinarum</b>	
	<b>Hyalosperma demissum</b> (A.Gray) Paul G.Wilson, Nuytsia 7: 85 (1989)	2:325
i	<b>Hypochoeris glabra</b> L., Sp. Pl. 2: 811 (1753)	2:386
i	<b>Hypochoeris radicata</b> L., Sp. Pl. 2: 811 (1753)	2:385
	Inula graveolens (L.) Desf. = <b>Dittrichia graveolens</b>	2:348
	<b>Isoetopsis graminifolia</b> Turcz., Bull. Soc. Imp. Naturalistes Moscou 24: 175 t.3 (1851)	2:360
	Ixiolaena supina F.Muell. = <b>Leiocarpa supina</b>	2:341
	Ixodia achlaena D.I.Morris = <b>Odixia achlaena</b>	
	Ixodia angusta (N.A.Wakef.) N.T.Burb. = <b>Odixia angusta</b>	2:343
i	<b>Lactuca saligna</b> L., Sp. Pl. 2: 796 (1753)	2:388

n i *	<i>Lactuca serriola</i> L. f. <i>integrifolia</i> (Gray) S.D.Prince & R.N.Carter, Cent. Pl. 2: 29 (1756)	2:389
i	<b>Lactuca serriola</b> L. f. <i>serriola</i> , Cent. Pl. 2: 29 (1756)	
	<i>Lagenophora billardierei</i> Cass. = <b>Lagenophora stipitata</b>	
	<i>Lagenophora emphysopus</i> Hook.f. nom. illeg. = <b>Solenogyne gunnii</b>	
	<b>Lagenophora gracilis</b> Steetz, Pl. Preiss. [J.G.C.Lehman] 1: 431 (1845)	
	<i>Lagenophora gunniana</i> Steetz = <b>Lagenophora huegelii</b>	
	<b>Lagenophora huegelii</b> Benth., Enum. Pl. [Endlicher]: 59 (1837)	2:290
	<i>Lagenophora latifolia</i> Hook.f. = <b>Lagenophora stipitata</b>	
	<b>Lagenophora montana</b> Hook.f., London J. Bot. 6: 113 (1847)	
	<b>Lagenophora stipitata</b> (Labill.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 630 (1917)	2:290
i	<b>Lapsana communis</b> L. subsp. <i>communis</i> , Sp. Pl. 2: 811 (1753)	
i t	<b>Lasiospermum bipinnatum</b> (Thunb.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 631 (1917)	2:350
	<b>Leiocarpa supina</b> (F.Muell.) Paul G.Wilson, Nuytsia 13: 604 (2001)	2:341
	<i>Leontodon hirtus</i> L. sensu Rodway (1903) = <b>Leontodon saxatilis</b> (misapplied in Tasmania)	
	<i>Leontodon leysseri</i> (Wallr.) Beck sensu Curtis (1963) = <b>Leontodon saxatilis</b> (misapplied in Tasmania)	2:386
i	<b>Leontodon rhagadioloides</b> (L.) Enke & Zidorn, Organisms Diversity Evol. 12: 14 (2012)	2:382
i	<b>Leontodon saxatilis</b> Lam., Fl. Franç. (Lamarck) 2: 115 (1779) (as "1778")	2:386
	<i>Leontodon taraxacoides</i> (Vill.) Mérat = <b>Leontodon saxatilis</b>	2:386
	<b>Leptinella filicula</b> (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 194 (1856)	2:357
	<i>Leptinella intricata</i> Hook.f. = <b>Leptinella reptans</b>	
	<b>Leptinella longipes</b> Hook.f., London J. Bot. 6: 117 (1847)	2:357
	<b>Leptinella reptans</b> (Benth.) D.G.Lloyd & C.J.Webb, New Zealand J. Bot. 25: 103 (1987)	2:357
	<b>Leptorhynchos elongatus</b> DC., Prodr. [A. P. de Candolle] 6: 160 (1838)	2:323
	<i>Leptorhynchos linearis</i> Less. sensu Curtis (1963) = <b>Leptorhynchos nitidulus</b> (misapplied in Tasmania)	2:323
	<b>Leptorhynchos nitidulus</b> DC., Prodr. [A. P. de Candolle] 6: 160 (1838)	2:323
	<b>Leptorhynchos squamatus</b> (Labill.) Less. subsp. <i>alpinus</i> Flann, Austral. Syst. Bot. 15: 217 (2002)	2:323
	<b>Leptorhynchos squamatus</b> (Labill.) Less. subsp. <i>squamatus</i> , Syn. Gen. Compos. 273 (1832)	2:323
i	<b>Leucanthemum × superbum</b> (Bergmans ex J.W.Ingram) D.H.Kent, Watsonia 18: 89 (1990)	
i	<b>Leucanthemum vulgare</b> Lam., Fl. Franç. (Lamarck) 2: 137 (1779) (as "1778")	2:353
n	<b>Leucochrysum albicans</b> (A.Cunn.) Paul G.Wilson subsp. <i>tricolor</i> (DC.) N.G.Walsh, Muelleria 34: 13 (2015)	2:324
	<b>Leucophyta brownii</b> Cass., Dict. Sci. Nat., ed. 2 [F.Cuvier] 26: 159 (1823)	2:345
i	<b>Logfia gallica</b> (L.) Coss. & Germ., Ann. Sci. Nat., Bot., sér. 2, 20: 291 (1843)	2:315
n i *	<i>Matricaria chamomilla</i> L., Sp. Pl. 2: 891 (1753)	
	<i>Matricaria chamomilla</i> L. sensu de Salas & Baker 2014 = <b>Matricaria discoidea</b> (misapplied in Tasmania)	
i	<b>Matricaria discoidea</b> DC., Prodr. [A. P. de Candolle] 6: 50 (1838)	2:354
	<i>Matricaria matricarioides</i> (Less.) Porter sensu Curtis (1963) = <b>Matricaria discoidea</b>	2:354
	<i>Matricaria perforata</i> Mérat = <b>Tripleurospermum maritimum</b> subsp. <i>inodorum</i>	
	<i>Matricaria recutita</i> L. sensu de Salas & Baker 2014 = <b>Matricaria chamomilla</b> (misapplied in Tasmania)	

	Microseris forsteri Hook.f. nom. illeg. = <b>Microseris lanceolata</b>	
	<b>Microseris lanceolata</b> (Walp.) Sch.Bip., Jahresber. Pollichia 22-24: 310 (1866)	2:381
	Microseris scapigera (Sol. ex A.Cunn.) Sch.Bip. sensu Curtis (1963) = <b>Microseris lanceolata</b> (misapplied in Tasmania)	2:381
n	<b>Microseris walteri</b> Gand., Bull. Soc. Bot. France 65: 52 (1918)	
	<b>Millotia muelleri</b> (Sond.) P.S.Short, Muelleria 7: 246 (1990)	
	<b>Millotia tenuifolia</b> Cass. var. <b>tenuifolia</b> , Ann. Sci. Nat. (Paris) 17: 417 (1829)	2:342
	Myriogyne minuta (G.Forst.) Less. sensu Bentham (1867) = <b>Centipeda elatinoides</b> (misapplied in Tasmania)	
e	<b>Nablonium calyceroides</b> Cass., Dict. Sci. Nat., ed. 2 [F.Cuvier] 34: 101 (1825)	2:349
e	<b>Odidia achlaena</b> (D.I.Morris) Orchard, Brunonia 4: 194 (1982)	
e	<b>Odidia angusta</b> (N.A.Wakef.) Orchard, Brunonia 4: 194 (1982)	2:343
	<b>Olearia algida</b> N.A.Wakef., Vict. Naturalist 73: 97 (1956)	2:308
	Olearia alpina (Hook.f.) W.M.Curtis nom. illeg. = <b>Olearia tasmanica</b>	2:304
e	<b>Olearia archeri</b> Lander, Muelleria 7: 117 (1989)	
	<b>Olearia argophylla</b> (Labill.) F.Muell. ex Benth., Fl. Austral. 3: 470 (1867)	2:303
	<b>Olearia axillaris</b> (DC.) F.Muell. ex Benth., Fl. Austral. 3: 475 (1867)	2:307
	<b>Olearia ciliata</b> (Benth.) F.Muell. ex Benth., Fl. Austral. 3: 488 (1867)	2:309
e	<b>Olearia ericoides</b> (Steetz) N.A.Wakef., Vict. Naturalist. 73: 97 (1956)	2:307
	<b>Olearia erubescens</b> (DC.) Dippel, Handb. Laubholz 1: 290 (1889)	2:303
	<b>Olearia floribunda</b> (Hook.f.) Benth., Fl. Austral. 3: 477 (1867)	2:308
	<b>Olearia glandulosa</b> (Labill.) Benth., Fl. Austral. 3: 483 (1867)	2:309
	<b>Olearia glutinosa</b> (Lindl.) Benth., Fl. Austral. 3: 482 (1867)	2:309
	Olearia gunniana (DC.) Hook.f. ex Hook. = <b>Olearia phlogopappa</b> subsp. <b>gunniana</b>	
e	<b>Olearia hookeri</b> (Sond.) Benth., Fl. Austral. 3: 483 (1867)	2:309
	Olearia lanceolata (Benth.) D.I.Morris sensu Morris (1977) = <b>Olearia archeri</b> (misapplied in Tasmania)	
e	<b>Olearia ledifolia</b> (DC.) Benth., Fl. Austral. 3: 472 (1867)	2:305
	<b>Olearia lepidophylla</b> (Pers.) Benth., Fl. Austral. 3: 477 (1867)	2:308
	<b>Olearia lirata</b> (Sims) Hutch., Gard. Chron., Ser. 3, 61: 14 fig.4 (1917)	2:306
	<b>Olearia myrsinoides</b> (Labill.) F.Muell. ex Benth., Fl. Austral. 3: 470 (1867)	2:303
	Olearia myrsinoides (Labill.) F.Muell. ex Benth. var. erubescens (DC.) Benth. = <b>Olearia erubescens</b>	
e	<b>Olearia obcordata</b> (Hook.f.) Benth., Fl. Austral. 3: 471 (1867)	2:304
e	<b>Olearia persoonioides</b> (DC.) Benth., Fl. Austral. 3: 471 (1867)	2:304
	Olearia persoonioides (DC.) Benth. var. alpina (Hook.f.) Benth. = <b>Olearia tasmanica</b>	
	Olearia persoonioides (DC.) Benth. var. lanceolata Benth. sensu Morris (1977) = <b>Olearia archeri</b>	
e	<b>Olearia phlogopappa</b> (Labill.) DC. subsp. <b>angustifolia</b> (Hook.f.) Messina, Austral. Syst. Bot. 26: 57 (2013)	2:306
e	<b>Olearia phlogopappa</b> (Labill.) DC. subsp. <b>gunniana</b> (DC.) Messina, Austral. Syst. Bot. 26: 60 (2013)	2:306
	<b>Olearia phlogopappa</b> (Labill.) DC. subsp. <b>insularis</b> Messina, Austral. Syst. Bot. 26: 63 (2013)	
e	<b>Olearia phlogopappa</b> (Labill.) DC. subsp. <b>phlogopappa</b> , Prodr. [A. P. de Candolle] 5: 272 (1836)	2:306

	<b>Olearia phlogopappa</b> (Labill.) DC. subsp. <b>salicina</b> (Hook.f.) Messina, Austral. Syst. Bot. 26: 63 (2013)	2:306
e	<b>Olearia phlogopappa</b> (Labill.) DC. subsp. <b>subrepanda</b> (DC.) Messina, Austral. Syst. Bot. 26: 67 (2013)	2:306
	<i>Olearia phlogopappa</i> (Labill.) DC. var. <i>angustifolia</i> (Hutch.) W.M.Curtis = <b>Olearia phlogopappa</b> subsp. <b>angustifolia</b>	2:306
	<i>Olearia phlogopappa</i> (Labill.) DC. var. <i>brevipes</i> (Hutch.) W.M.Curtis = <b>Olearia phlogopappa</b> subsp. <b>gunniana</b>	2:306
	<i>Olearia phlogopappa</i> (Labill.) DC. var. <i>microcephala</i> (Hutch.) W.M.Curtis = <b>Olearia phlogopappa</b> subsp. <b>gunniana</b>	2:306
	<i>Olearia phlogopappa</i> (Labill.) DC. var. <i>salicifolia</i> (Hutch.) W.M.Curtis = <b>Olearia phlogopappa</b> subsp. <b>salicina</b>	2:306
	<i>Olearia phlogopappa</i> (Labill.) DC. var. <i>subrepanda</i> (DC.) J.H.Willis = <b>Olearia phlogopappa</b> subsp. <b>subrepanda</b>	2:306
e	<b>Olearia pinifolia</b> (Hook.f.) Benth., Fl. Austral. 3: 471 (1867)	2:305
	<b>Olearia ramulosa</b> (Labill.) Benth., Fl. Austral. 3: 476 (1867)	2:307
e	<b>Olearia rugosa</b> (F.Muell. ex W.Archer bis) Hutch. subsp. <b>intermedia</b> Messina, Austral. Syst. Bot. 26: 76 (2013)	
	<b>Olearia stellulata</b> (Labill.) DC., Prodr. [A. P. de Candolle] 5: 272 (1836)	2:305
	<i>Olearia stellulata</i> (Labill.) DC. var. <i>lirata</i> (Sims) Benth. = <b>Olearia lirata</b>	
	<i>Olearia stellulata</i> (Labill.) DC. var. <i>quercifolia</i> (Sieber ex DC.) Benth. sensu Bentham (1867) = <b>Olearia rugosa</b> subsp. <b>intermedia</b> (misapplied in Tasmania)	
e	<b>Olearia tasmanica</b> W.M.Curtis, Vict. Naturalist 87: 251 (1970)	2:304
	<b>Olearia viscosa</b> (Labill.) Benth., Fl. Austral. 3: 468 (1867)	2:302
i	<b>Onopordum acanthium</b> L., Sp. Pl. 2: 827 (1753)	2:377
n i x	<b>Onopordum acaulon</b> L., Sp. Pl., ed. 2, 2: 1159 (1763)	
	<i>Osteospermum fruticosum</i> (L.) Norl. = <b>Dimorphotheca fruticosa</b>	
e	<b>Ozothamnus antennaria</b> (DC.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 203 (1856)	2:336
	<b>Ozothamnus argophyllus</b> (A.Cunn. ex DC.) Anderb., Opera Bot. 104: 89 (1991)	2:335
	<i>Ozothamnus backhousei</i> Hook.f. nom. illeg. = <b>Ozothamnus rodwayi</b>	
e	<b>Ozothamnus bracteolatus</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 203 (1856)	2:333
	<i>Ozothamnus cinereus</i> (Labill.) Sw. = <b>Ozothamnus turbinatus</b>	2:333
	<i>Ozothamnus cinereus</i> (Labill.) Sw. var. <i>bracteolatus</i> (Hook.f.) Rodway = <b>Ozothamnus bracteolatus</b>	
e	<b>Ozothamnus costatifructus</b> (R.V.Sm.) Anderb., Opera Bot. 104: 89 (1991)	2:332
e	<b>Ozothamnus ericifolius</b> Hook.f., London J. Bot. 6: 119 (1847)	2:334
e	<b>Ozothamnus × expansifolius</b> (P.Morris & J.H.Willis) Anderb., Opera Bot. 104: 89 (1991)	2:337
	<b>Ozothamnus ferrugineus</b> (Labill.) Sweet, Hort. Brit. [Sweet] I: 221 (1826)	2:335
	<i>Ozothamnus ferrugineus</i> DC. var. <i>gravesii</i> Rodway = <b>Ozothamnus argophyllus</b>	
	<i>Ozothamnus gunnii</i> Hook.f. = <b>Ozothamnus bracteolatus</b>	2:333
e	<b>Ozothamnus hookeri</b> Sond., Linnaea 25: 509 (1853)	2:337
e	<b>Ozothamnus ledifolius</b> (A.Cunn. ex DC.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 204 (1856)	2:334
e	<b>Ozothamnus lycopodioides</b> Hook.f., London J. Bot. 6: 119 (1847)	2:338
	<b>Ozothamnus obcordatus</b> DC., Prodr. [A. P. de Candolle] 6: 165 (1838)	2:336

e	<b>Ozothamnus purpurascens</b> DC., Prodr. [A. P. de Candolle] 6: 165 (1838)	2:334
e	<b>Ozothamnus reflexifolius</b> Leeson & Rozefelds, Austral Syst. Bot. 16: 319 (2003)	
e	<b>Ozothamnus reticulatus</b> (Labill.) DC., Prodr. [A. P. de Candolle] 6: 164 (1838)	2:332
e	<b>Ozothamnus rodwayi</b> Orchard var. <b>kingii</b> (W.M.Curtis) P.S.Short, Muelleria 7: 522 (1992)	2:337
e	<b>Ozothamnus rodwayi</b> Orchard var. <b>oreophilus</b> (W.M.Curtis) P.S.Short, Muelleria 7: 522 (1992)	2:337
e	<b>Ozothamnus rodwayi</b> Orchard var. <b>rodwayi</b> , Muelleria 7: 522 (1992)	2:337
	<b>Ozothamnus rosmarinifolius</b> (Labill.) Sweet, Hort. Brit. [Sweet] 1: 221 (1826)	2:335
	<i>Ozothamnus rosmarinifolius</i> (Labill.) Sweet var. <i>ericifolius</i> (Hook.f.) Rodway = <b>Ozothamnus ericifolius</b>	
e	<b>Ozothamnus scutellifolius</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 202 (1856)	2:338
e x	<b>Ozothamnus selaginoides</b> Sond. & F.Muell., Linnaea 25: 510 (1853)	2:339
	<b>Ozothamnus thyrsoides</b> DC., Prodr. [A. P. de Candolle] 6: 165 (1838)	2:335
	<b>Ozothamnus turbinatus</b> DC., Prodr. [A. P. de Candolle] 6: 164 (1838)	2:333
	<b>Pappochroma bellidioides</b> (Hook.f.) G.L.Nesom, Phytologia 85: 277 (1998)	
e	<b>Pappochroma gunnii</b> (Hook.f.) G.L.Nesom, Phytologia 76: 426 (1994)	2:312
e	<b>Pappochroma pappocromum</b> (Labill.) G.L.Nesom, Phytologia 85: 278 (1998)	2:311
e	<b>Pappochroma stellatum</b> (Hook.f.) G.L.Nesom, Phytologia 85: 278 (1998)	2:312
	<b>Pappochroma tasmanicum</b> (Hook.f.) G.L.Nesom, Phytologia 76: 426 (1994)	2:312
e	<b>Pappochroma trigonum</b> (S.J.Forbes & D.I.Morris) G.L.Nesom, Phytologia 85: 279 (1998)	
	<i>Paquerina graminea</i> (Labill.) Cass. ex Less. = <b>Brachyscome graminea</b>	2:295
	<i>Petasites fragrans</i> (Vill.) C.Presl = <b>Petasites pyrenaicus</b>	2:360
i	<b>Petasites pyrenaicus</b> (L.) G.López, Anales Jard. Bot. Madrid 42: 343 (1986)	2:360
	<b>Picris angustifolia</b> DC. subsp. <b>angustifolia</b> , Prodr. [A. P. de Candolle] 7: 130 (1838)	2:383
	<b>Picris angustifolia</b> DC. subsp. <b>merxmuelleri</b> Lack & S.Holzappel, Willdenowia 23: 190 (1993)	2:383
	<i>Picris echioides</i> L. = <b>Helminthotheca echioides</b>	2:383
	<i>Picris hieracioides</i> L. sensu Curtis (1963) = <b>Picris angustifolia</b> (misapplied in Tasmania)	2:383
i	<b>Pilosella aurantiaca</b> (L.) F.W.Schultz & Sch.Bip. subsp. <b>aurantiaca</b> , Flora 42: 426 (1862)	2:385
n i x	<b>Pilosella officinarum</b> Vaill. subsp. <b>officinarum</b> , Königl. Akad. Wiss. Paris Anat. Abh. 5: 703 (1754)	
	<i>Podolepis acuminata</i> R.Br. = <b>Podolepis jaceoides</b>	
	<b>Podolepis decipiens</b> Jeanes, Muelleria 33:26 (2015)	
	<b>Podolepis jaceoides</b> (Sims) Voss, Vilm. Blumengärtn., ed. 3, 1: 537 (1894)	2:348
	<i>Podosperma angustifolia</i> Labill. = <b>Podotheca angustifolia</b>	2:341
?x	<b>Podotheca angustifolia</b> (Labill.) Less., Syn. Gen. Compos. 272 (1832)	2:341
	<i>Pseudognaphalium luteoalbum</i> (L.) Hilliard & B.L.Burtt = <b>Helichrysum luteoalbum</b>	2:318
e	<b>Pterygopappus lawrencei</b> Hook.f., London J. Bot. 6: 120 (1847)	2:316
	<i>Raoulia catipes</i> (DC.) Hook.f. = <b>Ewartia catipes</b>	
	<i>Raoulia mereditheae</i> (F.Muell.) Rodway = <b>Ewartia meredithiae</b>	
	<i>Raoulia planchonii</i> (Hook.f.) Benth. = <b>Ewartia planchonii</b>	
	<i>Rhagadiolus hedynois</i> (L.) All. = <b>Leontodon rhagadioloides</b>	
	<b>Rhodanthe anthemoides</b> (Sieber ex Spreng.) Paul G.Wilson, Nuytsia 8: 386 (1992)	2:324
	<i>Richea glauca</i> Labill. = <b>Craspedia glauca</b>	
	<i>Rutidosis multiflora</i> (Nees) B.L.Rob. = <b>Siloxerus multiflorus</b>	2:342
	<i>Rutidosis pumilo</i> Benth. = <b>Siloxerus multiflorus</b>	

	Scleroleima forsteroides Hook.f. = <b>Abrotanella forsteroides</b>	
i	<b>Scorzonera laciniata</b> L. var. <b>laciniata</b> , Sp. Pl. 2: 791 (1753)	
e	<b>Senecio albogilvus</b> I.Thomps., Muelleria 20: 130 (2004)	2:364
n i	<b>Senecio angulatus</b> L.f., Suppl. Pl.369 (1782)	
	Senecio australis Willd. sensu Rodway (1903) = <b>Senecio linearifolius</b> (misapplied in Tasmania)	
	<b>Senecio biserratus</b> Belcher, Ann. Missouri Bot. Gard. 43: 43 (1956)	2:368
	<b>Senecio campylocarpus</b> I.Thomps., Muelleria 20: 139 (2004)	
	Senecio capillifolius Hook.f. = <b>Senecio pinnatifolius</b> var. <b>capillifolius</b>	2:365
	Senecio centropappus F.Muell. = <b>Centropappus brunonis</b>	2:365
i	<b>Senecio elegans</b> L., Sp. Pl. 2: 869 (1753)	2:366
	<b>Senecio extensus</b> I.Thomps., Muelleria 19: 150 (2004)	
x	<b>Senecio georgianus</b> DC., Prodr. [A. P. de Candolle] 6: 371 (1838)	
	Senecio glandulosus (A.Cunn. ex DC.) Sch.Bip. = <b>Senecio campylocarpus</b>	
	<b>Senecio glomeratus</b> Desf. ex Poir. subsp. <b>glomeratus</b> , Encycl. (Lamarck) Suppl. 5: 130 (1817)	2:370
	<b>Senecio glomeratus</b> Desf. ex Poir. subsp. <b>longifructus</b> I.Thomps., Muelleria 19: 148 (2004)	2:370
	<b>Senecio gunnii</b> (Hook.f.) Belcher, Ann. Missouri Bot. Gard. 43: 60 (1956)	2:370
	<b>Senecio hispidissimus</b> I.Thomps., Muelleria 19: 138 (2004)	
	<b>Senecio hispidulus</b> A.Rich., Voy. Astrolabe 2: 94 t.34 (1834)	2:370
i	<b>Senecio jacobaea</b> L., Sp. Pl. 2: 870 (1753)	2:366
	Senecio lautus G.Forst. ex Willd. sensu Curtis (1963) = <b>Senecio pinnatifolius</b> (sensu lato, misapplied in Tasmania)	2:365
e	<b>Senecio leptocarpus</b> DC., Prodr. [A. P. de Candolle] 6: 372 (1838)	2:364
	<b>Senecio linearifolius</b> A.Rich. var. <b>arachnoideus</b> I.Thomps., Muelleria 20: 98 (2004)	2:366
	<b>Senecio linearifolius</b> A.Rich. var. <b>denticulatus</b> I.Thomps., Muelleria 20: 93 (2004)	2:366
	<b>Senecio linearifolius</b> A.Rich. var. <b>latifolius</b> I.Thomps., Muelleria 20: 96 (2004)	2:366
	<b>Senecio linearifolius</b> A.Rich. var. <b>linearifolius</b> , Voy. Astrolabe 2: 129 (1834)	2:366
	<b>Senecio longipilus</b> I.Thomps., Muelleria 19: 193 (2004)	
x	<b>Senecio macrocarpus</b> F.Muell. ex Belcher, Muelleria 5: 119 (1983)	
	<b>Senecio microbasis</b> I.Thomps., Muelleria 19: 175 (2004)	
	Senecio mikanioides Otto ex Harv. = <b>Delairea odorata</b>	2:367
	<b>Senecio minimus</b> Poir., Encycl. (Lamarck) Suppl. 5: 130 (1817)	2:368
	<b>Senecio odoratus</b> Hornem., Enum. Pl. Hort. Hafn. Suppl. [1809] (1809)	2:367
	<b>Senecio × orarius</b> J.M.Black, Trans. & Proc. Roy. Soc. South Australia 52: 230 (1928)	
e	<b>Senecio papillosus</b> F.Muell., Trans. & Proc. Philos. Inst. Victoria 2: 69 (1857)	2:363
	Senecio pectinatus DC. var. leptocarpus = <b>Senecio leptocarpus</b>	
	Senecio pectinatus DC. var. major F.Muell. ex Belcher recorded in error	
	Senecio pectinatus DC. var. ochroleucus F.Muell. = <b>Senecio albogilvus</b>	2:364
e	<b>Senecio pectinatus</b> DC. var. <b>pectinatus</b> , Prodr. [A. P. de Candolle] 6: 372 (1838)	2:364
	Senecio pectinatus DC. var. pleiocephalus Benth. sensu Rodway (1903) = <b>Senecio pinnatifolius</b> var. <b>alpinus</b> (misapplied in Tasmania)	
	<b>Senecio phelleus</b> I.Thomps., Muelleria 19: 171 (2004)	
	<b>Senecio pinnatifolius</b> A.Rich. var. <b>alpinus</b> (Ali) I.Thomps., Muelleria 21: 53 (2005)	2:365
e	<b>Senecio pinnatifolius</b> A.Rich. var. <b>capillifolius</b> (Hook.f.) I.Thomps., Muelleria 21: 51 (2005)	2:365
	<b>Senecio pinnatifolius</b> A.Rich. var. <b>lanceolatus</b> (Benth.) I.Thomps., Muelleria 21: 49 (2005)	2:365
	<b>Senecio pinnatifolius</b> A.Rich. var. <b>maritimus</b> (Ali) I.Thomps., Muelleria 21: 54 (2005)	2:365

	<b>Senecio pinnatifolius</b> A.Rich. var. <b>pinnatifolius</b> , Voy. Astrolabe 2: 117 (1834)	2:365
	Senecio pinnatifolius A.Rich. var. pleiocephalus (Rodway) Belcher = <b>Senecio pinnatifolius</b> var. <b>alpinus</b>	
	<b>Senecio prenanthoides</b> A.Rich., Voy. Astrolabe 2: 96 (1834)	
e	<b>Senecio primulaefolius</b> F.Muell., Trans. & Proc. Philos. Inst. Victoria 2: 69 (1857)	2:364
	<b>Senecio psilocarpus</b> Belcher & Albr., Muelleria 8: 113 (1994)	
	<b>Senecio quadridentatus</b> Labill., Nov. Holl. Pl. 2: 48 t.194 (1806)	2:369
e	<b>Senecio spathulatus</b> A.Rich. var. <b>spathulatus</b> , Voy. Astrolabe 2: 125 (1834)	2:364
	<b>Senecio squarrosus</b> A.Rich., Voy. Astrolabe 2: 107 t.35 (1834)	2:369
e x	<b>Senecio tasmanicus</b> I.Thomps., Muelleria 19: 158 (2004)	
	<b>Senecio vagus</b> F.Muell. subsp. <b>vagus</b> , Trans. & Proc. Philos. Soc. Victoria 1: 46 (1855)	
	<b>Senecio velleioides</b> A.Cunn. ex DC., Prodr. [A. P. de Candolle] 6: 374 (1838)	2:365
i	<b>Senecio vulgaris</b> L., Sp. Pl. 2: 867 (1753)	2:367
n ?i #	<b>Sigesbeckia orientalis</b> L., Sp. Pl. 2: 900 (1753)	
	<b>Siloxerus multiflorus</b> Nees, Pl. Preiss. [J.G.C.Lehman] 2: 244 (1845)	2:342
i	<b>Silybum marianum</b> (L.) Gaertn., Fruct. Sem. Pl. 2: 378 t.162 (1791)	2:378
	Skirrhophorus eriocephalus Hook.f. ex A.Gray = <b>Angianthus preissianus</b>	
	Solenogyne bellioides Cass. var. gunnii (Hook.f.) G.L.Davis = <b>Solenogyne gunnii</b>	2:292
	<b>Solenogyne dominii</b> L.G.Adams, Brunonia 2: 58 (1979)	
	<b>Solenogyne gunnii</b> (Hook.f.) Cabrera, Blumea 14: 307 (1966)	2:292
i	<b>Soliva sessilis</b> Ruiz & Pav., Syst. Veg. Fl. Peruv. Chil. 225 (1798)	2:358
i	<b>Soliva valdiviana</b> Phil., Linnaea 33: 168 (1864)	
i	<b>Sonchus asper</b> (L.) Hill, Herb. Brit. 1: 47 t.34 (1769)	2:390
	Sonchus asper (L.) Hill subsp. glaucescens (Jord.) Ball = <b>Sonchus asper</b>	
	<b>Sonchus hydrophilus</b> Boulos, Suppl. Black's Fl. S. Austral. 331 (1965)	
	Sonchus megalocarpus (Hook.f.) J.M.Black = <b>Actites megalocarpus</b>	2:390
i	<b>Sonchus oleraceus</b> L., Sp. Pl. 2: 794 (1753)	2:389
i	<b>Symphotrichum novi-belgii</b> (L.) G.L.Nesom, Phytologia 77: 287 (1995) (as "1994")	
i	<b>Symphotrichum subulatum</b> (Michx.) G.L.Nesom, Phytologia 77: 293 (1995) (as "1994")	2:299
i t	<b>Tanacetum cinerariifolium</b> (Trevir.) Sch.Bip., Tanaceteeen: 58 (1844)	
i	<b>Tanacetum parthenium</b> (L.) Sch.Bip., Tanaceteeen: 55 (1844)	2:353
i	<b>Tanacetum vulgare</b> L., Sp. Pl. 2: 844 (1753)	2:360
	<b>Taraxacum aristum</b> G.E.Haglund & Markl., Bot. Not. 117: 197 (1964)	
	<b>Taraxacum cygnorum</b> Hand.-Mazz., Monogr. Taraxacum: 55 (1907)	
	Taraxacum dens-leonis Desf. sensu Rodway (1903) = <b>Taraxacum officinale</b> (misapplied in Tasmania)	
n	Taraxacum kok-saghyz L.E.Rodin – previously listed as naturalised but insufficient evidence exists to support this	2:387
i	<b>Taraxacum officinale</b> F.H.Wigg., Prim. Fl. Holsat. 56 (1780)	2:387
i	<b>Tolpis barbata</b> (L.) Gaertn., Fruct. Sem. Pl. 2: 372 (1791)	2:381
	Tolpis umbellata Bertol. = <b>Tolpis barbata</b>	2:381
i	<b>Tragopogon porrifolius</b> L. subsp. <b>porrifolius</b> , Sp. Pl. 2: 789 (1753)	2:390
	Tripleurospermum inodorum (L.) Sch.Bip. = <b>Tripleurospermum maritimum</b> subsp. <b>inodorum</b>	

i	<b>Tripleurospermum maritimum</b> (L.) W.D.J.Koch subsp. <b>inodorum</b> (L.) Appleg., Taxon 51: 760 (2002)	
	<b>Triptilodiscus pygmaeus</b> Turcz., Bull. Soc. Imp. Naturalistes Moscou 24: 66 (1851)	
i t	<b>Urospermum dalechampii</b> (L.) F.W.Schmidt, Samml. Phys.-Oekon. Aufsätze 1: 275 (1795)	2:391
i	<b>Vellereophyton dealbatum</b> (Thunb.) Hilliard & B.L.Burtt, Bot. J. Linn. Soc. 82: 210 (1981)	2:318
	<b>Vittadinia australasica</b> (Turcz.) N.T.Burb. var. <b>oricola</b> N.T.Burb., Brunonia 5: 44 (1982)	
	Vittadinia australis A.Rich. sensu Bentham (1867) misapplied to <b>V. australasica</b> and <b>V. cuneata</b>	
	Vittadinia australis A.Rich. var. megacephala F.Muell. ex Benth. = Vittadinia megacephala	
e	<b>Vittadinia burbridgeae</b> A.M.Gray & Rozefelds, Kanunnah 1: 9 (2005)	
	<b>Vittadinia cuneata</b> DC. var. <b>cuneata</b> , Prodr. [A. P. de Candolle] 5: 281 (1836)	2:313
	<b>Vittadinia gracilis</b> (Hook.f.) N.T.Burb., Brunonia 5: 54 (1982)	2:313
	Vittadinia megacephala (F.Muell. ex Benth.) J.M.Black recorded in error	2:314
	<b>Vittadinia muelleri</b> N.T.Burb., Proc. Linn. Soc. New South Wales 93: 440 fig.1D (1969)	
	Vittadinia scabra DC. sensu Hooker (1860) = <b>Vittadinia cuneata</b> (misapplied in Tasmania)	
	Vittadinia triloba (Gaudich.) DC. sensu Curtis (1963) = <b>Vittadinia cuneata</b> (misapplied in Tasmania)	2:313
	Vittadinia triloba (Gaudich.) DC. var. lanuginosa J.M.Black sensu Curtis (1963) = <b>Vittadinia gracilis</b> (misapplied in Tasmania)	2:313
	Vittadinia triloba (Gaudich.) DC. var. megacephala (F.Muell. ex Benth.) Ewart = Vittadinia megacephala	2:314
i	<b>Xanthium spinosum</b> L., Sp. Pl. 2: 987 (1753)	2:349
n e	<b>Xerochrysum alpinum</b> Paul G.Wilson, Nuytsia 28: 36 (2017)	
	<b>Xerochrysum bicolor</b> (Lindl.) R.J.Bayer, Kew Bull. 56: 1014 (2001)	2:331
	<b>Xerochrysum bracteatum</b> (Vent.) Tzvelev, Novosti Sist. Vyssh. Rast. 27: 151 (1990)	
e	<b>Xerochrysum collierianum</b> A.M.Buchanan, Muelleria 20: 49 (2004)	
n e	<b>Xerochrysum milliganii</b> (Hook.f.) Paul G.Wilson, Taxon 64: 106 (2015)	2:330
	<b>Xerochrysum palustre</b> (Flann) R.J.Bayer, Kew Bull. 56: 1014 (2001)	
	<b>Xerochrysum papillosum</b> (Labill.) R.J.Bayer, Kew Bull. 56: 1015 (2001)	2:331
	<b>Xerochrysum subundulatum</b> (Sch.Bip.) R.J.Bayer, Kew Bull. 56: 1015 (2001)	2:330
i	<b>Xerochrysum viscosum</b> (Sieber ex DC.) R.J.Bayer, Kew Bulletin 56: 1015 (2001)	
	<b>BASELLACEAE</b>	FTO 99
n i *	<i>Anredera cordifolia</i> (Ten.) Steenis, Fl. Malesiana 5: 303 fig.2 a-j (1957)	3:580
	<b>BERBERIDACEAE</b>	FTO 46
i	<b>Berberis darwinii</b> Hook., Icon. Pl. 7: t.672 (1844)	
	<b>BETULACEAE</b>	
n	<i>Alnus cordata</i> (Loisel.) Duby – previously listed as naturalised but insufficient evidence exists to support this	3:645
n i *	<i>Alnus glutinosa</i> (L.) Gaertn., Fruct. Sem. Pl. 2: 54 (1791)	
i	<b>Betula pendula</b> Roth, Tent. Fl. Germ. 1: 405 (1788)	
	<b>BIGNONIACEAE</b>	FTO 121
	<b>Pandorea pandorana</b> (Andrews) Steenis, Bull. Jard. Bot. Buitenzorg 10: 198 (1928)	3:538
	<b>BORAGINACEAE</b>	
i	<b>Amsinckia calycina</b> (Moris) Chater, Bot. J. Linn. Soc. 64: 380 (1971)	3:489
	<i>Amsinckia hispida</i> (Ruiz & Pav.) I.M.Johnst. = <b>Amsinckia calycina</b>	3:489



i	<b>Anchusa arvensis</b> (L.) M.Bieb., Fl. Taur.-Caucas. 1: 123 (1808)	
	<i>Anchusa sempervirens</i> L. = <b>Pentaglottis sempervirens</b>	3:491
	<b>Austrocynoglossum latifolium</b> (R.Br.) Popov ex R.R.Mill, Notes Roy. Bot. Gard. Edinburgh 46: 44 (1989)	3:488
i	<b>Borago officinalis</b> L., Sp. Pl. 1: 137 (1753)	3:490
i	<b>Buglossoides arvensis</b> (L.) I.M.Johnst., J. Arnold Arbor. 35: 42 (1954)	3:494
	<b>Cynoglossum australe</b> R.Br., Prodr. Fl. Nov. Holland. 495 (1810)	3:489
	<i>Cynoglossum latifolium</i> R.Br. = <b>Austrocynoglossum latifolium</b>	3:488
	<b>Cynoglossum suaveolens</b> R.Br., Prodr. Fl. Nov. Holland. 495 (1810)	3:488
i	<b>Echium candicans</b> L.f., Suppl. Pl. 131 (1782)	
	<i>Echium lycopsis</i> L. sensu Curtis (1967) = <b>Echium plantagineum</b> (misapplied in Tasmania)	3:495
i	<b>Echium plantagineum</b> L., Mant. Pl. 2: 202 (1771)	3:495
i	<b>Echium vulgare</b> L., Sp. Pl. 1: 139 (1753)	3:495
	<i>Exarrhena suaveolens</i> R.Br. = <b>Myosotis exarrhena</b>	
	<i>Lithospermum arvense</i> L. = <b>Buglossoides arvensis</b>	3:494
n i t	<b>Lithospermum officinale</b> L., Sp. Pl. 1: 132 (1753)	3:494
	<i>Lycopsis arvensis</i> L. = <b>Anchusa arvensis</b>	
i	<b>Myosotis arvensis</b> (L.) Hill, Veg. Syst. 7: 55 (1764)	
	<b>Myosotis australis</b> R.Br., Prodr. Fl. Nov. Holland. 495 (1810)	3:492
	<i>Myosotis caespitosa</i> Schultz = <b>Myosotis laxa</b> subsp. <b>caespitosa</b>	3:493
i	<b>Myosotis discolor</b> Pers., Syst. Veg., ed. 15 (J.A.Murray): 190 (1798)	3:493
	<b>Myosotis exarrhena</b> F.Muell., Syst. Census Austral. Pl. Suppl. 4: 7 (1889)	3:492
i	<b>Myosotis laxa</b> Lehm. subsp. <b>caespitosa</b> (Schultz) Hyl. ex Nordh., Norsk. Fl. (Nordhagen): 529 (1940)	3:493
i	<b>Myosotis scorpioides</b> L., Sp. Pl. 1: 131 (1753)	3:492
	<i>Myosotis suaveolens</i> (R.Br.) Poir. = <b>Myosotis exarrhena</b>	3:492
i	<b>Myosotis sylvatica</b> Hoffm., Deutschl. Fl. (Hoffm.): 61 (1791)	3:493
i	<b>Nonea lutea</b> (Desr.) Rchb. ex A.DC., Prodr. [A. P. de Candolle] 10: 28 (1846)	
i	<b>Pentaglottis sempervirens</b> (L.) Tausch ex L.H.Bailey, Man. Cult. Pl., ed. 2: 837 (1949)	3:491
	<i>Symphytum peregrinum</i> Ledeb. = <b>Symphytum uplandicum</b>	3:490
n i *	<i>Symphytum</i> × <i>uplandicum</i> Nyman, Syll. Fl. Eur. 80 (1855)	3:490
<b>BRASSICACEAE (CRUCIFERAE)</b>		
	<i>Alyssum maritimum</i> (L.) Lam. = <b>Lobularia maritima</b>	1:45
i	<b>Arabidopsis thaliana</b> (L.) Heynh., Fl. Sachsen 1: 538 (1842)	1:50
x	<b>Ballantinia antipoda</b> (F.Muell.) E.A.Shaw, Contr. Gray Herb. 205: 161 (1974)	1:42
e	<b>Barbarea australis</b> Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 14 (1852)	1:47
i	<b>Barbarea intermedia</b> Boreau, Fl. Centre France [Boreau] 2: 48 (1840)	1:48
i	<b>Barbarea verna</b> (Mill.) Asch., Fl. Brandenburg 1: 36 (1860)	1:48
	<i>Barbarea vulgaris</i> W.T.Aiton sensu Bentham (1863) = <b>Barbarea australis</b> (partly misapplied in Tasmania)	
i #	<b>Brassica</b> × <b>juncea</b> (L.) Czern., Conspect. Pl. Chark. 8 (1859)	
i	<b>Brassica</b> × <b>napus</b> L., Sp. Pl. 2: 666 (1753)	1:34
i	<b>Brassica nigra</b> (L.) W.D.J.Koch, Deutschl. Fl. (Röhling), ed. 3, 4: 713 (1833)	1:35
n i *	<i>Brassica oleracea</i> L., Sp. Pl. 2: 667 (1753)	
i	<b>Brassica rapa</b> L., Sp. Pl. 2: 666 (1753)	1:34

	Brassica sinapistrum Boiss. = <b>Sinapis arvensis</b>	
i	<b>Brassica tournefortii</b> Gouan, Ill. Observ. Bot. 3: 44, t.20A (1773)	
i	<b>Cakile edentula</b> (Bigelow) Hook., Fl. Bor.-Amer. (Hooker) 1: 59 (1830)	1:38
i	<b>Cakile maritima</b> Scop. subsp. <b>maritima</b> , Fl. Carniol., ed. 2, 2: 35 (1772)	1:38
	Camelina alyssum (Mill.) Thell. sensu Curtis (1975) = <b>Camelina sativa</b> (misapplied in Tasmania)	1:51
i	<b>Camelina sativa</b> (L.) Crantz, Stirp. Austr. Fasc.1: 17 (1762)	1:51
	Capsella antipoda F.Muell. = <b>Ballantinia antipoda</b>	
	Capsella australis (Hook.f.) Benth. = <b>Ballantinia antipoda</b>	
i	<b>Capsella bursa-pastoris</b> (L.) Medik., Pfl.-Gatt. 1: 85 (1792)	1:43
	Capsella elliptica C.A.Mey. nom. illeg. = <b>Hornungia procumbens</b>	
	Capsella procumbens (L.) Fr. = <b>Hornungia procumbens</b>	
	Capsella tasmanica (Hook.) F.Muell. ex B.D.Jacks. = <i>Hutchinsia tasmanica</i>	1:43
	<b>Cardamine astoniae</b> I.Thomps., Muelleria 9: 156 (1996)	
	Cardamine aff. corymbosa Hook.f. sensu Hewson (1982) = <b>Cardamine papillata</b>	
	Cardamine dictyosperma Hook. = <b>Rorippa dictyosperma</b>	
i	<b>Cardamine flexuosa</b> With., Arr. Brit. Pl., ed. 3, 3: 578 (1796)	
	<b>Cardamine franklinensis</b> I.Thomps., Muelleria 9: 152 (1996)	
	<b>Cardamine gunnii</b> Hewson, Fl. Australia 8: 390 (1982)	1:46
	Cardamine heterophylla Hook. = <b>Cardamine gunnii</b>	1:46
i	<b>Cardamine hirsuta</b> L., Sp. Pl. 2: 655 (1753)	1:46
	Cardamine hirsuta L. var. parvifolia Rodway – a name of uncertain application	
	Cardamine hirsuta L. var. tenuifolia (Hook.) Rodway = <b>Cardamine tenuifolia</b>	
	Cardamine intermedia Hook. sensu Curtis (1956) = <b>Cardamine lilacina</b>	
	<b>Cardamine lilacina</b> Hook., Companion Bot. Mag. 1: 273 (1836)	
	<b>Cardamine papillata</b> I.Thomps., Muelleria 9: 161 (1996)	
	<b>Cardamine paucijuga</b> Turcz., Bull. Soc. Imp. Naturalistes Moscou 27(2): 295 (1854)	
	Cardamine pratensis L. sensu Hooker (1860) = <b>Cardamine tenuifolia</b> (misapplied in Tasmania)	
	Cardamine pratensis L. var. lilacina (Hook.) Hook.f. = <b>Cardamine lilacina</b>	
	Cardamine radicata Hook.f. = <b>Pachycladon radicans</b>	
	Cardamine stylosa DC. = <b>Rorippa gigantea</b>	
	<b>Cardamine tenuifolia</b> Hook., J. Bot. (Hooker) 1: 247 (1837)	1:46
	<b>Cardamine tryssa</b> I.Thomps., Muelleria 18: 27 (2004)	
	Cardaria draba (L.) Desv. = <b>Lepidium draba</b>	1:42
n	<i>Carrichtera annua</i> (L.) DC. – previously listed as naturalised but insufficient evidence exists to support this	
	Cheesemaniania radicata (Hook.f.) O.E.Schulz = <b>Pachycladon radicans</b>	1:47
	Coronopus didymus (L.) Sm. = <b>Lepidium didymum</b>	1:41
	Coronopus squamatus (Forssk.) Asch. = <b>Lepidium coronopus</b>	1:41
	Cuphonotus antipodus (F.Muell.) J.M.Black = <b>Ballantinia antipoda</b>	1:42
i	<b>Diplotaxis muralis</b> (L.) DC., Syst. Nat. [Candolle] 2: 634 (1821)	1:36
i	<b>Diplotaxis tenuifolia</b> (L.) DC., Syst. Nat. [Candolle] 2: 632 (1821)	1:36
	<i>Draba muralis</i> L. sensu Curtis & Morris (1975) and earlier = <b>Draba nemorosa</b> (misapplied in Tasmania)	1:45
i	<b>Draba muralis</b> L., Sp. Pl. 2: 642 (1753)	1:45
i t	<b>Draba nemorosa</b> L., Sp. Pl. 2: 643 (1753)	1:45

i	<b>Erophila verna</b> (L.) Chevall. subsp. <b>praecox</b> (Steven) Walters, Feddes Repert. Spec. Nov. Regni Veg. 69: 57 (1964)	1:45
i	<b>Erophila verna</b> (L.) Chevall. subsp. <b>verna</b> , Fl. Gén. Env. Paris 2: 898 (1827)	1:45
n	<i>Eruca sativa</i> Mill. – previously listed as naturalised but insufficient evidence exists to support this	
	<i>Eruca vesicaria</i> (L.) Cav. subsp. <i>sativa</i> (Mill.) Thell. = <i>Eruca sativa</i>	
	<b>Geococcus pusillus</b> J.Drumm. ex Harv., Hooker's J. Bot. Kew Gard. Misc. 7: 52 (1855)	1:50
i	<b>Hirschfeldia incana</b> (L.) Lagr.-Foss., Fl. Tarn Garonne: 19 (1847)	1:35
i	<b>Hornungia procumbens</b> (L.) Hayek, Repert. Spec. Nov. Regni Veg. Beih. 30: 480 (1925)	1:44
	<i>Hutchinsia australis</i> Hook.f. = <b>Ballantinia antipoda</b>	
	<i>Hutchinsia tasmanica</i> Hook. – a name of uncertain application	
	<i>Hymenolobus procumbens</i> (L.) Nutt. = <b>Hornungia procumbens</b>	1:44
i	<b>Lepidium africanum</b> (Burm.f.) DC., Syst. Nat. [Candolle] 2: 522 (1821)	
i	<b>Lepidium bonariense</b> L., Sp. Pl. 2: 645 (1753)	
i	<b>Lepidium campestre</b> (L.) W.T.Aiton, Hortus Kew. (W.T.Aiton), ed. 2, 4: 88 (1812)	1:40
i	<b>Lepidium coronopus</b> (L.) Al-Shehbaz, Novon 14: 156 (2004)	1:41
	<i>Lepidium cuneifolium</i> DC. = <b>Lepidium foliosum</b>	
	<b>Lepidium desvauxii</b> Thell., Neue Denkschr. Allg. Schweiz. Ges. Gesamten Naturwiss. 41: 307 (1906)	1:39
i	<b>Lepidium didymum</b> L., Mant. Pl. 92 (1767)	1:41
i	<b>Lepidium draba</b> L., Sp. Pl. 2: 645 (1753)	1:42
t	<b>Lepidium flexicaule</b> Kirk, Trans. & Proc. New Zealand Inst. 14: 380 (1882)	
	<b>Lepidium foliosum</b> Desv., J. Bot. (Desvaux) 3: 164, 180 (1815)	1:39
	<i>Lepidium halmaturinum</i> J.M.Black = <b>Lepidium desvauxii</b>	1:39
n	<i>Lepidium heterophyllum</i> Benth. – previously listed as naturalised but insufficient evidence exists to support this	1:40
	<b>Lepidium hyssopifolium</b> Desv., J. Bot. (Desvaux) 3: 179 (1815)	
i	<b>Lepidium latifolium</b> L., Sp. Pl. 2: 644 (1753)	1:41
	<i>Lepidium praetervisum</i> Domin = <b>Lepidium desvauxii</b>	1:39
	<b>Lepidium pseudotasmanicum</b> Thell., Neue Denkschr. Allg. Schweiz. Ges. Gesamten Naturwiss. 41: 307 (1906)	1:39
	<i>Lepidium ruderales</i> L. sensu Bentham (1863) = <b>Lepidium foliosum</b> (misapplied in Tasmania)	
i	<b>Lepidium sativum</b> L., Sp. Pl. 2: 644 (1753)	1:40
	<i>Lepidium squamatum</i> Forssk. = <b>Lepidium coronopus</b>	1:41
	<i>Lepidium tasmanicum</i> Thell. – a name of uncertain application	
i	<b>Lobularia maritima</b> (L.) Desv., J. Bot. (Desvaux) 3: 162 (1815)	1:45
n i	<b>Lunaria annua</b> L., Sp. Pl. 2: 653 (1753)	
i	<b>Matthiola incana</b> (L.) W.T.Aiton, Hortus Kew. (W.T.Aiton), ed. 2, 4: 119 (1812)	
n i	<b>Nasturtium microphyllum</b> Boenn. ex Rchb., Fl. Germ. Excurs. 2: 683 (1830)	1:49
i	<b>Nasturtium officinale</b> W.T.Aiton, Hortus Kew., (W.T.Aiton), ed. 2, 4: 110 (1812)	1:49
	<i>Nasturtium palustre</i> (L.) DC. = <b>Rorippa palustris</b>	
	<i>Nasturtium terrestre</i> (With.) Aiton = <b>Rorippa palustris</b>	
e	<b>Pachycladon radicans</b> (Hook.f.) Heenan & A.D.Mitch., New Zealand J. Bot. 40: 558 (2002)	1:47
n i *	<i>Raphanus maritimus</i> Sm., Engl. Bot. 23: 1643 (1806)	1:37
i	<b>Raphanus raphanistrum</b> L., Sp. Pl. 2: 669 (1753)	1:37

i	<b>Rapistrum rugosum</b> (L.) All., Fl. Pedem. 1: 257 t.78 (1785)	1:37
	<b>Rorippa dictyosperma</b> (Hook.) L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 97 (1962)	1:49
	<b>Rorippa gigantea</b> (Hook.) Garn.-Jones, New Zealand J. Bot. 16: 119 (1978)	1:48
	<i>Rorippa islandica</i> (Oeder) Borbás sensu Curtis & Morris (1975) = <b>Rorippa palustris</b> (misapplied in Tasmania)	1:48
	<i>Rorippa microphylla</i> (Boenn. ex Rchb.) H.Hyl. = <b>Nasturtium microphyllum</b>	1:49
	<i>Rorippa nasturtium-aquaticum</i> (L.) Hayek = <b>Nasturtium officinale</b>	1:49
i	<b>Rorippa palustris</b> (L.) Besser, Enum. Pl. [Besser]: 27 (1822)	1:48
	<i>Rorippa stylosa</i> (DC.) Allan = <b>Rorippa gigantea</b>	1:48
i #	<b>Rorippa sylvestris</b> (L.) Besser, Enum. Pl. [Besser]: 27 (1822)	
	<i>Senebiera coronopus</i> (L.) Poir. = <b>Lepidium coronopus</b>	
	<i>Senebiera didyma</i> (L.) Pers. = <b>Lepidium didymum</b>	
i	<b>Sinapis alba</b> L., Sp. Pl. 2: 668 (1753)	
i	<b>Sinapis arvensis</b> L., Sp. Pl. 2: 668 (1753)	1:35
i	<b>Sisymbrium irio</b> L., Sp. Pl. 2: 659 (1753)	
i	<b>Sisymbrium officinale</b> (L.) Scop., Fl. Carniol., ed. 2, 2: 26 (1772)	1:50
i	<b>Sisymbrium orientale</b> L., Cent. Pl. 2: 24 (1756)	1:50
	<b>Stenopetalum lineare</b> R.Br. ex DC., Syst. Nat. [Candolle] 2: 513 (1821)	1:44
i t	<b>Teesdalia nudicaulis</b> (L.) W.T.Aiton, Hortus Kew. (W.T.Aiton), ed. 2, 4: 83 (1812)	1:43
n i *	<i>Thlaspi arvense</i> L., Sp. Pl. 2: 646 (1753)	
	<i>Thlaspi tasmanicum</i> (Hook.) Hook.f. = <i>Hutchinsia tasmanica</i>	
	<b>BRUNONIACEAE</b>	FTO 129
	<b>Brunonia australis</b> Sm. ex R.Br., Prodr. Fl. Nov. Holland. 590 (1810)	2:405
	<b>BUDDLEJACEAE</b>	
i	<b>Buddleja davidii</b> Franch., Nouv. Arch. Mus. Hist. Nat. Paris, Ser. 2, 10: 65 (1888)	
	<b>CALLITRICHACEAE</b>	
	<b>Callitriche brachycarpa</b> Hegelm., Verh. Bot. Vereins Prov. Brandenburg 10: 115 (1868)	1:195
n i *	<i>Callitriche brutia</i> Petagna subsp. <i>brutia</i> , Institutiones Botanicae 2: 10 (1787)	
	<b>Callitriche sonderi</b> Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 18 (1867)	
i	<b>Callitriche stagnalis</b> Scop., Fl. Carniol., ed. 2, 2: 251 (1772)	1:195
	<b>Callitriche umbonata</b> Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 19 (1867)	1:196
	<i>Callitriche verna</i> L. sensu Hooker (1860), Bentham (1864) = <b>Callitriche stagnalis</b> (misapplied in Tasmania)	
	<b>CAMPANULACEAE</b>	
	<i>Campanula gracilis</i> G.Forst. var. <i>littoralis</i> (Labill.) R.Br. = <b>Wahlenbergia gymnoclada</b>	
	<i>Campanula gracilis</i> G.Forst. var. <i>vincaeflora</i> R.Br. = <b>Wahlenbergia gracilis</b>	
	<i>Campanula littoralis</i> Labill. = <b>Wahlenbergia gymnoclada</b>	
n i *	<i>Campanula rapunculoides</i> L., Sp. Pl. 1: 165 (1753)	2:410
	<b>Isotoma fluviatilis</b> (R.Br.) F.Muell. ex Benth. subsp. <i>australis</i> McComb, Contr. New South Wales Natl. Herb. 4: 109 (1970)	2:411
	<i>Lobelia alata</i> Labill. = <b>Lobelia anceps</b>	2:414
	<b>Lobelia anceps</b> L.f., Suppl. Pl. 395 (1782)	2:414
	<b>Lobelia browniana</b> Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 5: 71 (1819)	

	<i>Lobelia cuneiformis</i> Labill. = <b>Lobelia anceps</b>	
	<b>Lobelia dentata</b> Cav., Icon. Pl. 6: 14, t. 522 (1800)	
i #	<b>Lobelia erinus</b> L., Sp. Pl. 2: 932 (1753)	
	<i>Lobelia fluviatilis</i> R.Br. = <b>Isotoma fluviatilis</b>	
	<b>Lobelia gibbosa</b> Labill., Nov. Holl. Pl. 1: 50, t.71 (1805)	2:414
	<i>Lobelia gibbosa</i> Labill. var. <i>browniana</i> (Schult.) F.M.Bailey = <b>Lobelia browniana</b>	
	<b>Lobelia irrigua</b> R.Br., Prodr. Fl. Nov. Holland. 563 (1810)	2:412
	<b>Lobelia pedunculata</b> R.Br., Prodr. Fl. Nov. Holland. 563 (1810)	2:413
	<i>Lobelia platycalyx</i> (F.Muell.) F.Muell. = <b>Lobelia irrigua</b>	
	<b>Lobelia pratioides</b> Benth., Fl. Austral. 4: 131 (1868)	2:415
	<b>Lobelia rhombifolia</b> de Vriese, Pl. Preiss. [J.G.C.Lehman] 1: 397 (1845)	2:414
	<b>Lobelia simplicicaulis</b> R.Br., Prodr. Fl. Nov. Holland. 564 (1810)	
	<b>Lobelia surrepens</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 237 (1856)	2:412
	<i>Pratia irrigua</i> (R.Br.) Benth. = <b>Lobelia irrigua</b>	2:412
	<i>Pratia pedunculata</i> (R.Br.) Benth. = <b>Lobelia pedunculata</b>	2:413
	<i>Pratia platycalyx</i> (F.Muell.) Benth. = <b>Lobelia irrigua</b>	2:412
	<i>Pratia surrepens</i> (Hook.f.) E.Wimm. = <b>Lobelia surrepens</b>	2:412
	<b>Wahlenbergia ceracea</b> Lothian, Victorian Nat. 72: 166 (1956)	2:407
	<i>Wahlenbergia consimilis</i> Lothian = <b>Wahlenbergia stricta</b>	2:408
	<b>Wahlenbergia gracilentata</b> Lothian, Proc. Linn. Soc. New South Wales 71: 217 (1947)	2:407
	<b>Wahlenbergia gracilis</b> (G.Forst.) A.DC., Monogr. Campan. 142 (1830)	2:409
	<b>Wahlenbergia gymnoclada</b> Lothian, Proc. Linn. Soc. New South Wales 71: 227 (1947)	2:408
	<b>Wahlenbergia littoricola</b> P.J.Sm. subsp. <i>littoricola</i> , Fl. S. Austral. [J.M.Black], ed. 4, 3: 1380 (1986)	
	<b>Wahlenbergia multicaulis</b> Benth., Enum. Pl. [Endlicher]: 75 (1837)	2:409
	<i>Wahlenbergia quadrifida</i> (R.Br.) A.DC. = <b>Wahlenbergia gracilis</b>	2:409
e	<b>Wahlenbergia saxicola</b> (R.Br.) A.DC., Monogr. Campan. 144 (1830)	2:410
	<b>Wahlenbergia stricta</b> (R.Br.) Sweet subsp. <i>stricta</i> , Hort. Brit. [Sweet], ed. 2: 593 (1830)	2:408
	<i>Wahlenbergia tadgellii</i> Lothian = <b>Wahlenbergia multicaulis</b>	2:409
<b>CANNABACEAE</b>		
i	<b>Humulus lupulus</b> L., Sp. Pl. 2: 1028 (1753)	
<b>CAPRIFOLIACEAE</b>		
i	<b>Leycesteria formosa</b> Wall., Fl. Ind. (Carey & Wallich ed.) 2: 182 (1824)	2:266
i	<b>Lonicera japonica</b> Thunb., Syst. Veg., ed. 14 (J.A.Murray): 216 (1784)	
i #	<b>Lonicera periclymenum</b> L., Sp. Pl. 1: 173 (1753)	2:266
	<b>Sambucus gaudichaudiana</b> DC., Prodr. [A. P. de Candolle] 4: 322 (1830)	2:265
i	<b>Sambucus nigra</b> L., Sp. Pl. 1: 269 (1753)	2:265
n i *	<i>Viburnum tinus</i> L., Sp. Pl. 1: 267 (1753)	
<b>CARYOPHYLLACEAE</b>		
i	<b>Agrostemma githago</b> L., Sp. Pl. 1: 435 (1753)	1:69
i	<b>Arenaria leptoclados</b> (Rchb.) Guss., Fl. Sicul. Syn. 2: 824 (1845)	
i	<b>Arenaria serpyllifolia</b> L., Sp. Pl. 1: 423 (1753)	1:72
i	<b>Cerastium balearicum</b> F.Herm., Verh. Bot. Vereins Prov. Brandenburg 54: 247 (1913)	
	<i>Cerastium fontanum</i> Baumg. subsp. <i>vulgare</i> (Hartm.) Greuter & Burdet = <b>Cerastium vulgare</b>	1:69

i	<b>Cerastium glomeratum</b> Thuill., Fl. Env. Paris, ed. 2: 226 (1799)	1:69
i	<b>Cerastium semidecandrum</b> L., Sp. Pl. 1: 438 (1753) Cerastium viscosum L. = <b>Cerastium vulgare</b>	
i	<b>Cerastium vulgare</b> Hartm., Hand. Skand. Fl. 182 (1820) Cerastium vulgatum L. sensu Curtis (1956) = <b>Cerastium vulgare</b> (misapplied in Tasmania)	1:69
	<b>Colobanthus affinis</b> (Hook.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 45 (1855)	1:74
	<b>Colobanthus apetalus</b> (Labill.) Druce var. <b>apetalus</b> , Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 616 (1917)	74
	Colobanthus billardierei Fenzl nom. illeg. = <b>Colobanthus apetalus</b> var. <b>apetalus</b>	
	<b>Colobanthus curtisiae</b> J.G.West, Aspects of Tasmanian Botany: 75 (1991)	
	<b>Colobanthus pulvinatus</b> F.Muell., Trans. Philos. Soc. Victoria 1: 101 (1855)	
i	<b>Dianthus armeria</b> L., Sp. Pl. 1: 410 (1753) Githago segetum Link = <b>Agrostemma githago</b>	
	Gypsophila australis (Schltdl.) A.Gray = <b>Gypsophila tubulosa</b>	1:66
i	<b>Gypsophila tubulosa</b> (Jaub. & Spach) Boiss., Diagn. Pl. Orient., Ser. 1, 1: 11 (1842) Kohlruschia prolifera (L.) Kunth sensu Curtis (1956) = <b>Petrorhagia nanteuillii</b> (misapplied in Tasmania)	1:66
	Kohlruschia velutina (Guss.) Rchb. = <b>Petrorhagia dubia</b>	
	Lychnis coronaria (L.) Desr. = <b>Silene coronaria</b>	
	Melandrium album (Mill.) Garcke = <b>Silene latifolia</b> subsp. <b>alba</b>	
	Minuartia hybrida (Vill.) Schischk. sensu Buchanan et al. (1989) = <b>Minuartia mediterranea</b> (misapplied in Tasmania)	
i	<b>Minuartia mediterranea</b> (Ledeb. ex Link) K.Maly, Glasn. Zemaljsk. Muz. Bosni Hercegovini 20: 563 (1908) Mniarum biflorum J.R.Forst. & G.Forst. = <b>Scleranthus biflorus</b> Mniarum fasciculatum R.Br. = <b>Scleranthus fasciculatus</b>	
i	<b>Moenchia erecta</b> (L.) G.Gaertn., B.Mey. & Scherb., Oekon. Fl. Wetterau 1: 219 (1799)	1:70
i	<b>Paronychia brasiliana</b> DC., Encycl. (Lamarck) 5: 23 (1804)	
i	<b>Petrorhagia dubia</b> (Raf.) G.López & Romo, Anales Jard. Bot. Madrid 45: 363 (1988)	1:66
i	<b>Petrorhagia nanteuillii</b> (Burnat) P.W.Ball & Heywood, Bull. Brit. Mus. (Nat. Hist.) Bot. 3: 164 (1964)	1:65
	Petrorhagia prolifera (L.) P.W.Ball & Heywood sensu Curtis & Morris (1975) = <b>Petrorhagia nanteuillii</b> (misapplied in Tasmania)	1:65
	Petrorhagia velutina (Guss.) P.W.Ball & Heywood = <b>Petrorhagia dubia</b>	1:66
i	<b>Polycarpon tetraphyllum</b> (L.) L., Syst. Nat., ed. 10, 2: 881 (1759)	1:76
i	<b>Sagina apetala</b> Ard., Animadv. Bot. Spec. Alt. 2: 22 (1764)	1:73
e	<b>Sagina diemensis</b> L.G.Adams, Muelleria 9: 64 (1996)	
i	<b>Sagina maritima</b> Don, Herb. Brit. [Don] 7: 155 (1806)	1:73
	<b>Sagina namadgi</b> L.G.Adams, Muelleria 9: 63 (1996)	
i	<b>Sagina procumbens</b> L., Sp. Pl. 1: 128 (1753)	1:73
i	<b>Saponaria officinalis</b> L., Sp. Pl. 1: 408 (1753) Saponaria tubulosa (Jaub. & Spach) F.Muell. = <b>Gypsophila tubulosa</b>	1:67
i	<b>Scleranthus annuus</b> L., Sp. Pl. 1: 406 (1753) <b>Scleranthus biflorus</b> (J.R.Forst. & G.Forst.) Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 74 (1852)	1:77 1:76

	<b>Scleranthus brockiei</b> P.A.Will., Rec. Domin. Mus. 3: 16 (1956)	
	<b>Scleranthus diander</b> R.Br., Prodr. Fl. Nov. Holland. 412 (1810)	1:77
	<b>Scleranthus fasciculatus</b> (R.Br.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.)1: 42 (1855)	
	<i>Silene alba</i> (Mill.) E.H.L.Krause = <b>Silene latifolia</b> subsp. <b>alba</b>	1:67
	<i>Silene colorata</i> Poir. – previously listed as naturalised but insufficient evidence exists to support this	
n	<i>Silene conica</i> L. – previously listed as naturalised but insufficient evidence exists to support this	
i	<b>Silene coronaria</b> (L.) Clairv., Man. Herbor. Suisse 145 (1811)	
i #	<b>Silene dichotoma</b> Ehrh., Beitr. Naturk. [Ehrhart] 7: 144 (1792)	
i	<b>Silene gallica</b> L. var. <b>gallica</b> , Sp. Pl. 1: 417 (1753)	1:68
i	<b>Silene gallica</b> L. var. <b>quinquevulnera</b> (L.) W.D.J.Koch, Syn. Fl. Germ. Helv. 1: 100 (1835)	1:68
i	<b>Silene latifolia</b> Poir. subsp. <b>alba</b> (Mill.) Greuter & Burdet, Willdenowia 12: 189 (1982)	1:67
	<i>Silene longicaulis</i> Pourr. ex Lag. sensu Buchanan (1995) = <i>Silene colorata</i> (misapplied in Tasmania)	
i	<b>Silene nocturna</b> L., Sp. Pl. 1: 416 (1753)	1:68
	<i>Silene pratensis</i> (Rafn) Godr. & Gren. = <b>Silene latifolia</b> subsp. <b>alba</b>	1:67
i	<b>Silene vulgaris</b> (Moench) Garcke, Fl. N. Mitt.-Deutschland, ed. 9: 64 (1869)	1:68
	<i>Spergula apetala</i> Labill. = <b>Colobanthus apetalus</b>	
i	<b>Spergula arvensis</b> L. var. <b>arvensis</b> , Sp. Pl. 1: 440 (1753)	
i	<b>Spergula arvensis</b> L. var. <b>sativa</b> (Boenn.) Mert. & W.D.J.Koch, Deutschl. Fl. (Mertens & W. D. J. Koch), ed. 3. 3: 360 (1831)	
i	<b>Spergularia bocconeii</b> (Scheele) Graebn., Syn. Mitteleur. Fl. [Ascherson & Graebner] 5: 849 (1919)	
	<i>Spergularia marginata</i> Kitt. sensu Curtis (1956) = <b>Spergularia tasmanica</b> (misapplied in Tasmania)	
i	<b>Spergularia marina</b> (L.) Griseb., Spic. Fl. Rumel. 1: 213 (1843)	1:75
	<i>Spergularia media</i> (L.) C.Presl sensu Curtis & Morris (1975) = <b>Spergularia tasmanica</b> (misapplied in Tasmania)	1:75
i	<b>Spergularia rubra</b> (L.) J.Presl & C.Presl, Fl. Cech. 94 (1819)	1:75
	<i>Spergularia rubra</i> (L.) J.Presl & C.Presl var. <i>marina</i> (Roth) A.Gray = <b>Spergularia marina</b>	
	<i>Spergularia salina</i> J.Presl & C.Presl = <b>Spergularia marina</b>	
	<b>Spergularia tasmanica</b> (Kindb.) L.G.Adams, Austral. Syst. Bot. 21: 258 (2008)	1:75
	<b>Stellaria angustifolia</b> Hook. subsp. <b>angustifolia</b> , J. Bot. (Hooker) 1: 250 (1834)	
	<b>Stellaria angustifolia</b> Hook. subsp. <b>tenella</b> (Benth.) C.H.Mill. & J.G.West, J. Adelaide Bot. Gard. 25: 36 (2012)	
	<i>Stellaria caespitosa</i> Hook.f. = <b>Stellaria angustifolia</b> subsp. <b>angustifolia</b>	
	<b>Stellaria flaccida</b> Hook., Companion Bot. Mag. 1: 275 (1836)	1:71
	<i>Stellaria glauca</i> With. sensu Hooker (1860) = <b>Stellaria angustifolia</b> (misapplied in Tasmania)	
n i *	<i>Stellaria graminea</i> L., Sp. Pl. 1: 422 (1753)	1:72
i	<b>Stellaria media</b> (L.) Vill., Hist. Pl. Dauphiné (Villars) 3: 615 (1789)	1:71
	<b>Stellaria multiflora</b> Hook. subsp. <b>multiflora</b> , Companion Bot. Mag. 1: 275 (1836)	
	<b>Stellaria multiflora</b> Hook. subsp. <b>nebulosa</b> C.H.Mill. & J.G.West, J. Adelaide Bot. Gard. 25: 48 (2012)	
i	<b>Stellaria pallida</b> (Dumort.) Crép., Man. Fl. Belgique, ed. 2: 19 (1866)	1:72
	<i>Stellaria palustris</i> Retz. sensu Curtis (1956), Curtis & Morris (1975) = <b>Stellaria angustifolia</b> (misapplied in Tasmania)	1:71

	<b>Stellaria pungens</b> Brongn., Voy. Monde, Atlas: t.78 (1826)	1:71
n	<i>Vaccaria hispanica</i> (Mill.) Rauschert – previously listed as naturalised but insufficient evidence exists to support this	1:66
	<i>Vaccaria pyramidata</i> Medik. = <i>Vaccaria hispanica</i>	1:66
	<i>Vaccaria segetalis</i> (Neck.) Garcke ex Asch. = <i>Vaccaria hispanica</i>	
	<b>CASUARINACEAE</b>	FTO 67
e	<b>Allocasuarina crassa</b> L.A.S.Johnson, Fl. Australia 3: 194 (1989)	
e	<b>Allocasuarina duncanii</b> L.A.S.Johnson & D.I.Morris, Telopea 5: 793 (1994)	
	<b>Allocasuarina littoralis</b> (Salisb.) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 76 (1982)	3:644
e	<b>Allocasuarina monilifera</b> (L.A.S.Johnson) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 76 (1982)	3:644
	<b>Allocasuarina paludosa</b> (Sieber ex Spreng.) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 77 (1982)	3:644
	<b>Allocasuarina verticillata</b> (Lam.) L.A.S.Johnson, J. Adelaide Bot. Gard. 6: 79 (1982)	3:644
e	<b>Allocasuarina zephyrea</b> L.A.S.Johnson, Fl. Australia 3: 199 (1989)	
	<i>Casuarina bicuspidata</i> Benth. sensu Rodway (1903) = <b>Allocasuarina paludosa</b> (misapplied in Tasmania)	3:644
	<i>Casuarina distyla</i> Vent. sensu Rodway (1903) = <b>Allocasuarina monilifera</b> (misapplied in Tasmania)	3:644
	<i>Casuarina littoralis</i> Salisb. = <b>Allocasuarina littoralis</b>	3:644
	<i>Casuarina monilifera</i> L.A.S.Johnson = <b>Allocasuarina monilifera</b>	3:644
	<i>Casuarina paludosa</i> Sieber ex Spreng. = <b>Allocasuarina paludosa</b>	3:644
	<i>Casuarina quadrivalvis</i> Labill. = <b>Allocasuarina verticillata</b>	3:644
	<i>Casuarina stricta</i> Dryand. = <b>Allocasuarina verticillata</b>	3:644
	<i>Casuarina suberosa</i> Otto & A.Dietr. = <b>Allocasuarina littoralis</b>	3:644
	<b>CELASTRACEAE</b>	
i	<b>Euonymus europaeus</b> L., Sp. Pl. 1: 197 (1753)	
i	<b>Euonymus japonicus</b> Thunb., Nova Acta Regiae Soc. Sci. Upsal. 3: 208 (1780)	
i	<b>Maytenus magellanica</b> (Lam.) Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.). 2: 254 (1845)	
	<b>CHENOPODIACEAE</b>	FTO 98
	<i>Arthrocnemum arbuscula</i> (R.Br.) Moq. = <b>Tecticornia arbuscula</b>	3:577
?i	<b>Atriplex australasica</b> Moq., Chenop. Monogr. Enum. 59 (1840)	
t	<b>Atriplex billardierei</b> (Moq.) Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 215 (1853)	3:575
	<b>Atriplex cinerea</b> Poir., Encycl. (Lamarck) Suppl. 1: 471 (1811)	3:574
	<i>Atriplex halimus</i> L. sensu Brown (1810) = <b>Atriplex cinerea</b> (misapplied in Tasmania)	
	<i>Atriplex hastata</i> L. sensu Curtis (1967) = <b>Atriplex prostrata</b> (misapplied in Tasmania)	3:575
i	<b>Atriplex hortensis</b> L., Sp. Pl. 2: 1053 (1753)	3:576
	<b>Atriplex paludosa</b> R.Br. subsp. <b>paludosa</b> , Prodr. Fl. Nov. Holland. 406 (1810)	3:574
i	<b>Atriplex patula</b> L., Sp. Pl. 2: 1053 (1753)	3:575
	<i>Atriplex patula</i> L. var. <i>angustifolia</i> Rodway – a name of uncertain application	
	<i>Atriplex patula</i> L. var. <i>littoralis</i> Rodway – a name of uncertain application	
i	<b>Atriplex prostrata</b> Boucher ex DC., Fl. Franc. (DC. & Lamarck), ed. 3, 3: 387 (1805)	3:575
n ?i #	<b>Atriplex semibaccata</b> R.Br., Prodr. Fl. Nov. Holland. 406 (1810)	
	<b>Atriplex suberecta</b> I.Verd., Bothalia 6: 418 (1954)	
n i *	<i>Bassia scoparia</i> (L.) A.J.Scott, Feddes Repert. 89: 108 (1978)	
i	<b>Beta vulgaris</b> L. subsp. <b>maritima</b> (L.) Arcang., Comp. Fl. Ital. [Arcangeli]: 593 (1882)	3:573



i	<b>Chenopodium album</b> L., Sp. Pl. 1: 219 (1753)	3:571
	Chenopodium ambiguum R.Br. = <b>Chenopodium glaucum</b>	
	Chenopodium baccatum Labill. sensu Labillardiere (1806) = <b>Rhagodia candolleana</b> (misapplied in Tasmania)	
	Chenopodium capitatum (L.) Ambrosi sensu Buchanan (2009) = Chenopodium foliosum (misapplied in Tasmania)	
x	<b>Chenopodium erosum</b> R.Br., Prodr. Fl. Nov. Holland. 407 (1810)	
n	Chenopodium foliosum (Moench) Asch. – previously listed as naturalised but insufficient evidence exists to support this	
	Chenopodium furfuraceum Moq. = <b>Rhagodia candolleana</b> subsp. <b>candolleana</b>	
?i	<b>Chenopodium glaucum</b> L., Sp. Pl. 1: 220 (1753)	3:572
	Chenopodium glaucum L. subsp. ambiguum (R.Br.) Murr & Thell. ex Thell. = <b>Chenopodium glaucum</b>	
i	<b>Chenopodium murale</b> L., Sp. Pl. 1: 219 (1753)	3:572
	Chenopodium pumilio R.Br. = <b>Dysphania pumilio</b>	3:573
i	<b>Chenopodium vulvaria</b> L., Sp. Pl. 1: 220 (1753)	
i	<b>Dysphania glomulifera</b> (Nees) Paul G.Wilson subsp. <b>glomulifera</b> , Nuytsia 4: 183 (1983)	
i	<b>Dysphania pumilio</b> (R.Br.) Mosyakin & Clemants, Ukrayins'k. Bot. Zhurn. 59: 382 (2002)	3:573
	<b>Einadia nutans</b> (R.Br.) A.J.Scott subsp. <b>nutans</b> , Feddes Repert. 89: 3 (1978)	3:570
	Rhagodia baccata (Labill.) Moq. sensu Curtis (1967) = <b>Rhagodia candolleana</b> (misapplied in Tasmania)	3:570
	Rhagodia billardierei R.Br. sensu Brown (1810) = <b>Rhagodia candolleana</b> (misapplied in Tasmania)	
	Rhagodia billardierei R.Br. var. congesta (Hook.f.) Benth. = <b>Chenopodium murale</b>	
	<b>Rhagodia candolleana</b> Moq. subsp. <b>candolleana</b> , Chenop. Monogr. Enum. 10 (1840)	3:570
	Rhagodia nutans R.Br. = <b>Einadia nutans</b>	3:570
	Salicornia arbuscula R.Br. = <b>Tecticornia arbuscula</b>	
	Salicornia australis Sol. ex Benth. nom. illeg. = <b>Sarcocornia quinqueflora</b> subsp. <b>quinqueflora</b>	
	Salicornia blackiana Ulbr. = <b>Sarcocornia blackiana</b>	3:579
	Salicornia indica R.Br. sensu Brown (1810) = <b>Sarcocornia quinqueflora</b> (misapplied in Tasmania)	
	Salicornia quinqueflora Bunge ex Ung.-Sternb. = <b>Sarcocornia quinqueflora</b>	3:578
	<b>Salsola australis</b> R.Br., Prodr. Fl. Nov. Holland. 411 (1810)	
	Salsola kali L. sensu Buchanan (2005) = <b>Salsola australis</b> (misapplied in Tasmania)	
	Salsola tragus L. sensu Buchanan (2009) = <b>Salsola australis</b> (misapplied in Tasmania)	
	<b>Sarcocornia blackiana</b> (Ulbr.) A.J.Scott, Bot. J. Linn. Soc. 75: 369 (1978)	3:579
	<b>Sarcocornia quinqueflora</b> (Bunge ex Ung.-Sternb.) A.J.Scott subsp. <b>quinqueflora</b> , Bot. J. Linn. Soc. 75: 368 (1978)	3:578
	<b>Sarcocornia quinqueflora</b> (Bunge ex Ung.-Sternb.) A.J.Scott subsp. <b>tasmanica</b> Paul G.Wilson, Nuytsia 3: 74 (1980)	3:578
	Sclerostegia arbuscula (R.Br.) Paul G.Wilson = <b>Tecticornia arbuscula</b>	3:577
	<b>Suaeda australis</b> (R.Br.) Moq., Ann. Sci. Nat. (Paris) 23: 318 (1831)	3:580
i	<b>Suaeda maritima</b> (L.) Dumort. subsp. <b>maritima</b> , Fl. Belg. (Dumortier): 22 (1827)	
	<b>Tecticornia arbuscula</b> (R.Br.) K.A.Sheph. & Paul G.Wilson, Austral. Syst. Bot. 20: 325 (2007)	3:577
	<b>Threlkeldia diffusa</b> R.Br., Prodr. Fl. Nov. Holland. 410 (1810)	3:577

**CISTACEAE**

- i **Cistus creticus** L., Sp. Pl., ed. 2: 738 (1762)
- i # **Cistus inflatus** Pourr. ex Demoly, Acta Bot. Gallica 144: 42 (1998)  
Cistus psilosepalus Sweet sensu Buchanan (2004) = **Cistus inflatus** (misapplied in Tasmania)

**CLUSIACEAE (GUTTIFERAE)**

- Ascyrum humifusum Labill. = **Hypericum pusillum**
- Ascyrum involutum Labill. = **Hypericum gramineum**
- i **Hypericum androsaemum** L., Sp. Pl. 2: 784 (1753) 1:82
- i **Hypericum calycinum** L., Mant. Pl. 1: 106 (1767) 1:82
- Hypericum gramineum** G.Forst., Fl. Ins. Austr. 53 (1786) 1:81
- n Hypericum humifusum L. – previously listed as naturalised but insufficient evidence exists to support this
- Hypericum japonicum** Thunb., Fl. Jap. (Thunberg): 295 (1784) 1:81
- i **Hypericum perforatum** L. subsp. **veronense** (Schränk) H.Lindb., Öfvers. Finska Vetensk.-Soc. Förh. 48: 73 (1906) 1:81
- i # t **Hypericum pulchrum** L., Sp. Pl. 2: 786 (1753)
- t **Hypericum pusillum** Choisy, Prodr. Monogr. Hyperic. 50 (1821)
- i **Hypericum tetrapterum** Fr. var. **tetrapterum**, Novit. Fl. Suec. Alt. 236 (1828)

**CONVOLVULACEAE**

FTO 111

- Calystegia marginata** R.Br., Prodr. Fl. Nov. Holland. 483 (1810)
- Calystegia sepium** (L.) R.Br., Prodr. Fl. Nov. Holland. 483 (1810) 3:498
- i **Calystegia silvatica** (Kit.) Griseb., Spic. Fl. Rumel. 1: 74 (1843) 3:498
- Calystegia soldanella** (L.) Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 4: 184 (1819) 3:499
- Convolvulus angustissimus** R.Br. subsp. **angustissimus**, Prodr. Fl. Nov. Holland. 482 (1810) 3:497
- i **Convolvulus arvensis** L., Sp. Pl. 1: 153 (1753) 3:497
- Convolvulus erubescens Sims sensu Curtis (1967) = **Convolvulus angustissimus** subsp. **angustissimus** (misapplied in Tasmania) 3:497
- Convolvulus sepium L. sensu Bentham (1868) = **Calystegia sepium**
- Convolvulus soldanella L. sensu Bentham (1868) = **Calystegia soldanella**
- Dichondra repens** J.R.Forst. & G.Forst., Char. Gen. Pl. 40: t.20 (1775) 3:499
- i **Ipomoea indica** (Burm.) Merr., Interpr. Herb. Amboin. 445 (1917)
- Wilsonia backhousei** Hook.f., London J. Bot. 6: 275 (1847) 3:500
- Wilsonia humilis** R.Br., Prodr. Fl. Nov. Holland. 490 (1810) 3:500
- Wilsonia rotundifolia** Hook., Icon. Pl. 5: t.160 (1842)

**CORNACEAE**

FTO 136

- i # t **Griselinia littoralis** (Raoul) Raoul, Choix Pl. Nouv.-Zél. 22, t.19 (1846)

**CRASSULACEAE**

- i **Aeonium arboreum** (L.) Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 1: 185 (1840)
- i # **Aeonium haworthii** Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 1: 193 (1840)
- i **Cotyledon orbiculata** L., Sp. Pl. 1: 429 (1753)
- Crassula closiana** (Gay) Reiche, Fl. Chile [Reiche] 2: 369 (1897)
- Crassula decumbens** Thunb. var. **decumbens**, Prodr. Pl. Cap. 1: 54 (1794) 1:185
- Crassula exserta** (Reader) Ostenf., Dansk Bot. Ark. 2: 47 (1918) 1:184

	<b>Crassula helmsii</b> (Kirk) Cockayne, Trans. New Zealand Inst. 39: 349 (1907)	1:185
	<i>Crassula macrantha</i> (Hook.f.) Diels & E.Pritz. = <b>Crassula decumbens</b>	1:185
t	<b>Crassula moschata</b> G.Forst., Commentat. Soc. Regiae Sci. Gott. 9: 26 (1789)	
i	<b>Crassula multicava</b> Lem. subsp. <b>multicava</b> , Ill. Hort. 9: 40 (1862)	
i #	<b>Crassula muscosa</b> L. var. <b>muscosa</b> , Pl. Rar. Afr. 10 (1760)	
n i	<b>Crassula natans</b> Thunb. var. <b>minus</b> (Eckl. & Zeyh.) G.D.Rowley, Cact. Succ. J. Gr. Brit. 40: 53 (1978)	
	<i>Crassula pedicellosa</i> (F.Muell.) Ostenf. = <b>Crassula closiana</b>	
	<b>Crassula peduncularis</b> (Sm.) F.Meigen, Bot. Jahrb Syst. 17: 239 (1893)	1:184
	<b>Crassula sieberiana</b> (Schult. & Schult.f.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 618 (1917)	1:184
	<i>Crassula sieberiana</i> (Schult. & Schult.f.) Druce subsp. <b>tetramera</b> Toelken = <b>Crassula tetramera</b>	
i #	<b>Crassula tetragona</b> L. subsp. <b>robusta</b> (Toelken) Toelken, J. S. African Bot. 41: 122 (1975)	
	<b>Crassula tetramera</b> (Toelken) A.P.Druce & Sykes, New Zealand J. Bot. 25: 128 (1987)	
i	<b>Sedum acre</b> L., Sp. Pl. 1: 432 (1753)	1:186
i	<b>Sedum album</b> L., Sp. Pl. 1: 432 (1753)	
i	<b>Sedum forsterianum</b> Sm., Engl. Bot. 26: t.1802 (1808)	
i	<b>Sedum praealtum</b> A. DC., Mém. Soc. Phys. Genève 11: 445 (1847)	
	<i>Sedum reflexum</i> L. sensu Buchanan (2007) = <b>Sedum rupestre</b> (misapplied in Tasmania)	
i	<b>Sedum rupestre</b> L., Sp. Pl. 1: 431 (1753)	
i	<b>Sedum sediforme</b> (Jacq.) Pau, Acta y Memorias Prim. Congr. Nat. Esp. 246 (1909)	
i t	<b>Sedum sexangulare</b> L., Sp. Pl. 1: 432 (1753)	1:186
	<i>Tillaea exerta</i> Reader = <b>Crassula exserta</b>	
	<i>Tillaea macrantha</i> Hook.f. = <b>Crassula decumbens</b>	
	<i>Tillaea purpurata</i> Hook.f. = <b>Crassula peduncularis</b>	
	<i>Tillaea recurva</i> (Hook.f.) Hook.f. = <b>Crassula helmsii</b>	
	<i>Tillaea sieberiana</i> Schult. & Schult.f. = <b>Crassula sieberiana</b>	
	<i>Tillaea verticillaris</i> DC. = <b>Crassula sieberiana</b>	
	<b>CUCURBITACEAE</b>	FTO 65
i	<b>Cucumis myriocarpus</b> Naudin, Ann. Sci. Nat., Bot., sér. 4, 11: 22 (1859)	
i #	<b>Ecballium elaterium</b> (L.) A.Rich., Dict. Class. Hist. Nat. [Bory] 6: 19 (1824)	
	<i>Sicyos angulatus</i> L. sensu Curtis (1963) = <b>Sicyos australis</b> (misapplied in Tasmania)	2:237
	<b>Sicyos australis</b> Endl., Prodr. Fl. Norfolk. 67 (1833)	2:237
	<b>CUNONIACEAE</b>	FTO 72
e	<b>Anodopetalum biglandulosum</b> (A.Cunn. ex Hook.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 148 (1856)	1:180
	<b>Bauera rubioides</b> Andrews, Bot. Repos. 3: t.198 (1801)	1:180
	<b>CUSCUTACEAE</b>	FTO 111
i	<b>Cuscuta epithymum</b> (L.) L., Syst. Veg., ed. 13: 140 (1774)	3:501
n i x	<b>Cuscuta suaveolens</b> Ser., Ann. Sci. Phys. Nat. Lyon 3: 519 (1840)	
	<b>Cuscuta tasmanica</b> Engelm., Trans. Acad. Sci. St. Louis 1: 512 (1859)	3:501
	<b>DILLENIACEAE</b>	FTO 92
	<i>Dillenia procumbens</i> Labill. = <b>Hibbertia procumbens</b>	
	<b>Hibbertia acicularis</b> (Labill.) F.Muell., Pl. Victoria 1: 17 (1862)	1:23

	Hibbertia angustifolia (R.Br. ex DC.) Benth. = <b>Hibbertia procumbens</b>	
	<b>Hibbertia appressa</b> Toelken, J. Adelaide Bot. Gard. 19: 54 (2000)	
	Hibbertia aspera DC. sensu Curtis & Morris (1975) = <b>Hibbertia hirticalyx</b> (misapplied in Tasmania)	1:23
e	<b>Hibbertia basaltica</b> A.M.Buchanan & Schah., Muelleria 22: 105 (2005)	
	Hibbertia billardierei F.Muell. nom. illeg. = <b>Hibbertia empetrifolia</b>	
	Hibbertia billardierei F.Muell. var. monadelpha F.Muell. ex Benth. = <b>Hibbertia hirticalyx</b>	
	Hibbertia billardierei F.Muell. var. obovata R.Br. ex Benth. = <b>Hibbertia hirticalyx</b>	
	<b>Hibbertia calycina</b> (DC.) N.A.Wakef., Vict. Naturalist 72: 122 (1955)	
	Hibbertia densiflora (Hook.f.) F.Muell. = <b>Hibbertia sericea</b>	
	<b>Hibbertia dispar</b> Toelken, J. Adelaide Bot. Gard. 26: 31-69 (2013)	
	<b>Hibbertia empetrifolia</b> (DC.) Hoogland subsp. <b>empetrifolia</b> , Kew Bull. 29: 155 (1974)	1:23
	<b>Hibbertia ericifolia</b> Hook.f. subsp. <b>ericifolia</b> , Bot. Antarct. Voy. III. (Fl. Tasman.) I: 14 (1855) (as H. ericaefolia)	
	Hibbertia fasciculata R.Br. ex DC. sensu Curtis & Morris (1975) = <b>Hibbertia prostrata</b>	1:24
	<b>Hibbertia hirsuta</b> (Hook.) Benth., Fl. Austral. 1: 26 (1863)	1:22
	<b>Hibbertia hirticalyx</b> Toelken, J. Adelaide Bot. Gard. 18: 146 (1998)	1:23
	Hibbertia linearis R.Br. var. obtusifolia (DC.) Benth. = <b>Hibbertia obtusifolia</b>	
x	<b>Hibbertia obtusifolia</b> DC., Syst. Nat. [Candolle] 1: 429 (1817)	1:24
	<b>Hibbertia procumbens</b> (Labill.) DC., Syst. Nat. [Candolle] 1: 427 (1817)	1:24
	<b>Hibbertia prostrata</b> Hook., J. Bot. (Hooker) 1: 246 (1834)	1:24
	<b>Hibbertia riparia</b> (R.Br. ex DC.) Hoogland, Kew Bull. 29: 155 (1974)	1:22
	<b>Hibbertia rufa</b> N.A.Wakef., Vict. Naturalist 72: 119 (1955)	1:23
	<b>Hibbertia sericea</b> (R.Br. ex DC.) Benth. var. <b>sericea</b> , Fl. Austral. 1: 26 (1863)	1:22
	<b>Hibbertia serpyllifolia</b> R.Br. ex DC., Syst. Nat. [Candolle] 1: 430 (1817)	1:24
	Hibbertia stricta (DC.) F.Muell. = <b>Hibbertia riparia</b>	
	Hibbertia stricta (DC.) F.Muell. var. canescens = <b>Hibbertia riparia</b>	
	Hibbertia stricta (DC.) F.Muell. var. glabriuscula = <b>Hibbertia riparia</b>	
	<b>Hibbertia virgata</b> R.Br. ex DC., Syst. Nat. [Candolle] 1: 428 (1817)	1:24
	Pleurandra acicularis Labill. = <b>Hibbertia acicularis</b>	
	Pleurandra hirsuta Hook. = <b>Hibbertia hirsuta</b>	
	Pleurandra ovata Labill. = <b>Hibbertia empetrifolia</b> and <b>Hibbertia appressa</b>	
	Pleurandra riparia R.Br. ex DC. (and vars.) = <b>Hibbertia riparia</b>	
	Pleurandra sericea R.Br. ex DC. = <b>Hibbertia sericea</b>	
	<b>DIPSACACEAE</b>	FTO 134
i	<b>Dipsacus fullonum</b> L., Sp. Pl. 1: 97 (1753)	2:280
	Dipsacus sylvestris Huds. = <b>Dipsacus fullonum</b>	2:280
i	<b>Scabiosa atropurpurea</b> L., Sp. Pl. 1: 100 (1753)	2:280
i	<b>Scabiosa farinosa</b> Coss., Ill. Fl. Atlant. 6: 58, t. 134. (1893)	
	<b>DONATIACEAE</b>	FTO 126
t	<b>Donatia novae-zelandiae</b> Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 81 (1852)	2:397
	<b>DROSERACEAE</b>	FTO 96
	<b>Drosera arcturi</b> Hook., J. Bot. (Hooker) 1: 247 (1834)	1:187
	<b>Drosera auriculata</b> Backh. ex Planch., Ann. Sci. Nat., Bot., sér. 3, 9: 295 (1848)	1:188

	<b>Drosera binata</b> Labill., Nov. Holl. Pl. 1: 78 t.105 (1805)	1:188
	<i>Drosera foliosa</i> Hook.f. ex Planch. nom. illeg. = <b>Drosera hookeri</b>	
	<b>Drosera glanduligera</b> Lehm., Nov. Stirp. Pug. [Lehmann] 8: 37 (1844)	1:187
	<b>Drosera gracilis</b> Hook.f. ex Planch., Ann. Sci. Nat., Bot., sér. 3, 9:297 (1848)	1:188
	<b>Drosera hookeri</b> R.P.Gibson, B.J. Conn & Conran, J. Adelaide Bot. Gard. 24: 41 (2010)	
	<b>Drosera macrantha</b> Endl., Enum. Pl. [Endlicher] 6 (1837)	1:189
	<i>Drosera macrantha</i> Endl. subsp. <i>planchonii</i> (Hook.f. ex Planch.) N.G.Marchant = <b>Drosera macrantha</b>	1:189
	<i>Drosera menziesii</i> R.Br. ex DC. sensu Rodway (1903) = <b>Drosera macrantha</b> (misapplied in Tasmania)	
	<i>Drosera menziesii</i> R.Br. ex DC. var. <i>albiflora</i> Benth. = <b>Drosera macrantha</b>	
e	<b>Drosera murfetii</b> Lowrie & Conran, J. Adelaide Bot. Gard. 27:17 (2014)	
	<b>Drosera peltata</b> Thunb., <i>Drosera</i> 7 (1797)	1:188
	<i>Drosera peltata</i> Thunb. subsp. <i>auriculata</i> (Backh. ex Planch.) Conn = <b>Drosera auriculata</b>	1:188
	<i>Drosera peltata</i> Thunb. var. <i>foliosa</i> (Hook.f. ex Planch.) Benth. = <b>Drosera hookeri</b>	
	<i>Drosera peltata</i> Thunb. var. <i>gracilis</i> (Hook.f. ex Planch.) Benth. = <b>Drosera gracilis</b>	
	<i>Drosera planchonii</i> Hook.f. = <b>Drosera macrantha</b>	1:189
	<b>Drosera pygmaea</b> DC., Prodr. [A. P. de Candolle] 1: 317 (1824)	1:187
	<b>Drosera spatulata</b> Labill. var. <i>spatulata</i> , Nov. Holl. Pl. 1: 79 t.106 fig.1 (1805)	1:187
	<b>ELAEOCARPACEAE</b>	FTO 71
e	<b>Aristotelia peduncularis</b> (Labill.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 52 (1855)	1:91
	<i>Elaeocarpus cyaneus</i> Sims = <b>Elaeocarpus reticulatus</b>	
	<i>Elaeocarpus peduncularis</i> Labill. = <b>Aristotelia peduncularis</b>	
	<b>Elaeocarpus reticulatus</b> Sm., Cycl. 12, no.6 (1809)	1:91
	<b>ELATINACEAE</b>	FTO 76
	<i>Elatine americana</i> (Pursh) Arn. sensu Bentham (1863) = <b>Elatine gratioloides</b> (misapplied in Tasmania)	
	<b>Elatine gratioloides</b> A.Cunn., Ann. Nat. Hist. 4: 26 (1840)	1:80
	<b>EPACRIDACEAE</b>	FTO 105
	<b>Acrothamnus hookeri</b> (Sond.) Quinn, Austral. Syst. Bot. 18: 452 (2005)	2:438
	<b>Acrothamnus montanus</b> (R.Br.) Quinn, Austral. Syst. Bot. 18: 452 (2005)	2:433
	<b>Acrotriche affinis</b> DC., Prodr. [A. P. de Candolle] 7: 757 (1839)	
	<b>Acrotriche cordata</b> (Labill.) R.Br., Prodr. Fl. Nov. Holland. 548 (1810)	
	<i>Acrotriche patula</i> sensu Hooker (1860) = <b>Acrotriche serrulata</b>	
	<b>Acrotriche serrulata</b> (Labill.) R.Br., Prodr. Fl. Nov. Holland. 547 (1810)	2:440
e	<b>Androstoma verticillata</b> (Hook.f.) Quinn, Austral. Syst. Bot. 18: 450 (2005)	2:439
e	<b>Archeria comberi</b> Melville, Kew Bull. 12: 389 (1957)	2:453
e	<b>Archeria eriocarpa</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 263 t.80B (1857)	2:452
e	<b>Archeria hirtella</b> (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 263 t.81 (1857)	2:452
e	<b>Archeria minor</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 264 (1857)	2:452
e	<b>Archeria serpyllifolia</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 263 t.80A (1857)	2:452
	<i>Archeria serpyllifolia</i> Hook.f. var. <i>minor</i> (Hook.f.) Benth. = <b>Archeria minor</b>	2:452
	<b>Astroloma humifusum</b> (Cav.) R.Br., Prodr. Fl. Nov. Holland. 538 (1810)	2:424
	<b>Astroloma pinifolium</b> (R.Br.) Benth., Fl. Austral. 4: 159 (1868)	2:424

	<b>Brachyloma ciliatum</b> (R.Br.) Benth., Fl. Austral. 4: 173 (1868)	2:432
	Brachyloma ciliatum (R.Br.) Benth. var. intermedium Rodway = <b>Brachyloma ciliatum</b>	
	Brachyloma daphnoides Benth. sensu Bentham (1868) mistakenly attributed to Tasmania	
	<b>Brachyloma depressum</b> (F.Muell.) Benth., Fl. Austral. 4: 173 (1868)	2:432
	Cyathodes abietina (Labill.) R.Br. = <b>Leptecophylla abietina</b>	2:427
	Cyathodes acerosa R.Br. ex Roem. & Schult. = Leptecophylla juniperina	
	Cyathodes adscendens Hook.f. = <b>Planocarpa petiolaris</b>	
	Cyathodes dealbata R.Br. = <b>Montitega dealbata</b>	2:427
	Cyathodes disticha Labill. = <b>Trochocarpa disticha</b>	
	Cyathodes divaricata Hook.f. = <b>Leptecophylla divaricata</b>	2:428
e	<b>Cyathodes glauca</b> Labill., Nov. Holl. Pl. 1: 57 t.81 (1805)	2:426
	Cyathodes juniperina (J.R.Forst. & G.Forst.) Druce = Leptecophylla juniperina	2:427
	Cyathodes macrantha Hook.f. = <b>Cyathodes straminea</b>	
	Cyathodes nitida Jarman = <b>Planocarpa nitida</b>	
	Cyathodes oxycedrus (Labill.) R.Br. = <b>Leptecophylla oxycedrus</b>	
	Cyathodes parvifolia R.Br. = <b>Leptecophylla parvifolia</b>	2:428
	Cyathodes pendulosa Jarman = <b>Leptecophylla pendulosa</b>	
	Cyathodes petiolaris (DC.) Druce = <b>Planocarpa petiolaris</b>	2:426
e	<b>Cyathodes platystoma</b> C.M.Weiller, Austral. Syst. Bot. 9: 502 (1996)	
e	<b>Cyathodes straminea</b> R.Br., Prodr. Fl. Nov. Holland. 539 (1810)	2:426
	Cyathodes straminea var. macrantha (Hook.f.) Rodway = <b>Cyathodes straminea</b>	
	Cyathodes sulcata Mihaich = <b>Planocarpa sulcata</b>	
	Cystanthe acerosa (Lindl.) F.Muell. = <b>Richea acerosa</b>	
	Cystanthe dracophylla (Hook.f.) Kuntze = <b>Richea dracophylla</b>	
	Cystanthe gunnii (Hook.f.) Kuntze = <b>Richea gunnii</b>	
	Cystanthe milliganii (Hook.f.) F.Muell. = <b>Richea milliganii</b>	
	Cystanthe pandanifolia (Hook.f.) Kuntze = <b>Richea pandanifolia</b>	
	Cystanthe procera F.Muell. = <b>Richea procera</b>	
	Cystanthe scoparia (Hook.f.) Kuntze = <b>Richea scoparia</b>	
	Cystanthe sprengeioides R.Br. = <b>Richea sprengeioides</b>	
	Decaspora cunninghamii DC. = <b>Trochocarpa cunninghamii</b>	
	Decaspora disticha (Labill.) R.Br. = <b>Trochocarpa disticha</b>	
	Decaspora gunnii Hook.f. = <b>Trochocarpa gunnii</b>	
	Decaspora thymifolia R.Br. = <b>Trochocarpa thymifolia</b>	
e	<b>Dracophyllum milliganii</b> Hook.f., Icon. Pl. 9: t.845 (1852)	2:461
e	<b>Dracophyllum minimum</b> F.Muell., Fragm. (Mueller) 1: 39 (1859)	2:461
e	<b>Epacris acuminata</b> Benth., Fl. Austral. 4: 240 (1868)	2:451
e	<b>Epacris apsleyensis</b> Crowden, Pap. & Proc. Roy. Soc. Tasmania 120: 17 (1986)	
e	<b>Epacris barbata</b> Melville, Kew Bull. 7: 176 (1952)	2:450
	Epacris campanulata Lodd. ex DC. = <b>Epacris impressa</b>	
e	<b>Epacris cerasicollina</b> Crowden, Muelleria 25: 124 (2007)	
	Epacris ceriflora Graham = <b>Epacris impressa</b>	
	Epacris cerinthoides Labill. = <b>Prionotes cerinthoides</b>	
e	<b>Epacris corymbiflora</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 261 (1857)	2:447
e	<b>Epacris curtisiae</b> Jarman, Pap. & Proc. Roy. Soc. Tasmania 122: 115 (1988)	

e	<b>Epacris exserta</b> R.Br., Prodr. Fl. Nov. Holland. 551 (1810)	2:448
	Epacris exserta var. virgata (Hook.f.) Benth. = <b>Epacris virgata</b>	
e	<b>Epacris franklinii</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 261 (1857)	2:448
e	<b>Epacris glabella</b> Jarman, Aspects of Tasmanian Botany: 100 (1991)	
e	<b>Epacris grandis</b> Crowden, Pap. & Proc. Roy. Soc. Tasmania 120: 19 (1986)	
e	<b>Epacris graniticola</b> Crowden, Muelleria 25: 126 (2007)	
	<b>Epacris gunnii</b> Hook.f., London J. Bot. 6: 272 (1847)	2:451
e	<b>Epacris heteronema</b> Labill. var. <b>gigantea</b> F.Muell., Fragm (Mueller) 4: 127 (1864)	
e	<b>Epacris heteronema</b> Labill. var. <b>heteronema</b> , Nov. Holl. Pl. I: 42 t.56 (1805)	
	Epacris heteronema var. planifolia = <b>Epacris heteronema</b> (Tasmanian material)	
	Epacris hirtella Hook.f. = <b>Archeria hirtella</b>	
	<b>Epacris impressa</b> Labill., Nov. Holl. Pl. I: 43 t.58 (1805)	2:446
	Epacris impressa f. ceriflora (Graham) Siebert & Voss = <b>Epacris impressa</b>	
	Epacris impressa f. diemenica Gand. = <b>Epacris impressa</b>	
	Epacris impressa f. lucida Gand. = <b>Epacris impressa</b>	
	Epacris impressa f. milliganii Gand. = <b>Epacris impressa</b>	
	Epacris impressa f. ruscifolia (R.Br.) Siebert & Voss = <b>Epacris impressa</b>	
	Epacris impressa var. campanulata (Lodd. ex DC.) Hook.f. = <b>Epacris impressa</b>	
	Epacris impressa var. ceriflora (Graham) Rodway = <b>Epacris impressa</b>	
	Epacris impressa var. nivea Hook.f. = <b>Epacris impressa</b>	
	Epacris impressa var. ovata Benth. = <b>Epacris impressa</b>	
	Epacris impressa var. ruscifolia (R.Br.) Rodway = <b>Epacris impressa</b>	
	Epacris impressa var. variabilis (Lodd. ex Paxton) Hook.f. = <b>Epacris impressa</b>	
	<b>Epacris lanuginosa</b> Labill., Nov. Holl. Pl. I: 42 t.57 (1805)	2:448
e	<b>Epacris limbata</b> K.J.Williams & F.Duncan, Aspects of Tasmanian Botany: 95 (1991)	
e	<b>Epacris marginata</b> Melville, Kew Bull. 7: 175 (1952)	2:449
	Epacris micranthera F.Muell. = <b>Archeria serpyllifolia</b>	
	Epacris microphylla Hook.f. nom. illeg., sensu Hooker (1847) = <b>Archeria minor</b>	
	Epacris microphylla R.Br. sensu Bentham (1868), Rodway (1903) p.p. = <b>Epacris gunnii</b> (Tasmanian material)	
	Epacris microphylla var. gunnii (Hook.f.) Benth. = <b>Epacris gunnii</b>	
e	<b>Epacris moscaliana</b> Crowden, Muelleria 25: 127 (2007)	
e	<b>Epacris mucronulata</b> R.Br., Prodr. Fl. Nov. Holland. 552 (1810)	2:448
	Epacris mucronulata sensu Hooker (1860) p.p. = <b>Epacris acuminata</b>	
e	<b>Epacris myrtifolia</b> Labill., Nov. Holl. Pl. I: 41 t.55 (1805)	2:447
	Epacris myrtifolia var. corymbiflora (Hook.f.) Rodway = <b>Epacris corymbiflora</b>	
e	<b>Epacris navicularis</b> Jarman, Pap. & Proc. Roy. Soc. Tasmania 112: 3 (1978)	
	<b>Epacris obtusifolia</b> Sm., Exot. Bot. 1: 77 t.40 (1804)	2:447
	<b>Epacris paludosa</b> R.Br., Prodr. Fl. Nov. Holland. 551 (1810)	2:449
	<b>Epacris petrophila</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) I: 261 (1857)	2:446
	Epacris ruscifolia R.Br. = <b>Epacris impressa</b>	
e	<b>Epacris serpyllifolia</b> R.Br., Prodr. Fl. Nov. Holland. 551 (1810)	2:450
	Epacris serpyllifolia var. squarrosa (Hook.f.) Benth. = <b>Epacris tasmanica</b>	
	Epacris squarrosa Hook.f. = <b>Epacris tasmanica</b>	2:450
e	<b>Epacris stuartii</b> Stapf, Bull. Misc. Inform. Kew 1910: 217 (1910)	2:449

e	<b>Epacris tasmanica</b> W.M.Curtis, Taxon 18: 244 (1969)	2:450
	Epacris variabilis Lodd. ex Paxton nom. illeg., non Courtois (1833) = <b>Epacris impressa</b>	
e	<b>Epacris virgata</b> Hook.f., London J. Bot. 6: 271 (1847)	2:448
e	<b>Leptecophylla abietina</b> (Labill.) C.M.Weiller, Muelleria 12: 211 (1999)	2:427
e	<b>Leptecophylla divaricata</b> (Hook.f.) C.M.Weiller, Muelleria 12: 207 (1999)	2:428
n	Leptecophylla juniperina (J.R.Forst. & G.Forst.) C.M.Weiller subsp. juniperina sensu Curtis & Morris (1975), de Salas & Baker (2016) = <b>Leptecophylla oxycedrus</b> and <b>L. pogonocalyx</b> subsp. <b>decipiens</b> (misapplied in Tasmania)	2:427
n	Leptecophylla juniperina (J.R.Forst. & G.Forst.) C.M.Weiller subsp. oxycedrus (Labill.) C.M.Weiller = <b>Leptecophylla oxycedrus</b>	
n	Leptecophylla juniperina (J.R.Forst. & G.Forst.) C.M.Weiller subsp. parvifolia (R.Br.) C.M.Weiller = <b>Leptecophylla parvifolia</b>	
n	<b>Leptecophylla oxycedrus</b> (Labill.) Jarman, Swainsona 31: 1	
n e	<b>Leptecophylla parvifolia</b> (R.Br.) Jarman, Swainsona 31: 7	2:428
e	<b>Leptecophylla pendulosa</b> (Jarman) C.M.Weiller, Muelleria 12: 209 (1999)	
n e	<b>Leptecophylla pogonocalyx</b> C.M.Weiller subsp. <b>pogonocalyx</b> , Muelleria 12: 206 (1999)	
n e	<b>Leptecophylla pogonocalyx</b> C.M.Weiller subsp. <b>decipiens</b> Jarman, Swainsona 31: 13 (2017)	
	<b>Leucopogon affinis</b> R.Br., Prodr. Fl. Nov. Holland: 541 (1810)	2:435
	<b>Leucopogon australis</b> R.Br., Prodr. Fl. Nov. Holland. 541 (1810)	2:436
	Leucopogon ciliatus A.Cunn. ex DC. = <b>Leucopogon collinus</b>	
	Leucopogon ciliatus var. [alpha] Hook.f. = <b>Leucopogon collinus</b>	
	Leucopogon ciliatus var. [beta] Hook.f. = <b>Leucopogon oreophilus</b> and <b>Leucopogon pilifer</b>	
	<b>Leucopogon collinus</b> (Labill.) R.Br., Prodr. Fl. Nov. Holland. 543 (1810)	2:436
	Leucopogon collinus var. billardierei DC. = <b>Leucopogon collinus</b>	
	Leucopogon collinus var. brownii DC. = <b>Leucopogon collinus</b>	
	<b>Leucopogon ericoides</b> (Sm.) R.Br., Prodr. Fl. Nov. Holland. 543 (1810)	2:438
	Leucopogon ericoides var. [alpha] Hook.f. = <b>Leucopogon ericoides</b>	
	Leucopogon ericoides var. [beta] Hook.f. = <b>Leucopogon ericoides</b>	
	<b>Leucopogon esquamatus</b> R.Br., Prodr. Fl. Nov. Holland. 546 (1810)	2:439
	<b>Leucopogon fraseri</b> A.Cunn., Ann. Nat. Hist. 2: 47 (1839)	2:439
	Leucopogon hookeri Sond. = <b>Acrothamnus hookeri</b>	2:438
	Leucopogon lanceolatus (Sm.) R.Br. var. lanceolatus nom. illeg. = <b>Leucopogon affinis</b>	2:435
	Leucopogon milliganii (F.Muell.) Rodway = <b>Androstoma verticillata</b>	2:439
	Leucopogon montanus (R.Br.) J.H.Willis = <b>Acrothamnus montanus</b>	2:433
	Leucopogon obtusatus nom. illeg., sensu Hooker (1847) = <b>Acrothamnus hookeri</b>	2:438
e	<b>Leucopogon oreophilus</b> J.M.Powell, Aspects of Tasmanian Botany: 108 (1991)	
	<b>Leucopogon parviflorus</b> (Andrews) Lindl., Edwards's Bot. Reg. 18: t.1560 (1832)	2:435
	<b>Leucopogon pilifer</b> N.A.Wakef., Vict. Naturalist 73: 58 (1956)	
	Leucopogon richei (Labill.) R.Br. = <b>Leucopogon parviflorus</b>	
	Leucopogon richei var. [gamma] Hook.f. = <b>Leucopogon affinis</b>	
	Leucopogon richei var. parviflorus (Andrews) Hook.f. = <b>Leucopogon parviflorus</b>	
	Leucopogon rufus Lindl. sensu Bentham (1868) recorded in error	
	Leucopogon stuartii F.Muell. ex Sond. = <b>Leucopogon fraseri</b>	2:439
	Leucopogon trichocarpus (Labill.) R.Br. = <b>Leucopogon ericoides</b>	
	<b>Leucopogon virgatus</b> (Labill.) R.Br. var. <b>brevifolius</b> Benth., Fl. Austral. 4: 202 (1868)	2:438



	<b>Leucopogon virgatus</b> (Labill.) R.Br. var. <b>virgatus</b> , Prodr. Fl. Nov. Holland. 543 (1810)	2:438
	Lissanthe ciliata R.Br. = <b>Brachyloma ciliatum</b>	
	Lissanthe daphnoides (Sm.) R.Br. sensu Brown (1810), Hooker (1860) attributed in error	
	Lissanthe montana R.Br. = <b>Acrothamnus montanus</b>	2:433
	<b>Lissanthe strigosa</b> (Sm.) R.Br. subsp. <b>subulata</b> (R.Br.) J.M.Powell, Telopea 5: 741 (1994)	2:433
	<b>Monotoca elliptica</b> (Sm.) R.Br., Prodr. Fl. Nov. Holland. 546 (1810)	2:441
e	<b>Monotoca empetrifolia</b> R.Br., Prodr. Fl. Nov. Holland. 547 (1810)	2:443
	<b>Monotoca glauca</b> (Labill.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 636 (1917)	2:442
	Monotoca lineata R.Br. = <b>Monotoca glauca</b>	
	Monotoca lineata var. linifolia Rodway = <b>Monotoca linifolia</b>	
e	<b>Monotoca linifolia</b> (Rodway) W.M.Curtis subsp. <b>algida</b> Jarman, Aspects of Tasmanian Botany: 102 (1991)	2:442
e	<b>Monotoca linifolia</b> (Rodway) W.M.Curtis subsp. <b>linifolia</b> , The Student's Flora of Tasmania 2: 464 (1963)	2:442
	<b>Monotoca scoparia</b> (Sm.) R.Br., Prodr. Fl. Nov. Holland. 547 (1810)	2:442
	Monotoca scoparia var. submutica Benth. = <b>Monotoca submutica</b>	
e	<b>Monotoca submutica</b> (Benth.) Jarman var. <b>autumnalis</b> Jarman, Pap. & Proc. Roy. Soc. Tasmania 112: 1 (1978)	
e	<b>Monotoca submutica</b> (Benth.) Jarman var. <b>submutica</b> , Pap. & Proc. Roy. Soc. Tasmania 112: 1 (1978)	
t	<b>Montitega dealbata</b> (R.Br.) C.M.Weiller, Austral. Syst. Bot. 23: 328 (2010)	2:427
e	<b>Pentachondra ericifolia</b> Hook.f., London J. Bot. 6: 271 (1847)	2:430
e	<b>Pentachondra involucrata</b> R.Br., Prodr. Fl. Nov. Holland. 549 (1810)	2:429
	Pentachondra mucronata Hook.f. = <b>Leucopogon fraseri</b>	
	<b>Pentachondra pumila</b> (J.R.Forst. & G.Forst.) R.Br., Prodr. Fl. Nov. Holland. 549 (1810)	2:429
	Pentachondra verticillata Hook.f. = <b>Androstoma verticillata</b>	
	Pilitis acerosa Lindl. = <b>Richea acerosa</b>	
	Pilitis milliganii Hook.f. = <b>Richea milliganii</b>	
e	<b>Planocarpa nitida</b> (Jarman) C.M.Weiller, Austral. Syst. Bot. 9: 517 (1996)	
e	<b>Planocarpa petiolaris</b> (DC.) C.M.Weiller, Austral. Syst. Bot. 9: 514 (1996)	2:426
e	<b>Planocarpa sulcata</b> (Mihaiich) C.M.Weiller, Austral. Syst. Bot. 9: 516 (1996)	
e	<b>Prionotes cerinthoides</b> (Labill.) R.Br., Prodr. Fl. Nov. Holland. 553 (1810)	2:453
e	<b>Richea acerosa</b> (Lindl.) F.Muell., Fragm. (Mueller) 6: 69 (1867)	2:458
e	<b>Richea alpina</b> Menadue, Austral. Syst. Bot. 13: 798 (2000)	
	Richea angustifolia B.L.Burtt = <b>Richea scoparia</b>	2:460
e	<b>Richea × curtisiae</b> A.M.Gray, Muelleria 2: 143 (1971) = <b>R. pandanifolia</b> × <b>R. scoparia</b>	
e	<b>Richea dracophylla</b> R.Br., Prodr. Fl. Nov. Holland. 555 (1810)	2:460
e	<b>Richea gunnii</b> Hook.f., London J. Bot. 6: 273 (1847)	2:459
e	<b>Richea milliganii</b> (Hook.f.) F.Muell., Fragm. (Mueller) 6: 69 (1867)	2:458
e	<b>Richea pandanifolia</b> Hook.f. subsp. <b>pandanifolia</b> , Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 50 (1844)	2:460
e	<b>Richea pandanifolia</b> Hook.f. subsp. <b>ramulosa</b> Menadue, Austral. Syst. Bot. 13: 795 (2000)	2:460
e	<b>Richea procera</b> (F.Muell.) F.Muell., Fragm. (Mueller) 6: 68 (1867)	2:458
e	<b>Richea scoparia</b> Hook.f., London J. Bot. 6: 273 (1847)	2:459
e	<b>Richea sprengelioides</b> (R.Br.) F.Muell., Fragm. (Mueller) 6: 68 (1867)	2:457

	Sprengelia sp. Mt Field (A.M.Gray 1513) Tas Herbarium = <b>Sprengelia minima</b>	
e	<b>Sprengelia sp. Mt Read</b> , Telopea 15: 63 (2013) (R.K. Crowden 0802 006) Tas Herbarium	
e	<b>Sprengelia distichophylla</b> (Rodway) W.M.Curtis, The Student's Flora of Tasmania 2: 464 (1963)	2:456
	<b>Sprengelia incarnata</b> Sm., Kongl. Vetensk. Acad. Nya Handl. 15: 260 (1794)	2:454
	Sprengelia incarnata f. montana (R.Br.) Siebert & Voss = <b>Sprengelia montana</b>	
	Sprengelia incarnata f. propinqua (A.Cunn. ex DC.) Siebert & Voss = <b>Sprengelia propinqua</b>	
	Sprengelia incarnata var. distichophylla Rodway = <b>Sprengelia distichophylla</b>	
	Sprengelia incarnata var. montana (R.Br.) Domin = <b>Sprengelia montana</b>	
	Sprengelia macrantha Hook.f. = <b>Sprengelia propinqua</b>	
e	<b>Sprengelia minima</b> Crowden, Telopea 15: 62 (2013)	
e	<b>Sprengelia montana</b> R.Br., Prodr. Fl. Nov. Holland. 555 (1810)	2:455
e	<b>Sprengelia propinqua</b> A.Cunn. ex DC., Prodr. [A. P. de Candolle] 7: 768 (1839)	
n	Sprengelia propinqua var. demissa F.Muell. = <b>Sprengelia minima</b>	
	Stenantha pinifolia R.Br. = <b>Astroloma pinifolium</b>	
	Styphelia abietina Labill. = <b>Leptecophylla abietina</b>	
	<b>Styphelia adscendens</b> R.Br., Prodr. Fl. Nov. Holland. 537 (1810)	2:423
	Styphelia billardierei = <b>Cyathodes glauca</b>	
	Styphelia ciliata (R.Br.) F.Muell. = <b>Brachyloma ciliatum</b>	
	Styphelia collina Labill. = <b>Leucopogon collinus</b>	
	Styphelia empetrifolia (R.Br.) F.Muell. = <b>Monotoca empetrifolia</b>	
	Styphelia glauca Labill. = <b>Monotoca glauca</b>	
	Styphelia hookeri = <b>Acrothamnus hookeri</b>	
	Styphelia lanceolata Sm. = <b>Leucopogon affinis</b>	
	Styphelia montana (R.Br.) F.Muell. = <b>Acrothamnus montanus</b>	
	Styphelia mucronata (Hook.f.) J.H.Willis = <b>Leucopogon fraseri</b>	
	Styphelia oxycedrus Labill. = <b>Leptecophylla oxycedrus</b>	
	Styphelia serrulata Labill. = <b>Acrotriche serrulata</b>	
	Styphelia straminea (R.Br.) Spreng. = <b>Cyathodes straminea</b>	
	Styphelia strigosa Sm. = <b>Lissanthe strigosa</b>	
	Styphelia trichocarpa Labill. = <b>Leucopogon ericoides</b>	
	Styphelia virgata Labill. = <b>Leucopogon virgatus</b>	
e	<b>Trochocarpa cunninghamii</b> (DC.) W.M.Curtis, The Student's Flora of Tasmania 2: 463 (1963)	2:431
e	<b>Trochocarpa disticha</b> (Labill.) Spreng., Syst. Veg. (ed. 16) [Sprengel] 1: 660 (1824)	2:430
	Trochocarpa disticha var. cunninghamii (DC.) Benth. = <b>Trochocarpa cunninghamii</b>	
	Trochocarpa disticha var. microphylla F.Muell. = <b>Trochocarpa disticha</b>	
	Trochocarpa glauca (Labill.) Spreng. = <b>Cyathodes glauca</b>	
e	<b>Trochocarpa gunnii</b> (Hook.f.) Benth., Fl. Austral. 4: 167 (1868)	2:431
e	<b>Trochocarpa thymifolia</b> (R.Br.) Spreng., Syst. Veg. (ed. 16) [Sprengel] 1: 660 (1824)	2:431
	<b>ERICACEAE</b>	FTO 105
n i *	Arbutus unedo L., Sp. Pl. 1: 395 (1753)	
	Brossaea lanceolata (Hook.f.) Kuntze = <b>Gaultheria lanceolata</b>	
i t	<b>Calluna vulgaris</b> (L.) Hull, Brit. Fl., ed. 2, 1: 114 (1808)	2:418
	Erica andromedaeflora Andrews sensu Curtis (1963) = <b>Erica holosericea</b>	2:419
i	<b>Erica arborea</b> L., Sp. Pl. 1: 353 (1753)	2:420

i	<b>Erica baccans</b> L., Mant. Pl. 2: 233 (1771)	2:419
i	<b>Erica caffra</b> L., Sp. Pl. 1: 353 (1753)	2:421
i t	<b>Erica holosericea</b> Salisb., Trans. Linn. Soc. London 6: 352 (1802)	2:419
i	<b>Erica lusitanica</b> Rudolphi, J. Bot. (Schrader) 2: 286 (1800)	2:420
i t	<b>Erica scoparia</b> L., Sp. Pl. 1: 353 (1753)	
	Erica aff. willmorei Knowles & Westc. sensu Curtis (1963) recorded in error	2:420
	Gaultheria antipoda G.Forst. sensu Hooker (1860), Bentham (1868), Rodway (1903) = <b>Gaultheria depressa</b> (misapplied in Tasmania)	
	Gaultheria antipoda G.Forst. var. depressa (Hook.f.) Hook.f. = <b>Gaultheria depressa</b>	
t	<b>Gaultheria depressa</b> Hook.f., London J. Bot. 6: 267 (1847)	2:417
	<b>Gaultheria hispida</b> R.Br., Prodr. Fl. Nov. Holland. 559 (1810)	2:417
e	<b>Gaultheria lanceolata</b> Hook.f., London J. Bot. 6: 267 (1847)	2:416
e	<b>Gaultheria tasmanica</b> (Hook.f.) D.J.Middleton, Edinburgh J. Bot. 47: 299 (1990)	2:416
	Pernettya lanceolata (Hook.f.) B.L.Burt & A.W.Hill = <b>Gaultheria lanceolata</b>	2:416
	Pernettya tasmanica Hook.f. = <b>Gaultheria tasmanica</b>	2:416
i	<b>Rhododendron ponticum</b> L., Sp. Pl., ed. 2: 562 (1762)	
<b>ESCALLONIACEAE</b>		FTO 52, 131
e	<b>Anopterus glandulosus</b> Labill., Nov. Holl. Pl. 1: 86 t.112 (1805)	1:181
e	<b>Tetracarpaea tasmanica</b> Hook., Icon. Pl. 3: t.264 (1840)	1:182
<b>EUCRYPHIACEAE</b>		FTO 72
	Carpodontos lucida Labill. = <b>Eucryphia lucida</b>	
	Eucryphia billardierei Spach nom. illeg., nom. superfl. = <b>Eucryphia lucida</b>	
	Eucryphia billardierei Spach var. milliganii (Hook.f.) Benth. nom. illeg. = <b>Eucryphia milliganii</b>	
e	<b>Eucryphia lucida</b> (Labill.) Baill., Hist. Pl. (Baillon) 1: 402 (1869)	1:183
e	<b>Eucryphia milliganii</b> Hook.f. subsp. <b>milliganii</b> , Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 54 (1855)	1:183
e	<b>Eucryphia milliganii</b> Hook.f. subsp. <b>pubescens</b> R.W.Barnes, G.J.Jord., R.S.Hill & McCoull, Austral. J. Bot. 48: 488 (2000)	
<b>EUPHORBIACEAE</b>		
	Adriana quadripartita (Labill.) Müll.Arg. sensu Bentham (1873) recorded in error	
	Amperea spartioides Brongn. = <b>Amperea xiphoclada</b> var. <b>xiphoclada</b>	
	<b>Amperea xiphoclada</b> (Sieber ex Spreng.) Druce var. <b>xiphoclada</b> , Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 604 (1917)	3:637
	Bertya rosmarinifolia (A.Cunn.) Planch. sensu Curtis (1967) = <b>Bertya tasmanica</b> subsp. <b>tasmanica</b> (misapplied in Tasmania)	
e	<b>Bertya tasmanica</b> (Sond. & F.Muell.) Müll.Arg. subsp. <b>tasmanica</b> , Linnaea 34: 63 (1865)	3:637
	Beyeria backhousei Hook.f. = <b>Beyeria lechenaultii</b>	
	<b>Beyeria lechenaultii</b> (DC.) Baill., Adansonia 6: 307 (1866)	3:635
	Beyeria lechenaultii (DC.) Baill. var. latifolia Grüning = <b>Beyeria lechenaultii</b>	
	Beyeria oblongifolia (Klotzsch) Hook.f. = <b>Beyeria viscosa</b>	
	Beyeria opaca F.Muell. sensu Bentham (1873) = <b>Beyeria lechenaultii</b> (misapplied in Tasmania)	
	<b>Beyeria viscosa</b> (Labill.) Miq., Ann. Sci. Nat., Bot., sér. 3, 1: 350 t.15 (1844)	3:635
	Chamaesyce drummondii (Boiss.) D.C.Hassall = Euphorbia drummondii	3:632
i	<b>Euphorbia cyparissias</b> L., Sp. Pl. 1: 461 (1753)	3:633
	Euphorbia drummondii Boiss. sensu Curtis (1967) recorded in error	3:632

i	<b>Euphorbia exigua</b> L., Sp. Pl. 1: 456 (1753)	
i	<b>Euphorbia helioscopia</b> L., Sp. Pl. 1: 459 (1753)	3:632
i	<b>Euphorbia lathyris</b> L., Sp. Pl. 1: 457 (1753)	3:632
i	<b>Euphorbia paralias</b> L., Sp. Pl. 1: 458 (1753)	3:633
i	<b>Euphorbia peplus</b> L., Sp. Pl. 1: 456 (1753)	3:633
n i * t	<i>Euphorbia stricta</i> L., Syst. Nat., ed. 10, 2: 1049 (1759)	
	<b>Micrantheum hexandrum</b> Hook.f., London J. Bot. 6: 283 (1847)	3:634
e	<b>Micrantheum serpentinum</b> Orchard, Aspects of Tasmanian Botany: 60 (1991)	
	<i>Oreoporanthera petalifera</i> Orchard & J.B.Davies = <b>Poranthera petalifera</b>	
	<b>Phyllanthus australis</b> Hook.f., London J. Bot. 6: 284 (1847)	3:638
	<b>Phyllanthus gunnii</b> Hook.f., London J. Bot. 6: 284 (1847)	3:638
	<b>Poranthera microphylla</b> Brongn., Ann. Sci. Nat. (Paris) 29: 385 (1833)	3:634
e	<b>Poranthera petalifera</b> (Orchard & J.B.Davies) Halford & R.J.F.Hend., Austrobaileya 7: 22 (2005)	
	<i>Pseudanthus divaricatissimus</i> (Müll.Arg.) Benth. sensu Buchanan (1999) recorded in error	
	<b>Pseudanthus ovalifolius</b> F.Muell., Trans. & Proc. Philos. Inst. Victoria 2: 66 (1857)	3:634
	<i>Pseudanthus tasmanicus</i> Rodway = <b>Muehlenbeckia axillaris</b> (Polygonaceae)	3:593
	<i>Ricinocarpos major</i> F.Muell. sensu Bentham (1873) – a name of uncertain application	
	<b>Ricinocarpos pinifolius</b> Desf., Mém. Mus. Hist. Nat. 3: 459 t.22 (1817)	3:636
<b>FABACEAE (LEGUMINOSAE)</b>		
	<b>Almaleea subumbellata</b> (Hook.) Crisp & P.H.Weston, Telopea 4: 310 (1991)	1:141
	<i>Anthyllis vulneraria</i> L. recorded in error, only known from cultivated plants	
	<b>Aotus ericoides</b> (Vent.) G.Don, Gen. Hist. 2: 120 (1832)	1:137
	<i>Aotus ferruginea</i> Labill. = <b>Aotus ericoides</b>	
	<i>Aotus villosa</i> (Andrews) Sm. = <b>Aotus ericoides</b>	1:137
	<b>Bossiaea cinerea</b> R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 4: 268 (1812)	1:147
	<i>Bossiaea cinerea</i> R.Br. var. <i>rigida</i> Rodway = <b>Bossiaea tasmanica</b>	
	<b>Bossiaea cordifolia</b> Sweet, Fl. Australas. (Sweet) 20: 20 (1827)	
	<b>Bossiaea cordigera</b> Benth. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 95 t.16 (1856)	1:147
	<i>Bossiaea ensata</i> Sieber sensu Hooker (1860) non Sieber = <b>Bossiaea riparia</b>	
	<b>Bossiaea heterophylla</b> Vent., Descr. Pl. Nouv. 1: 7, pl.7 (1800)	
	<i>Bossiaea microphylla</i> Sm. sensu Curtis (1956) = <b>Bossiaea tasmanica</b>	
	<i>Bossiaea obcordata</i> (Vent.) Druce = <b>Bossiaea tasmanica</b>	1:148
	<b>Bossiaea prostrata</b> R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 4: 268 (1812)	1:147
	<b>Bossiaea riparia</b> A.Cunn. ex Benth., Fl. Austral. 2: 166 (1864)	1:148
e	<b>Bossiaea tasmanica</b> I.Thomps., Muellera 30: 144 (2012)	1:148
i	<b>Callistachys lanceolata</b> Vent., Jard. Malmaison 2: 115, t. 115 (1805)	
i	<b>Chamaecytisus palmensis</b> (Christ) F.A.Bisby & K.W.Nicholls, Bot. J. Linn. Soc. 74: 114 (1977)	1:151
	<i>Chamaecytisus proliferus</i> Link sensu Curtis & Morris (1975) = <b>Chamaecytisus palmensis</b>	1:151
	<i>Coronilla varia</i> L. = <b>Securigera varia</b>	1:165
	<b>Cullen microcephalum</b> (Rchb. ex Kunze) J.W.Grimes, Muellera 9: 195 (1996)	1:163
	<i>Cytisus monspessulanus</i> L. = <b>Genista monspessulana</b>	
i	<b>Cytisus multiflorus</b> (Aiton) Sweet, Hort. Brit. [Sweet] 112 (1826)	
	<i>Cytisus palmensis</i> (H.Christ) Hutch. = <b>Chamaecytisus palmensis</b>	1:151
	<i>Cytisus proliferus</i> L.f. sensu Curtis (1956) = <b>Chamaecytisus palmensis</b>	
i	<b>Cytisus scoparius</b> (L.) Link, Enum. Hort. Berol. Alt. 2: 241 (1822)	1:151

	<b>Daviesia latifolia</b> R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 20 (1811)	1:136
	<b>Daviesia sejugata</b> G.Chandler & Crisp, Austral. Syst. Bot. 10: 33 (1997)	
	<b>Daviesia ulicifolia</b> Andrews subsp. <b>ruscifolia</b> (A.Cunn. ex Benth.) G.Chandler & Crisp, Austral. Syst. Bot. 10: 44 (1997)	1:136
	<b>Daviesia ulicifolia</b> Andrews subsp. <b>ulicifolia</b> , Bot. Repos. 5: t.304 (1803)	1:136
	<i>Daviesia ulicina</i> Sm. nom. illeg., nom. superfl. = <b>Daviesia ulicifolia</b>	
	<i>Daviesia ulicina</i> Sm. f. <i>communis</i> Benth. = <b>Daviesia ulicifolia</b>	
	<i>Daviesia ulicina</i> Sm. f. <i>ruscifolia</i> (A.Cunn. ex Benth.) Benth. = <b>Daviesia ulicifolia</b> subsp. <b>ruscifolia</b>	
	<i>Daviesia ulicina</i> Sm. f. <i>subumbellata</i> Benth. = <b>Daviesia ulicifolia</b>	
	<i>Daviesia umbellata</i> Labill. = <b>Daviesia ulicifolia</b>	
	<b>Desmodium gunnii</b> Benth. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 101 (1856)	1:166
	<b>Desmodium varians</b> (Labill.) G.Don, Gen. Hist. 2: 298 (1832)	1:166
	<i>Desmodium varians</i> (Labill.) G.Don var. <i>gunnii</i> (Benth. ex Hook.f.) Benth. = <b>Desmodium gunnii</b>	
	<b>Dillwynia cinerascens</b> R.Br. ex Sims, Bot. Mag. 48: t.2247 (1821)	1:145
	<i>Dillwynia ericifolia</i> Sm. f. <i>glaberrima</i> (Sm.) Benth. sensu Bentham (1864) = <b>Dillwynia glaberrima</b> (misapplied in Tasmania)	
	<i>Dillwynia floribunda</i> Sm. sensu Bentham (1864), Rodway (1903), Curtis (1956) = <b>Dillwynia sericea</b> (misapplied in Tasmania)	
	<b>Dillwynia glaberrima</b> Sm., Ann. Bot. [König & Sims] 1: 510 (1805)	1:144
	<b>Dillwynia sericea</b> A.Cunn., Geogr. Mem. New South Wales: 347 (1825)	1:144
i	<b>Dipogon lignosus</b> (L.) Verdc., Taxon 17: 537 (1968)	
	<i>Dolichos lignosus</i> L. = <b>Dipogon lignosus</b>	
	<b>Eutaxia microphylla</b> (R.Br.) C.H.Wright & Dewar, Johnson's Gard. Dict. 1052 (1894)	1:143
	<i>Eutaxia microphylla</i> (R.Br.) C.H.Wright & Dewar var. <i>microphylla</i> = <b>Eutaxia microphylla</b>	
i	<b>Genista linifolia</b> L., Sp. Pl., ed. 2, 2: 997 (1763)	
	<i>Genista maderensis</i> (Webb & Berthel.) Lowe sensu Curtis & Morris (1975) = <b>Genista stenopetala</b>	1:152
i	<b>Genista monspessulana</b> (L.) L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 98 (1962)	1:151
i	<b>Genista × spachiana</b> Webb, Bot. Mag. 71: t.4195 (1845)	
i	<b>Genista stenopetala</b> Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 3(2): 39 (1836)	1:152
	<b>Glycine clandestina</b> J.C.Wendl., Bot. Beob. [Wendland]: 54 (1798)	1:168
	<i>Glycine clandestina</i> J.C.Wendl. var. <i>latrobeana</i> Rodway = <b>Glycine latrobeana</b>	
	<b>Glycine latrobeana</b> (Meisn.) Benth., Fl. Austral. 2: 244 (1864)	1:168
	<b>Glycine microphylla</b> (Benth.) Tindale, Brunonia 9: 181 (1987)	
	<b>Glycine tabacina</b> (Labill.) Benth., Fl. Austral. 2: 244 (1864)	
	<b>Gompholobium ecostatum</b> Kuchel, Suppl. Black's Fl. S. Austral. 182 (1965)	1:135
	<i>Gompholobium ellipticum</i> Labill. = <b>Oxylobium ellipticum</b>	
	<b>Gompholobium huegelii</b> Benth., Enum. Pl. [Endlicher]: 29 (1837)	1:135
	<i>Gompholobium latifolium</i> Labill. nom. illeg., non Sm. = <b>Gompholobium huegelii</b>	
	<b>Goodia lotifolia</b> Salisb., Parad. Lond. 1: t.41 (1806)	1:149
	<i>Goodia lotifolia</i> Salisb. var. <i>lotifolia</i> = <b>Goodia lotifolia</b>	1:149
	<i>Goodia lotifolia</i> Salisb. var. <i>pubescens</i> (Sims) H.B.Will. = <b>Goodia pubescens</b>	1:149
	<b>Goodia pubescens</b> Sims, Bot. Mag. 32: 1310 (1810)	1:149
	<i>Hardenbergia monophylla</i> (Vent.) Benth. = <b>Hardenbergia violacea</b>	

	<b>Hardenbergia violacea</b> (Schneev.) Stearn, J. Bot. 78: 70 (1940)	1:169
n	Hedysarum coronarium L. – previously listed as naturalised but insufficient evidence exists to support this	1:165
	<b>Hovea corrickiae</b> J.H.Ross, Muelleria 7: 203 (1990)	
	<b>Hovea heterophylla</b> A.Cunn. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 93 (1856)	1:148
	Hovea lanceolata Sims sensu Buchanan (1999) = <b>Hovea tasmanica</b>	
	Hovea linearis (Sm.) R.Br. sensu Buchanan (1999) = <b>Hovea heterophylla</b>	
	Hovea longifolia R.Br. sensu Curtis (1956), Curtis & Morris (1975) misapplied to <b>H. montana</b> & <b>H. tasmanica</b>	1:149
	<b>Hovea magnibractea</b> I.Thomps., Austral. Syst. Bot. 14: 74 (2001)	
	<b>Hovea montana</b> (Hook.f.) J.H.Ross, Muelleria 6: 427 (1988)	1:149
	Hovea purpurea Sweet var. montana Hook.f. = <b>Hovea montana</b>	
e	<b>Hovea tasmanica</b> I.Thomps. & J.H.Ross, Austral. Syst. Bot. 14: 68 (2001)	1:149
	Indigofera australis Willd. f. angulata Benth. = <b>Indigofera australis</b> subsp. <b>australis</b>	
	<b>Indigofera australis</b> Willd. subsp. <b>australis</b> , Sp. Pl., ed. 4 [Willdenow], 3: 1235 (1802)	1:164
i	<b>Kennedia nigricans</b> Lindl., Edwards's Bot. Reg. 20: 1715, pl. 1715 (1835)	
	<b>Kennedia prostrata</b> R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 4: 299 (1812)	1:169
i	<b>Kennedia rubicunda</b> (Schneev.) Vent., Jard. Malmaison 2: 104 t.104 (1804)	
i #	<b>Laburnum anagyroides</b> Medik., Vorles. Churpfälz. Phys.-Öcon. Ges. 2: 363 (1787)	
i #	<b>Lathyrus nissolia</b> L., Sp. Pl. 2: 729 (1753)	1:168
i	<b>Lathyrus tingitanus</b> L., Sp. Pl. 2: 732 (1753)	
	Leptocyamus clandestinus (J.C.Wendl.) Benth. var. clandestinus = <b>Glycine clandestina</b>	
	Leptocyamus clandestinus (J.C.Wendl.) Benth. var. microphylla (Benth.) Hook.f. = <b>Glycine microphylla</b>	
	Leptocyamus tasmanicus Benth. ex Hook.f. = <b>Glycine latrobeana</b>	
i #	<b>Lotus angustissimus</b> L., Sp. Pl. 2: 774 (1753)	
	<b>Lotus australis</b> Andrews, Bot. Repos. 10 t.624 (1811)	1:162
i	<b>Lotus corniculatus</b> L. var. <b>corniculatus</b> , Sp. Pl. 2: 776 (1753)	1:162
i	<b>Lotus corniculatus</b> L. var. <b>tenuifolius</b> L., Sp. Pl. 2: 776 (1753)	1:162
	Lotus hispidus Desf. ex DC. = <b>Lotus subbiflorus</b>	1:163
	Lotus pedunculatus Cav. sensu Curtis & Morris (1975) = <b>Lotus uliginosus</b>	1:163
	Lotus suaveolens Pers. nom. illeg. = <b>Lotus subbiflorus</b>	1:163
i	<b>Lotus subbiflorus</b> Lag., Varied. Ci. 2(4): 213 (1805)	1:163
	Lotus tenuis Waldst. & Kit. = <b>Lotus corniculatus</b> var. <b>tenuifolius</b>	
i	<b>Lotus uliginosus</b> Schkuhr, Handb. [C.Schkuhr] 2: 412 (1796)	1:163
i #	<b>Lupinus angustifolius</b> L., Sp. Pl. 2: 721 (1753)	1:150
i	<b>Lupinus arboreus</b> Sims, Bot. Mag. 18: t.682 (1803)	1:150
i	<b>Lupinus × regalis</b> Bergmans, Vaste Pl. Rotsheesters: 328 (1924)	
i	<b>Medicago arabica</b> (L.) Huds., Fl. Angl. (Hudson): 288 (1762)	1:154
i #	<b>Medicago arborea</b> L., Sp. Pl. 2: 778 (1753)	
	Medicago denticulata Willd. = <b>Medicago polymorpha</b>	
n	Medicago falcata L. sensu Curtis & Morris (1975) = <b>Medicago sativa</b> nothosubsp × <b>varia</b> (misapplied in Tasmania)	1:153
	Medicago hispida Gaertn. var. apiculata (Willd.) Urb. = <b>Medicago polymorpha</b>	
	Medicago hispida Gaertn. var. confinis (W.D.J.Koch) Burnat = <b>Medicago polymorpha</b>	

	Medicago hispida Gaertn. var. denticulata (Willd.) Urb. = <b>Medicago polymorpha</b>	
i	<b>Medicago lupulina</b> L., Sp. Pl. 2: 779 (1753)	1:153
	Medicago maculata Willd. = <b>Medicago arabica</b>	
i	<b>Medicago minima</b> (L.) L. ex Bartal., Cat. Piante Siena 61 (1776)	1:154
i	<b>Medicago polymorpha</b> L., Sp. Pl. 2: 779 (1753)	1:154
i	<b>Medicago sativa</b> L., Sp. Pl. 2: 778 (1753)	1:153
n i #	<b>Medicago sativa</b> L. nothosubsp. × <b>varia</b> (Martyn) Arcang., Comp. Fl. Ital. 160 = <b>Medicago falcata</b> × <b>M. sativa</b>	1:153
i	<b>Medicago scutellata</b> (L.) Mill., Gard. Dict., ed. 8, no. 2 (1768)	1:154
	Melilotus alba Desr. nom. illeg. = <b>Melilotus albus</b>	
i	<b>Melilotus albus</b> Medik., Vorles. Churpfälz. Phys.-Öcon. Ges. 2: 382 (1787)	1:155
	Melilotus arvensis Wallr. = <b>Melilotus officinalis</b>	
i	<b>Melilotus indicus</b> (L.) All., Fl. Pedem. 1: 308 (1785)	1:155
i	<b>Melilotus officinalis</b> (L.) Pall., Reise Russ. Reich. 3: 537 (1776)	1:155
	Melilotus parviflorus Desf. = <b>Melilotus indicus</b>	
	<b>Mirbelia oxylobioides</b> F.Muell., Fragm. (Mueller) 2: 154 (1861)	
n	Onobrychis viciifolia Scop. – previously listed as naturalised but insufficient evidence exists to support this	1:165
	Ononis arvensis L. sensu Rodway (1903) = <b>Ononis spinosa</b>	1:152
	Ononis repens L. sensu Curtis (1956), Curtis & Morris (1975) = <b>Ononis spinosa</b>	1:152
i	<b>Ononis spinosa</b> L., Sp. Pl. 2: 716 (1753)	1:152
i	<b>Ornithopus compressus</b> L., Sp. Pl. 2: 744 (1753)	
i	<b>Ornithopus pinnatus</b> (Mill.) Druce, J. Bot. 45: 420 (1907)	
i #	<b>Ornithopus sativus</b> Brot., Fl. Lusit. 2: 160 (1805)	
	Ornithopus sativus Brot. subsp. sativus = <b>Ornithopus sativus</b>	
	<b>Oxylobium arborescens</b> R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 10 (1811)	1:134
	<b>Oxylobium ellipticum</b> (Vent.) R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 10 (1811)	1:134
	Oxylobium ellipticum (Vent.) R.Br. var. angustifolium Benth. = <b>Oxylobium arborescens</b>	
e	<b>Phyllota diffusa</b> (Hook.f.) F.Muell., Fragm. (Mueller) 1: 8 (1858)	1:137
	Platylobium formosum Sm. sensu Bentham (1864) = <b>Platylobium parviflorum</b> (misapplied in Tasmania)	1:146
	Platylobium formosum Sm. subsp. parviflorum (Sm.) A.T.Lee = <b>Platylobium parviflorum</b>	
	Platylobium murrayanum Hook.f. = <b>Platylobium triangulare</b>	
	<b>Platylobium obtusangulum</b> Hook., Bot. Mag. 60: t.3258 (1833)	1:146
	<b>Platylobium parviflorum</b> Sm., Spec. Bot. New Holland 18 (1795)	
	<b>Platylobium triangulare</b> R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 4: 266 (1812)	1:146
i	<b>Podalyria sericea</b> (Andrews) R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 7 (1811)	
	Psoralea adscendens F.Muell. = <b>Cullen microcephalum</b>	1:163
i	<b>Psoralea arborea</b> Sims, Bot. Mag. 46: t. 2090 (1819)	
	Psoralea gunnii Hook.f. = <b>Cullen microcephalum</b>	
i	<b>Psoralea pinnata</b> L., Sp. Pl. 2: 762 (1753)	1:163
	Pultenaea cordata Hook. = <b>Pultenaea juniperina</b>	
	<b>Pultenaea daphnoides</b> J.C.Wendl., Bot. Beob. [Wendland]: 49 (1798)	1:139
	Pultenaea daphnoides J.C.Wendl. var. obcordata (Andrews) Hook.f. = <b>Pultenaea daphnoides</b>	
	<b>Pultenaea dentata</b> Labill., Nov. Holl. Pl. 1: 103 t.131 (1805)	1:141

	Pultenaea diffusa Hook.f. = <b>Phyllota diffusa</b>	
	<b>Pultenaea fasciculata</b> Benth., Commentat. Legum. Gen. 18 (1837)	1:142
	<b>Pultenaea gunnii</b> Benth. subsp. <b>gunnii</b> , Commentat. Legum. Gen. 18 (1837)	1:140
	Pultenaea gunnii Benth. var. <b>baeckeoides</b> (A.Cunn. ex Benth.) Rodway = <b>Pultenaea gunnii</b> subsp. <b>gunnii</b>	1:140
	Pultenaea hibbertioides Hook.f. = <b>Pultenaea mollis</b>	1:141
	<b>Pultenaea humilis</b> Benth. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 91 (1856)	1:142
	<b>Pultenaea juniperina</b> Labill., Nov. Holl. Pl. 1: 102 t.130 (1805)	1:142
	Pultenaea juniperina Labill. var. <b>latifolia</b> Benth. = <b>Pultenaea juniperina</b>	
	<b>Pultenaea mollis</b> Lindl., Three Exped. Australia [Mitchell] 2: 258 (1838)	1:141
	Pultenaea paleacea Willd. var. <b>sericea</b> Benth. = <b>Pultenaea sericea</b>	1:140
	<b>Pultenaea pedunculata</b> Hook., Bot. Mag. 55: t.2859 (1828)	1:140
	Pultenaea pimelioides Hook.f. = <b>Pultenaea dentata</b>	
	<b>Pultenaea prostrata</b> Benth. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 89 (1856)	1:142
	Pultenaea selaginoides Hook.f. = <b>Stonesiella selaginoides</b>	1:141
	<b>Pultenaea sericea</b> (Benth.) Corrick, Muelleria 8: 392 (1995)	1:140
	<b>Pultenaea stricta</b> Sims, Bot. Mag. 38: t.1588 (1813)	1:140
	Pultenaea subumbellata Hook. = <b>Almaleea subumbellata</b>	1:141
	<b>Pultenaea tenuifolia</b> R.Br. ex Sims, Bot. Mag. 46: t.2086 (1819)	1:143
	Sarothamnus scoparius (L.) Wimm. ex W.D.J.Koch = <b>Cytisus scoparius</b>	1:151
n i	<b>Securigera varia</b> (L.) Lassen, Svensk. Bot. Tidskr. 83: 86 (1989)	1:165
i	<b>Spartium junceum</b> L., Sp. Pl. 2: 708 (1753)	
	<b>Sphaerolobium minus</b> Labill., Nov. Holl. Pl. 1: 108, t.138 (1805)	
	<b>Sphaerolobium vimineum</b> Sm., Ann. Bot. [König & Sims] 1: 509 (1805)	1:135
e	<b>Stonesiella selaginoides</b> (Hook.f.) Crisp & P.H.Weston, Taxon 48: 712 (1999)	1:141
	<b>Swainsona lessertiifolia</b> DC., Ann. Sci. Nat., Bot., sér. 1, 4: 99 (1825)	1:164
	Trifolium agrarium L. sensu Rodway (1903) = <b>Trifolium campestre</b>	
i	<b>Trifolium alexandrinum</b> L., Cent. Pl. 1: 25 (1755)	
i	<b>Trifolium angustifolium</b> L., Sp. Pl. 2: 769 (1753)	1:161
i	<b>Trifolium arvense</b> L., Sp. Pl. 2: 769 (1753)	1:160
i	<b>Trifolium campestre</b> Schreb., Die Kleearten Deutschlands 1(4): t.17.4 (1800)	1:157
i	<b>Trifolium cernuum</b> Brot., Phytogr. Lusitan. Select. (1816-1827) 1: 150 (1816)	1:159
i	<b>Trifolium dubium</b> Sibth., Fl. Oxon. 231 (1794)	1:157
i	<b>Trifolium fragiferum</b> L., Sp. Pl. 2: 772 (1753)	1:158
i	<b>Trifolium glomeratum</b> L., Sp. Pl. 2: 770 (1753)	1:159
i	<b>Trifolium hybridum</b> L., Sp. Pl. 2: 766 (1753)	1:159
i	<b>Trifolium incarnatum</b> L., Sp. Pl. 2: 769 (1753)	1:160
i	<b>Trifolium lappaceum</b> L., Sp. Pl. 2: 768 (1753)	
	Trifolium medium Huds. = <b>Trifolium pratense</b> (misapplied in Tasmania)	1:159
i	<b>Trifolium micranthum</b> Viv., Fl. Libyc. Spec. 45, pl. 19, Fig. 1 (1824)	1:158
i	<b>Trifolium ornithopodioides</b> L., Sp. Pl. 2: 766 (1753)	1:152
i	<b>Trifolium pratense</b> L., Sp. Pl. 2: 768 (1753)	1:160
	Trifolium procumbens L. sensu Rodway (1903) = <b>Trifolium dubium</b> (misapplied in Tasmania)	
i	<b>Trifolium repens</b> L., Sp. Pl. 2: 767 (1753)	1:158
i	<b>Trifolium resupinatum</b> L., Sp. Pl. 2: 771 (1753)	1:158



i	<b>Trifolium scabrum</b> L., Sp. Pl. 2: 770 (1753)	1:160
i	<b>Trifolium squamosum</b> L., Amoen. Acad., Linnaeus ed. 4: 105 (1759)	1:161
i	<b>Trifolium stellatum</b> L., Sp. Pl. 2: 769 (1753)	1:161
i	<b>Trifolium striatum</b> L., Sp. Pl. 2: 770 (1753)	1:160
i	<b>Trifolium subterraneum</b> L., Sp. Pl. 2: 767 (1753)	1:159
i	<b>Trifolium suffocatum</b> L., Mant. Pl. 2: 276 (1771)	1:159
i	<b>Trifolium tomentosum</b> L., Sp. Pl. 2: 771 (1753)	1:158
n	<i>Trifolium uniflorum</i> L. – previously listed as naturalised but insufficient evidence exists to support this	
	<i>Trigonella ornithopodioides</i> (L.) DC. = <b>Trifolium ornithopodioides</b>	1:152
i	<b>Ulex europaeus</b> L., Sp. Pl. 2: 741 (1753)	1:150
	<i>Vicia angustifolia</i> L. = <b>Vicia sativa</b>	1:167
	<i>Vicia cracca</i> L. sensu Curtis & Morris (1975) = <b>Vicia villosa</b>	1:167
i	<b>Vicia hirsuta</b> (L.) Gray, Nat. Arr. Brit. Pl. 2: 614 (1821)	1:166
	<i>Vicia lathyroides</i> L. recorded in error	
i	<b>Vicia sativa</b> L. subsp. <b>nigra</b> (L.) Ehrh., Hannover. Mag. 15: 229 (1780)	1:167
i	<b>Vicia sativa</b> L. subsp. <b>sativa</b> , Sp. Pl. 2: 736 (1753)	1:167
i	<b>Vicia tetrasperma</b> (L.) Schreb., Spic. Fl. Lips. 26 (1771)	1:167
i	<b>Vicia villosa</b> Roth subsp. <b>eriocarpa</b> (Hauskn.) P.W.Ball, Feddes Repert. 79: 45 (1968)	1:167
	<i>Viminaria denudata</i> (Vent.) Sm. = <b>Viminaria juncea</b>	
	<b>Viminaria juncea</b> (Schrader & J.C.Wendl.) Hoffmanns., Verz. Pfl.-Kult. 200 (1824)	1:136
	<b>FAGACEAE</b>	FTO 66
	<i>Fagus cunninghamii</i> Hook. = <b>Nothofagus cunninghamii</b>	
	<i>Fagus gunnii</i> Hook.f. = <b>Nothofagus gunnii</b>	
	<i>Fuscospora gunnii</i> (Hook.f.) Heenan & Smissen = <b>Nothofagus gunnii</b>	
	<i>Lophozonia cunninghamii</i> (Hook.) Heenan & Smissen = <b>Nothofagus cunninghamii</b>	
	<b>Nothofagus cunninghamii</b> (Hook.) Oerst., Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd. (ser. 5), 9: 355 (1871)	3:647
e	<b>Nothofagus gunnii</b> (Hook.f.) Oerst., Kongel. Danske Vidensk. Selsk. Skr., Naturvidensk. Math. Afd. (ser. 5), 9: 354 (1871)	3:646
	<b>FRANKENIACEAE</b>	FTO 93
	<b>Frankenia pauciflora</b> DC. var. <b>gunnii</b> Summerh., J. Linn. Soc. Bot. 48: 366 (1930)	1:64
	<b>FUMARIACEAE</b>	FTO 45
i	<b>Fumaria bastardii</b> Boreau, Rev. Bot. Recueil Mans. 2: 359 (1847)	1:30
i	<b>Fumaria densiflora</b> DC., Cat. Pl. Horti Monsp. 113 (1813)	1:30
i	<b>Fumaria muralis</b> Sond. ex W.D.J.Koch subsp. <b>muralis</b> , Syn. Fl. Germ. Helv., ed. 2: 1017 (1845)	1:30
n i *	<i>Fumaria officinalis</i> L. subsp. <i>officinalis</i> , Sp. Pl. 2: 700 (1753)	
n	<i>Pseudofumaria alba</i> (Mill.) Lidén subsp. <i>alba</i> – previously listed as naturalised but insufficient evidence exists to support this	
	<b>GENTIANACEAE</b>	
	<i>Centaurium australe</i> (R.Br.) Druce = <b>Schenkia australis</b>	3:479
i	<b>Centaurium erythraea</b> Rafn, Danm. Holst. Fl. 2: 75 (1800)	3:480
	<i>Centaurium pulchellum</i> (Sw.) Druce sensu Curtis (1967) = <b>Centaurium tenuiflorum</b>	3:480
	<i>Centaurium spicatum</i> (L.) Fritsch ex Janch. sensu Buchanan (1999) = <b>Schenkia australis</b>	3:479

i	<b>Centaurium tenuiflorum</b> (Hoffmanns. & Link) Fritsch ex Janch., Mitt. Naturwiss. Vereins Univ. Wien 5: 97 (1907)	3:480
	Chionogentias brevisepala L.G.Adams = <b>Gentianella brevisepala</b>	3:482
	Chionogentias cunninghamii L.G.Adams = <b>Gentianella cunninghamii</b>	3:482
	Chionogentias demissa L.G.Adams = <b>Gentianella demissa</b>	3:482
	Chionogentias diemensis (Griseb.) L.G.Adams = <b>Gentianella diemensis</b>	3:482
	Chionogentias eichleri L.G.Adams = <b>Gentianella eichleri</b>	3:482
e ?x	<b>Chionogentias grandis</b> L.G.Adams, Austral. Syst. Bot. 8: 960 (1995)	3:482
	Chionogentias gunniana L.G.Adams = <b>Gentianella gunniana</b>	3:482
	Chionogentias pleurogynoides (Griseb.) L.G.Adams = <b>Gentianella pleurogynoides</b>	3:482
	Chionogentias polysperes L.G.Adams = <b>Gentianella polysperes</b>	3:482
i	<b>Cicendia filiformis</b> (L.) Delarbre, Fl. Auvergne (Delarbre), ed. 2, 1: 29 (1800)	3:481
	Erythraea australis R.Br. = <b>Schenkia australis</b>	
	Exacum ovatum Labill. = <b>Sebaea ovata</b>	
	Gentiana montana R.Br. = <b>Gentianella gunniana</b> and <b>Gentianella pleurogynoides</b>	
	Gentiana saxosa G.Forst. sensu Rodway (1903) = <b>Gentianella</b> spp.	
e	<b>Gentianella brevisepala</b> (L.G.Adams) Glenny, New Zealand J. Bot. 42: 518 (2004)	3:482
x	<b>Gentianella cunninghamii</b> (L.G.Adams) Glenny subsp. <b>cunninghamii</b> , New Zealand J. Bot. 42: 518 (2004)	3:482
e	<b>Gentianella demissa</b> (L.G.Adams) Glenny, Muelleria 26: 95 (2008)	3:482
e	<b>Gentianella diemensis</b> (Griseb.) J.H.Willis subsp. <b>diemensis</b> , Vict. Naturalist 73: 199 (1957)	3:482
e	<b>Gentianella diemensis</b> (Griseb.) J.H.Willis subsp. <b>plantaginea</b> (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004)	3:482
e	<b>Gentianella eichleri</b> (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004)	3:482
	Gentianella grandis (L.G.Adams) Glenny = <b>Chionogentias grandis</b>	3:482
	<b>Gentianella gunniana</b> (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004)	3:482
e	<b>Gentianella pleurogynoides</b> (Griseb.) Glenny subsp. <b>milliganii</b> (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004)	3:482
e	<b>Gentianella pleurogynoides</b> (Griseb.) Glenny subsp. <b>pleurogynoides</b> , New Zealand J. Bot. 42: 519 (2004)	3:482
	<b>Gentianella polysperes</b> (L.G.Adams) Glenny, New Zealand J. Bot. 42: 519 (2004)	3:482
	<b>Schenkia australis</b> (R.Br.) G.Mans., Taxon 53: 725 (2004)	3:479
	<b>Sebaea albidiflora</b> F.Muell., Trans. Philos. Soc. Victoria 1: 46 (1855)	3:479
	<b>Sebaea ovata</b> (Labill.) R.Br., Prodr. Fl. Nov. Holland. 452 (1810)	3:478
	<b>GERANIACEAE</b>	FTO 54
i	<b>Erodium botrys</b> (Cav.) Bertol., Amoen. Ital. 35 (1819)	1:98
i	<b>Erodium cicutarium</b> (L.) L'Hér. ex Aiton, Hortus Kew. (W.Aiton) 2: 414 (1789)	1:97
n i *	<b>Erodium malacoides</b> (L.) L'Hér. ex Aiton, Hortus Kew. (W.Aiton) 2: 415 (1789)	1:98
i	<b>Erodium moschatum</b> (L.) L'Hér. ex Aiton, Hortus Kew. (W.Aiton) 2: 414 (1789)	1:97
	<b>Geranium</b> sp. <b>Pale Pink Flowers (M.Gray 5847) Vic. Herbarium</b>	
	<b>Geranium brevicaule</b> Hook., J. Bot. (Hooker) 1: 252 (1834)	
i	<b>Geranium dissectum</b> L., Cent. Pl. 1: 21 (1755)	1:96
	<b>Geranium dissectum</b> L. var. <b>australe</b> Benth. sensu Bentham (1863) = <b>G. potentilloides</b> , <b>G. retrorsum</b> and <b>G. solanderi</b>	
	<b>Geranium dissectum</b> L. var. <b>pilosum</b> Benth. = <b>Geranium solanderi</b>	

	Geranium dissectum L. var. potentilloides Benth. = <b>Geranium potentilloides</b>	
?i	<b>Geranium homeanum</b> Turcz., Bull. Soc. Imp. Naturalistes Moscou 36: 591 (1863)	
	Geranium microphyllum Hook.f. = <b>Geranium potentilloides</b> var. <b>potentilloides</b>	
i	<b>Geranium molle</b> L., Sp. Pl. 2: 682 (1753)	1:96
	Geranium molle L. var. molle = <b>Geranium molle</b>	
	Geranium pilosum G.Forst. nom. inval., nom. nud. = <b>Geranium solanderi</b>	
	<b>Geranium potentilloides</b> L'Hér. ex DC. var. <b>potentilloides</b> , Prodr. [A. P. de Candolle] 1: 639 (1824)	1:95
	<b>Geranium retrorsum</b> L'Hér. ex DC., Prodr. [A. P. de Candolle] 1: 644 (1824)	
i t	<b>Geranium rotundifolium</b> L., Sp. Pl. 2: 683 (1753)	1:96
	Geranium sessiliflorum Cav. = <b>Geranium brevicaule</b>	
	Geranium sessiliflorum Cav. subsp. brevicaule (Hook.) Carolin = <b>Geranium brevicaule</b>	1:95
	<b>Geranium solanderi</b> Carolin, Proc. Linn. Soc. New South Wales 89: 350 (1965)	1:95
n i	<b>Geranium yeoi</b> Aedo & Muñoz Garm., Kew Bull. 52: 727 (1997)	
	Pelargonium acugnaticum Thouars sensu Hooker (1860) = <b>Pelargonium inodorum</b> (misapplied in Tasmania)	
i	<b>Pelargonium × asperum</b> Ehrh. ex Willd., Sp. Pl., ed. 4 [Willdenow] 3: 678 (1800)	1:99
	<b>Pelargonium australe</b> Willd., Sp. Pl., ed. 4 [Willdenow] 3: 675 (1800)	1:98
i	<b>Pelargonium capitatum</b> (L.) L'Hér. ex Aiton, Hortus Kew. (W.Aiton)2: 425 (1789)	
i	<b>Pelargonium × domesticum</b> L.H.Bailey, Stand. Cycl. Hort. 2532 (1916)	1:99
	<b>Pelargonium inodorum</b> Willd., Enum. Pl. [Willdenow] 2: 702 (1809)	1:99
	<b>Pelargonium littorale</b> Endl., Bot. Arch.(Hügel): t.5 (1837)	
	<b>GOODENIACEAE</b>	FTO 129
x	<b>Coopernookia barbata</b> (R.Br.) Carolin, Proc. Linn. Soc. New South Wales 92: 213 (1968)	2:401
	<b>Dampiera stricta</b> (Sm.) R.Br., Prodr. Fl. Nov. Holland. 589 (1810)	2:404
	Goodenia amplexans F.Muell. sensu Curtis (1963) recorded in error	2:400
	Goodenia barbata R.Br. = <b>Coopernookia barbata</b>	2:401
	<b>Goodenia elongata</b> Labill., Nov. Holl. Pl. 1: 52, t.75 (1805)	2:401
	<b>Goodenia geniculata</b> R.Br., Prodr. Fl. Nov. Holland. 577 (1810)	2:401
	Goodenia geniculata R.Br. var. lanata (R.Br.) Rodway = <b>Goodenia lanata</b>	
	Goodenia hederacea R.Br. = <b>Goodenia lanata</b> (misapplied in Tasmania)	
	<b>Goodenia humilis</b> R.Br., Prodr. Fl. Nov. Holland. 575 (1810)	2:402
	<b>Goodenia lanata</b> R.Br., Prodr. Fl. Nov. Holland. 577 (1810)	2:401
	<b>Goodenia ovata</b> Sm., Trans. Linn. Soc. London 2: 347 (1794)	2:400
x	<b>Goodenia pinnatifida</b> Schltldl., Linnaea 21: 450 (1848)	
	Goodenia repens Labill. = <b>Selliera radicans</b>	
	<b>Scaevola aemula</b> R.Br., Prodr. Fl. Nov. Holland. 584 (1810)	2:404
	<b>Scaevola albida</b> (Sm.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916, Suppl. 2: 644 (1917)	2:404
	Scaevola calendulacea (Andrews) Druce sensu Curtis (1963) recorded in error	
	Scaevola cuneiformis Labill. recorded in error	
	<b>Scaevola hookeri</b> (de Vriese) F.Muell. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 231 (1856)	2:403
	Scaevola microcarpa Cav. = <b>Scaevola albida</b>	
	<b>Selliera radicans</b> Cav., Anales. Hist. Nat. 1: 41, t.5 (1799)	2:402
	<b>Velleia montana</b> Hook.f., London J. Bot. 6: 265 (1847)	2:399

	<b>Velleia paradoxa</b> R.Br., Prodr. Fl. Nov. Holland. 580 (1810)	2:398
	<b>GROSSULARIACEAE</b>	FTO 50
i t	<b>Ribes sanguineum</b> Pursh, Fl. Amer. Sept. (Pursh) 1: 164 (1813)	
	<b>GUNNERACEAE</b>	FTO 49
e	<b>Gunnera cordifolia</b> (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 125 (1856)	1:194
	<b>GYROSTEMONACEAE</b>	FTO 83
	Didymotheca thesioides Hook.f. = <b>Gyrostemon thesioides</b>	3:581
	<b>Gyrostemon thesioides</b> (Hook.f.) A.S.George, Fl. Australia 8: 392 (1982)	3:581
	<b>HALORAGACEAE</b>	
	<b>Gonocarpus humilis</b> Orchard, Bull. Auckland Inst. Mus. 10: 195 (1975)	
	<b>Gonocarpus micranthus</b> Thunb. subsp. <b>micranthus</b> , Nov. Gen. Pl. [Thunberg] 3: 55 (1783)	1:190
	<b>Gonocarpus montanus</b> (Hook.f.) Orchard, Bull. Auckland Inst. Mus. 10: 172 (1975)	1:192
	<b>Gonocarpus serpyllifolius</b> Hook.f., Icon. Pl. 3: t.290 (1840)	1:192
	<b>Gonocarpus tetragynus</b> Labill., Nov. Holl. Pl. 1: 39, t.53 (1805)	1:191
	<b>Gonocarpus teucrioides</b> DC., Prodr. [A. P. de Candolle] 3: 66 (1828)	1:191
	<b>Haloragis aspera</b> Lindl., J. Exped. Trop. Australia [Mitchell]: 306 (1848)	1:191
	<b>Haloragis brownii</b> (Hook.f.) Schindl., Pflanzenr. (Engler) 23: 54 (1905)	1:192
	Haloragis ceratophylla Endl. sensu Bentham (1864) = <b>Haloragis aspera</b> (misapplied in Tasmania)	
	Haloragis depressa (A.Cunn.) Walp. sensu Bentham (1864) = <b>Gonocarpus montanus</b> and <b>G. serpyllifolius</b>	
	Haloragis depressa (A.Cunn.) Walp. var. montana (Hook.f.) Hook.f. = <b>Gonocarpus montanus</b>	
	Haloragis gunnii Hook.f. = <b>Gonocarpus teucrioides</b>	
	<b>Haloragis heterophylla</b> Brongn., Voy. Monde, Atlas: t.68A (1826)	1:190
	Haloragis micrantha (Thunb.) R.Br. ex Sieber & Zucc. = <b>Gonocarpus micranthus</b>	1:190
	Haloragis montana Hook.f. = <b>Gonocarpus montanus</b>	1:192
	<b>Haloragis myriocarpa</b> Orchard, Bull. Auckland Inst. Mus. 10: 132 (1975)	
	Haloragis pinnatifida A.Gray sensu Hooker (1860) = <b>Haloragis aspera</b>	
	Haloragis serpyllifolia (Hook.f.) Walp. = <b>Gonocarpus serpyllifolius</b>	1:192
	Haloragis tetragyna (Labill.) Hook.f. = <b>Gonocarpus tetragynus</b>	1:191
	Haloragis teucrioides (DC.) Schldl. = <b>Gonocarpus teucrioides</b> & <b>G. humilis</b>	1:191
	Meionectes brownii Hook.f. = <b>Haloragis brownii</b>	
	<b>Myriophyllum amphibium</b> Labill., Nov. Holl. Pl. 2: 70, t.220 (1806)	1:193
i	<b>Myriophyllum aquaticum</b> (Vell.) Verdc., Kew Bull. 28: 36 (1973)	1:193
e	<b>Myriophyllum austropygmaeum</b> Orchard, Brunonia 8: 280 (1986)	
	Myriophyllum brasiliense Cambess. = <b>Myriophyllum aquaticum</b>	1:193
?i	<b>Myriophyllum crispatum</b> Orchard, Brunonia 8: 210 (1986)	
	Myriophyllum elatinoides Gaudich. sensu Curtis (1956), Curtis & Morris (1975) =	
	<b>Myriophyllum salsugineum</b>	1:193
x	<b>Myriophyllum glomeratum</b> Schindl., Pflanzenr. (Engler) 23: 103 (1905)	
	<b>Myriophyllum integrifolium</b> (Hook.f.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 123 (1856)	1:194
	<b>Myriophyllum muelleri</b> Sond., Linnaea 28: 233 (1856)	
	<b>Myriophyllum pedunculatum</b> Hook.f. subsp. <b>longibracteolatum</b> (Schindl.) Orchard, Brunonia 8: 273 (1986)	1:194

	<b>Myriophyllum pedunculatum</b> Hook.f. subsp. <b>pedunculatum</b> , London J. Bot. 6: 474 (1847)	1:194
	Myriophyllum propinquum A.Cunn. sensu Curtis (1956), Curtis & Morris (1975) =	
	<b>Myriophyllum simulans</b> & <b>M. variifolium</b>	1:193
	<b>Myriophyllum salsugineum</b> Orchard, Brunonia 4: 48 (1981)	1:193
	<b>Myriophyllum simulans</b> Orchard, Brunonia 8: 203 (1986)	1:193
	<b>Myriophyllum variifolium</b> Hook.f., Icon. Pl. 3: t.289 (1840)	1:193
<b>HYDROPHYLLACEAE</b>		
i	<b>Phacelia tanacetifolia</b> Benth., Edwards's Bot. Reg. 20: t.1696 (1835)	3:486
<b>LAMIACEAE (LABIATAE)</b>		
	<b>Ajuga australis</b> R.Br., Prodr. Fl. Nov. Holland. 503 (1810)	3:557
i	<b>Ajuga reptans</b> L., Sp. Pl. 2: 561 (1753)	
i	<b>Cedronella canariensis</b> (L.) Webb & Berthel., Hist. Nat. Illes Canaries (Phytogr.) 3: 87 (1845)	3:548
i	<b>Lamium amplexicaule</b> L., Sp. Pl. 2: 579 (1753)	3:552
i	<b>Lamium purpureum</b> L., Sp. Pl. 2: 579 (1753)	3:552
i	<b>Lavandula stoechas</b> L. subsp. <b>stoechas</b> , Sp. Pl. 2: 573 (1753)	
i	<b>Leonotis leonurus</b> (L.) R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 3: 410 (1811)	
	<b>Lycopus australis</b> R.Br., Prodr. Fl. Nov. Holland. 500 (1810)	3:546
i	<b>Marrubium vulgare</b> L., Sp. Pl. 2: 583 (1753)	3:550
i	<b>Melissa officinalis</b> L., Sp. Pl. 2: 592 (1753)	3:546
	<b>Mentha australis</b> R.Br., Prodr. Fl. Nov. Holland. 505 (1810)	3:544
	<b>Mentha diemenica</b> Spreng. var. <b>diemenica</b> , Syst. Veg. (ed. 16) [Sprengel] 2: 724 (1825)	3:544
	<b>Mentha diemenica</b> Spreng. var. <b>serpyllifolia</b> (Benth.) J.H.Willis, Muelleria 1: 144 (1967)	3:544
	<i>Mentha gracilis</i> R.Br. = <b>Mentha diemenica</b>	
i	<b>Mentha × piperita</b> L., Sp. Pl. 2: 576 (1753)	3:545
i	<b>Mentha pulegium</b> L., Sp. Pl. 2: 577 (1753)	3:545
	<i>Mentha serpyllifolia</i> Benth. = <b>Mentha diemenica</b> var. <b>serpyllifolia</b>	
i #	<b>Mentha spicata</b> L., Sp. Pl. 2: 576 (1753)	3:545
i	<b>Nepeta cataria</b> L., Sp. Pl. 2: 570 (1753)	3:548
x	<b>Prostanthera cuneata</b> Benth., Prodr. [A. P. de Candolle] 12: 560 (1848)	3:554
	<b>Prostanthera lasianthos</b> Labill. var. <b>lasianthos</b> , Nov. Holl. Pl. 2: 18 t.157 (1806)	3:553
	<i>Prostanthera retusa</i> R.Br. = <b>Prostanthera rotundifolia</b>	
	<b>Prostanthera rotundifolia</b> R.Br., Prodr. Fl. Nov. Holland. 509 (1810)	3:553
i	<b>Prunella laciniata</b> (L.) L., Sp. Pl., ed. 2, 2: 837 (1763)	3:550
i	<b>Prunella vulgaris</b> L., Sp. Pl. 2: 600 (1753)	3:550
	<i>Salvia horminoides</i> Pourr. = <b>Salvia verbenaca</b>	3:547
i	<b>Salvia verbenaca</b> L. var. <b>verbenaca</b> , Sp. Pl. 1: 25 (1753)	3:547
	<b>Scutellaria humilis</b> R.Br., Prodr. Fl. Nov. Holland. 507 (1810)	3:549
i	<b>Stachys arvensis</b> (L.) L., Sp. Pl., ed. 2, 2: 814 (1763)	3:551
i t	<b>Stachys palustris</b> L., Sp. Pl. 2: 580 (1753)	3:551
	<b>Teucrium corymbosum</b> R.Br., Prodr. Fl. Nov. Holland. 504 (1810)	3:556
i t	<b>Teucrium scorodonia</b> L., Sp. Pl. 2: 564 (1753)	
e	<b>Westringia angustifolia</b> R.Br., Prodr. Fl. Nov. Holland. 501 (1810)	3:555
e	<b>Westringia brevifolia</b> Benth., Labiat. Gen. Spec. 459 (1834)	3:555
	<i>Westringia brevifolia</i> Benth. var. <i>raleighii</i> (B.Boivin) W.M.Curtis = <b>Westringia brevifolia</b>	
	<i>Westringia dampieri</i> R.Br. sensu Bentham (1870) misapplied to <b>W. angustifolia</b> or <b>W. rigida</b>	

	<b>Westringia rigida</b> R.Br., Prodr. Fl. Nov. Holland. 501 (1810)	3:555
e	<b>Westringia rubiifolia</b> R.Br., Prodr. Fl. Nov. Holland. 501 (1810)	3:556
	<b>LAURACEAE</b>	FTO 5
	<b>Cassytha glabella</b> R.Br. f. <b>dispar</b> (Schtdl.) J.Z.Weber, J. Adelaide Bot. Gard. 3: 209 (1981)	3:597
	<b>Cassytha glabella</b> R.Br. f. <b>glabella</b> , Prodr. Fl. Nov. Holland. 404 (1810)	3:597
	<b>Cassytha melantha</b> R.Br., Prodr. Fl. Nov. Holland. 404 (1810)	3:597
e	<b>Cassytha pedicellosa</b> J.Z.Weber, J. Adelaide Bot. Gard. 3: 214 (1981)	
	<b>Cassytha pubescens</b> R.Br., Prodr. Fl. Nov. Holland. 404 (1810)	3:597
	<b>LENTIBULARIACEAE</b>	
	Polypompholyx tenella (R.Br.) Lehm. = <b>Utricularia tenella</b>	3:537
	<b>Utricularia australis</b> R.Br., Prodr. Fl. Nov. Holland. 430 (1810)	3:536
	<b>Utricularia barkeri</b> R.W.Jobson, Telopea 15: 131 (2013)	
	<b>Utricularia dichotoma</b> Labill., Nov. Holl. Pl. 1: 11, t.8 (1805)	3:536
	Utricularia dichotoma Labill. var. uniflora (R.Br.) Benth. = <b>Utricularia uniflora</b>	
	Utricularia flexuosa Vahl = <b>Utricularia australis</b>	3:536
n i *	Utricularia gibba L., Sp. Pl. 1: 18 (1753)	
	<b>Utricularia lateriflora</b> R.Br., Prodr. Fl. Nov. Holland. 431 (1810)	3:536
	<b>Utricularia monanthos</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 299 (1857)	3:537
	<b>Utricularia tenella</b> R.Br., Prodr. Fl. Nov. Holland. 432 (1810)	3:537
	<b>Utricularia uniflora</b> R.Br., Prodr. Fl. Nov. Holland. 431 (1810)	3:536
	<b>Utricularia violacea</b> R.Br., Prodr. Fl. Nov. Holland. 431 (1810)	
	<b>LINACEAE</b>	FTO 77
	Linum albidum Ewart & Jean White = <b>Linum marginale</b>	
i t	<b>Linum bienne</b> Mill., Gard. Dict., ed. 8, no.8 (1768)	
i t	<b>Linum catharticum</b> L., Sp. Pl. 1: 281 (1753)	1:92
	Linum gallicum L. = <b>Linum trigynum</b>	
	<b>Linum marginale</b> A.Cunn., Geogr. Mem. New South Wales [Field]: 357 (1825)	1:92
i	<b>Linum trigynum</b> L., Sp. Pl. 1: 279 (1753)	1:93
i	<b>Linum usitatissimum</b> L., Sp. Pl. 1: 277 (1753)	1:92
	<b>LOGANIACEAE</b>	
	Mitrasacme archeri Hook.f. = <b>Schizacme archeri</b>	3:475
	Mitrasacme distylis F.Muell. = <b>Phyllangium distylis</b>	3:477
	Mitrasacme divergens Hook.f. = <b>Phyllangium divergens</b>	3:476
	Mitrasacme montana Hook.f. ex Benth. = <b>Schizacme montana</b>	3:475
	Mitrasacme paradoxa R.Br. sensu Curtis (1967) = <b>Phyllangium divergens</b>	3:476
	<b>Mitrasacme pilosa</b> Labill. var. <b>pilosa</b> , Nov. Holl. Pl. 1: 36, t.49 (1805)	3:476
	<b>Mitrasacme pilosa</b> Labill. var. <b>stuartii</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 274 (1857)	3:476
	<b>Mitrasacme serpyllifolia</b> R.Br., Prodr. Fl. Nov. Holland. 454 (1810)	3:475
	<b>Phyllangium distylis</b> (F.Muell.) Dunlop, Fl. Australia 28: 315 (1996)	3:477
	<b>Phyllangium divergens</b> (Hook.f.) Dunlop, Fl. Australia 28: 315 (1996)	3:476
e	<b>Schizacme archeri</b> (Hook.f.) Dunlop, Fl. Australia 28: 314 (1996)	3:475
	<b>Schizacme montana</b> (Hook.f. ex Benth.) Dunlop, Fl. Australia 28: 314 (1996)	3:475

	<b>LYTHRACEAE</b>	FTO 55
	<b>Lythrum hyssopifolia</b> L., Sp. Pl. 1: 447 (1753)	2:227
	<b>Lythrum salicaria</b> L., Sp. Pl. 1: 446 (1753)	2:226
	<b>MALVACEAE</b>	
i t	<b>Anisodonteia capensis</b> (L.) D.M.Bates, Gentes Herb. 10: 327 (1969)	
e	<b>Asterotrichion discolor</b> (Hook.) Melville, Kew Bull. 20: 512 (1967)	1:86
	<b>Gynatrix pulchella</b> (Willd.) Alef., Oesterr. Bot. Z. 12: 35 (1862)	1:87
n	Hibiscus trionum L. – previously listed as naturalised but insufficient evidence exists to support this	1:88
	Lavatera arborea L. = <b>Malva arborea</b>	1:84
	Lavatera cretica L. has also been missapplied to <b>M. preissiana</b> in = <b>Malva pseudolavatera</b>	1:84
	Lavatera plebeia Sims var. tomentosa Hook.f. = <b>Malva preissiana</b>	1:83
	<b>Lawrenzia spicata</b> Hook., Icon. Pl. 3: t.261 (1840)	1:88
?i #	<b>Lawrenzia squamata</b> Nees, Pl. Preiss. [J.G.C.Lehman] 1: 242 (1845)	
i	<b>Malva arborea</b> (L.) Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.) 1: 30 (1836)	1:84
	Malva australiana M.F.Ray = <b>Malva preissiana</b>	1:83
	Malva dendromorpha M.F.Ray nom. illeg., nom. superfl. = <b>Malva arborea</b>	
	Malva linnaei M.F.Ray nom. illeg., nom. superfl. = <b>Malva pseudolavatera</b>	
i	<b>Malva moschata</b> L., Sp. Pl. 2: 690 (1753)	1:85
i	<b>Malva neglecta</b> Wallr., Syll. Pl. Nov. 1: 140 (1824)	1:85
i	<b>Malva nicaeensis</b> All., Fl. Pedem. 2: 40 (1785)	1:85
i	<b>Malva parviflora</b> L., Demonstr. Pl. 18 (1753)	1:85
	<b>Malva preissiana</b> Miq., Pl. Preiss. [J.G.C.Lehman] 1: 238 (1845)	1:83
n i	<b>Malva pseudolavatera</b> Webb & Berthel., Hist. Nat. Iles Canaries (Phytogr.). 3(2(1)): 29 (1836)	1:84
	Malva rotundifolia L. sensu Rodway (1903) = <b>Malva neglecta</b> (misapplied in Tasmania)	
i	<b>Malva sylvestris</b> L., Sp. Pl. 2: 689 (1753)	1:85
i	<b>Modiola caroliniana</b> (L.) G.Don, Gen. Hist. 1: 466 (1831)	1:86
	Plagianthus pulchellus (Willd.) A.Gray ex Hook.f. = <b>Gynatrix pulchella</b>	1:87
	Plagianthus pulchellus (Willd.) Hook.f. var. tomentosus Rodway = <b>Gynatrix pulchella</b>	
	Plagianthus sidoides Hook. = <b>Asterotrichion discolor</b>	1:86
	Plagianthus spicatus (Hook.) Benth. = <b>Lawrenzia spicata</b>	1:88
	<b>MENYANTHACEAE</b>	FTO 128
	Limnanthemum exaltatum (Sol. ex Sims) F.Muell. = <b>Liparophyllum exaltatum</b>	
	Limnanthemum exiguum F.Muell. = <b>Liparophyllum exiguum</b>	
	Limnanthemum gunnii (Hook.f.) Hook.f. = <b>Liparophyllum gunnii</b>	
	<b>Liparophyllum exaltatum</b> (Sol. ex Sims) Tippet & Les, Novon 19: 408 (2009)	3:483
e	<b>Liparophyllum exiguum</b> (F.Muell.) Tippet & Les, Novon 19: 408 (2009)	3:483
t	<b>Liparophyllum gunnii</b> Hook.f., London J. Bot. 6: 473 bis (1847)	3:484
	Nymphoides crenata (F.Muell.) Kuntze sensu Curtis (1967) recorded in error	3:483
	Nymphoides exigua (F.Muell.) Kuntze = <b>Liparophyllum exiguum</b>	3:483
	<b>Ornduffia reniformis</b> (R.Br.) Tippet & Les, Novon 19: 410 (2009)	3:483
	<b>Ornduffia umbricola</b> (Aston) Tippet & Les var. <b>umbricola</b> , Novon 19: 410 (2009)	
	Villarsia exaltata (Sol. ex Sims) G.Don = <b>Liparophyllum exaltatum</b>	3:483
	Villarsia parnassifolia (Labill.) R.Br. = <b>Liparophyllum exaltatum</b>	
	Villarsia reniformis R.Br. = <b>Ornduffia reniformis</b>	3:483

Villarsia umbricola Aston = **Ornduffia umbricola**

**MIMOSACEAE**

- Acacia armata R.Br. = **Acacia paradoxa** 1:127
- e **Acacia axillaris** Benth., London J. Bot. 1: 341 (1842) 1:127
- n i **Acacia baileyana** F.Muell., Trans. & Proc. Roy. Soc. Victoria 24: 168 (1888)
- Acacia botrycephala (Vent.) Desf. = **Acacia terminalis** 1:129
- Acacia crassiuscula H.L.Wendl. sensu Hooker (1860), Bentham (1864) = **Acacia uncifolia**
- Acacia dealbata** Link subsp. **dealbata**, Enum. Hort. Berol. Alt. 2: 445 (1822) 1:130
- i **Acacia decurrens** Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1072 (1806) 1:130
- Acacia decurrens Willd. f. mollis (Lindl.) Benth. = **Acacia dealbata**
- Acacia dependens A.Cunn. ex Benth. nom. illeg. = **Acacia mucronata** subsp. **dependens**
- e **Acacia derwentiana** A.M.Gray, Muellera 21: 107 (2005)
- Acacia diffusa Ker Gawl. = **Acacia genistifolia**
- Acacia discolor (Andrews) Willd. = **Acacia terminalis**
- Acacia dissitiflora Benth. = **Acacia mucronata** subsp. **longifolia**
- i **Acacia floribunda** (Vent.) Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1051 (1806)
- Acacia genistifolia** Link, Enum. Hort. Berol. Alt. 2: 442 (1822) 1:126
- Acacia gunnii** Benth., London J. Bot. 1: 332 (1842) 1:127
- i **Acacia howittii** F.Muell., Vict. Naturalist 10: 16 (1893)
- n Acacia implexa Benth. sensu Buchanan (1999) = **Acacia uncifolia** (misapplied in Tasmania)
- Acacia juniperina (Vent.) Willd. sensu Bentham (1864) = **Acacia ulicifolia** (misapplied in Tasmania)
- Acacia leprosa** Sieber ex DC. var. **graveolens** Maslin & D.J.Murphy, Muellera 27: 201 (2009) 1:127
- Acacia linearis (J.C.Wendl.) Sims sensu Hooker (1860), Bentham (1864) = **Acacia mucronata** subsp. **mucronata** (misapplied in Tasmania)
- Acacia longifolia (Andrews) Willd. f. dissitiflora (Benth.) Benth. = **Acacia mucronata** subsp. **longifolia**
- Acacia longifolia (Andrews) Willd. f. mucronata (Willd. ex H.L.Wendl.) Benth. = **Acacia mucronata**
- Acacia longifolia** (Andrews) Willd. subsp. **longifolia**, Sp. Pl., ed. 4 [Willdenow], 4: 1052 (1806) 1:129
- Acacia longifolia** (Andrews) Willd. subsp. **sophorae** (Labill.) Court, Fl. Australia 11B: 491 (2001) 1:129
- Acacia mearnsii** De Wild., Pl. Bequaert. 3: 61 (1925) 1:130
- Acacia melanoxylon** R.Br., Hortus Kew. (W.T.Aiton), ed. 2, 5: 462 (1813) 1:128
- Acacia mollissima Willd. sensu Hooker (1860) = **Acacia mearnsii**
- e **Acacia mucronata** Willd. ex H.L.Wendl. subsp. **dependens** (Hook.f.) Court, Fl. Australia 11B: 491 (2001) 1:129
- Acacia mucronata** Willd. ex H.L.Wendl. subsp. **longifolia** (Benth.) Court, Fl. Australia 11B: 491 (2001) 1:129
- e **Acacia mucronata** Willd. ex H.L.Wendl. subsp. **mucronata**, Comm. Acac. Aphyll. 46 t.12 (1820) 1:129
- Acacia mucronata Willd. ex H.L.Wendl. var. dissitiflora Benth. = **Acacia mucronata** subsp. **longifolia**
- Acacia mucronata Willd. ex H.L.Wendl. var. linearis (Sims) Rodway = **Acacia mucronata** subsp. **mucronata**



	<b>Acacia myrtifolia</b> (Sm.) Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1054 (1806)	1:128
	Acacia ovoidea Benth. = <b>Acacia verticillata</b> subsp. <b>ovoidea</b>	
	Acacia oxycedrus Sieber ex DC. sensu Bentham (1864) recorded in error	
i	<b>Acacia paradoxa</b> DC., Cat. Pl. Horti Monsp. 74 (1813)	1:127
e	<b>Acacia pataczekii</b> D.I.Morris, Records of the Queen Victoria Museum 50: 1 (1974)	1:128
	Acacia penninervis Sieber ex DC. sensu Bentham (1864) recorded in error	
i	<b>Acacia pravissima</b> F.Muell. ex Benth., Linnaea 26: 608 (1854)	
i	<b>Acacia provincialis</b> A.Camus, Bull. Soc. Dendrol. France 64: 68 (1927)	
i	<b>Acacia pycnantha</b> Benth., London J. Bot. 1: 351 (1842)	
i	<b>Acacia retinodes</b> Schltld., Linnaea 20: 664 (1847)	
	Acacia retinodes Schltld. var. retinodes = <b>Acacia retinodes</b>	
	Acacia retinodes Schltld. var. uncifolia J.M.Black = <b>Acacia uncifolia</b>	1:128
e	<b>Acacia riceana</b> Hensl., Botanist 3: t.135 (1839)	1:127
	<b>Acacia siculiformis</b> A.Cunn. ex Benth., London J. Bot. 1: 337 (1842)	1:126
	Acacia sophorae (Labill.) R.Br. = <b>Acacia longifolia</b> subsp. <b>sophorae</b>	1:129
	<b>Acacia stricta</b> (Andrews) Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1052 (1806)	1:127
	Acacia stuartiana F.Muell. ex Benth. sensu Hooker (1860) = <b>Acacia siculiformis</b>	
	<b>Acacia suaveolens</b> (Sm.) Willd., Sp. Pl., ed. 4 [Willdenow], 4: 1050 (1806)	1:128
e	<b>Acacia sp. Sugarloaf (A.Moscal 8301) Tas Herbarium</b>	
	<b>Acacia terminalis</b> (Salisb.) J.F.Macbr., Contr. Gray Herb. 59: 7 (1919)	1:129
	<b>Acacia ulicifolia</b> (Salisb.) Court, Vict. Naturalist 73: 173 (1957)	1:126
	<b>Acacia uncifolia</b> (J.M.Black) O'Leary, J. Adelaide Bot. Gard. 21: 100 (2007)	1:128
	Acacia verniciflua A.Cunn. sensu Curtis (1963) = <b>Acacia leprosa</b> var. <b>graveolens</b>	1:127
	<b>Acacia verticillata</b> (L'Hér.) Willd. subsp. <b>ovoidea</b> (Benth.) Court, Fl. Australia 11B: 492 (2001)	1:126
	<b>Acacia verticillata</b> (L'Hér.) Willd. subsp. <b>ruscifolia</b> (A.Cunn. ex G.Don) Court, Fl. Australia 11B: 492 (2001)	1:126
	<b>Acacia verticillata</b> (L'Hér.) Willd. subsp. <b>verticillata</b> , Sp. Pl., ed. 4 [Willdenow], 4: 1049 (1806)	1:126
	Acacia verticillata (L'Hér.) Willd. var. latifolia Benth. = <b>Acacia verticillata</b> subsp. <b>ruscifolia</b>	
	Acacia vomeriformis A.Cunn. ex Benth. = <b>Acacia gunnii</b>	1:127
	Albizia lophantha (Willd.) Benth. = <b>Paraserianthes lophantha</b>	1:130
	Mimosa sophorae Labill. = <b>Acacia longifolia</b> subsp. <b>sophorae</b>	
	Mimosa suaveolens Sm. = <b>Acacia suaveolens</b>	
i	<b>Paraserianthes lophantha</b> (Willd.) I.C.Nielsen subsp. <b>lophantha</b> , Bull. Mus. Natl. Hist. Nat., B, Adansonia 5: 326 (1983) - Possibly native in some Furneaux populations, see Harris et al. (2001)	1:130
	Racosperma axillare (Benth.) Pedley = <b>Acacia axillaris</b>	1:127
	Racosperma dealbatum (Link) Pedley = <b>Acacia dealbata</b>	1:130
	Racosperma decurrens (Willd.) Pedley = <b>Acacia decurrens</b>	1:130
	Racosperma gunnii (Benth.) Pedley = <b>Acacia gunnii</b>	1:127
	Racosperma implexum (Benth.) Pedley = <b>Acacia uncifolia</b>	
	Racosperma mearnsii (De Wild.) Pedley = <b>Acacia mearnsii</b>	1:130
	Racosperma melanoxyton (R.Br.) Pedley = <b>Acacia melanoxyton</b>	1:128
	Racosperma myrtifolium (Sm.) Mart. = <b>Acacia myrtifolia</b>	1:128
	Racosperma paradoxum (DC.) Mart. = <b>Acacia paradoxa</b>	1:127
	Racosperma riceanum (Hensl.) Pedley = <b>Acacia riceana</b>	1:127

	Racosperma sophorae (Labill.) Mart. = <b>Acacia longifolia</b> subsp. <b>sophorae</b>	1:129
	Racosperma strictum (Andrews) Mart. = <b>Acacia stricta</b>	1:127
	Racosperma suaveolens (Sm.) Mart. = <b>Acacia suaveolens</b>	1:128
	Racosperma ulicifolium (Salisb.) Pedley = <b>Acacia ulicifolia</b>	1:126
	Racosperma vernicifluum (A.Cunn.) Pedley = <b>Acacia leprosa</b> var. <b>graveolens</b>	1:127
	Racosperma verticillatum (L'Hér.) Pedley = <b>Acacia verticillata</b>	1:126
	<b>MONIMIACEAE</b>	FTO 3, 4
	<b>Atherosperma moschatum</b> Labill. subsp. <b>moschatum</b> , Nov. Holl. Pl. 2: 74, t.224 (1806)	3:594
	<b>Hedycarya angustifolia</b> A.Cunn., Ann. Nat. Hist. Ser. I 1: 215 (1838)	3:595
	<b>MYOPORACEAE</b>	
	Myoporum adscendens R.Br. sensu Rodway (1903) = <b>Myoporum insulare</b>	
	<b>Myoporum insulare</b> R.Br., Prodr. Fl. Nov. Holland. 516 (1810)	3:540
	<b>Myoporum parvifolium</b> R.Br., Prodr. Fl. Nov. Holland. 516 (1810)	3:540
	Myoporum serratum R.Br. nom. illeg., sensu Bentham (1870) = <b>Myoporum insulare</b> (misapplied in Tasmania)	
	Myoporum serratum R.Br. f. obovatum = <b>Myoporum insulare</b>	
	Myoporum tasmanicum A.DC. = <b>Myoporum insulare</b>	
	<b>MYRTACEAE</b>	FTO 57
	Baeckea diffusa Sieber ex DC. = <b>Euryomyrtus ramosissima</b>	
	<b>Baeckea gunniana</b> Schauer, Repert. Bot. Syst. (Walpers) 2: 920 (1843)	1:198
e	<b>Baeckea leptocaulis</b> Hook.f., Icon. Pl. 3: t.298 (1840)	1:199
	Baeckea ramosissima A.Cunn. = <b>Euryomyrtus ramosissima</b>	1:198
	Baeckea thymifolia Hook.f. = <b>Euryomyrtus ramosissima</b>	
	Callistemon pallidus (Bonpl.) DC. = <b>Melaleuca pallida</b>	1:203
	Callistemon paludosus F.Muell. sensu Curtis (1956), Curtis & Morris (1975) = <b>Melaleuca pallida</b>	1:203
	Callistemon salignus (Sm.) Colvill ex Sweet sensu Bentham (1867) = <b>Melaleuca pallida</b> (misapplied in Tasmania)	
	Callistemon salignus (Sm.) Colvill ex Sweet var. australis Benth. = <b>Melaleuca pallida</b>	
	Callistemon salignus (Sm.) Colvill ex Sweet var. hebestachys Benth. = <b>Melaleuca pallida</b>	
	Callistemon salignus (Sm.) Colvill ex Sweet var. viridiflora Benth. = <b>Melaleuca virens</b>	
	Callistemon viridiflorus (Sims) Sweet = <b>Melaleuca virens</b>	1:203
	Calycothrix glabra (R.Br.) Hook.f. = <b>Calytrix tetragona</b>	
	<b>Calytrix tetragona</b> Labill., Nov. Holl. Pl. 2: 8, t.146 (1806)	1:197
	Eucalyptus acervula Hook. sensu Rodway (1903) = <b>Eucalyptus ovata</b>	
	Eucalyptus aggregata H.Deane & Maiden sensu Curtis (1956) = <b>Eucalyptus rodwayi</b>	
	Eucalyptus ambigua DC. can not be confidently applied	
e	<b>Eucalyptus amygdalina</b> Labill., Nov. Holl. Pl. 2: 14, t.154 (1806)	1:224
	Eucalyptus amygdalina Labill. var. nitida (Hook.f.) Benth. = <b>Eucalyptus nitida</b>	
	Eucalyptus amygdalina Labill. var. radiata (DC.) Benth. = <b>Eucalyptus radiata</b> subsp. <b>radiata</b>	
e	<b>Eucalyptus archeri</b> Maiden & Blakely, Crit. Rev. Eucalyptus 8: 58 (1929)	1:215
e	<b>Eucalyptus barberi</b> L.A.S.Johnson & Blaxell, Contr. N.S.W. Natl. Herb. 4: 288 (1972)	1:211
	Eucalyptus bicostata Maiden, Blakely & Simmonds sensu Curtis & Morris (1975) = <b>Eucalyptus globulus</b> subsp. <b>pseudoglobulus</b>	1:219

	<b>Eucalyptus brookeriana</b> A.M.Gray, Austral. Forest Res. 9: 111 (1979)	
e	<b>Eucalyptus coccifera</b> Hook.f., London J. Bot. 6: 477 (1847)	1:225
	<i>Eucalyptus coccifera</i> Hook.f. var. <i>parviflora</i> Benth. = <b>Eucalyptus coccifera</b>	
e	<b>Eucalyptus cordata</b> Labill. subsp. <b>cordata</b> , Nov. Holl. Pl. 2: 13, t.152 (1806)	1:218
e	<b>Eucalyptus cordata</b> Labill. subsp. <b>quadrangulosa</b> D.Nicolle, B.M.Potts & McKinnon, Pap. & Proc. Roy. Soc. Tasmania 142: 75 (2008)	
	<i>Eucalyptus coriacea</i> Schauer = <b>Eucalyptus pauciflora</b> subsp. <b>pauciflora</b>	
	<b>Eucalyptus dalrympleana</b> Maiden subsp. <b>dalrympleana</b> , Forest Fl. N.S.W. 7: 137, t.241 (1920)	1:214
e	<b>Eucalyptus delegatensis</b> R.T.Baker subsp. <b>tasmaniensis</b> Boland, Austral. Forest Res. 15: 177 (1985)	1:220
	<i>Eucalyptus gigantea</i> Hook.f. = <b>Eucalyptus delegatensis</b> subsp. <b>tasmaniensis</b>	1:220
	<b>Eucalyptus globulus</b> Labill. subsp. <b>globulus</b> , Voy. Rech. Perouse 1: 153, t.13 (1800)	1:218
	<b>Eucalyptus globulus</b> Labill. subsp. <b>pseudoglobulus</b> (Naudin) J.B.Kirkp., Bot. J. Linn. Soc. 69: 101 (1975)	1:219
e	<b>Eucalyptus gunnii</b> Hook.f. subsp. <b>divaricata</b> (McAulay & Brett) B.M.Potts, Pap. & Proc. Roy. Soc. Tasmania 135: 57 (2001)	
e	<b>Eucalyptus gunnii</b> Hook.f. subsp. <b>gunnii</b> , London J. Bot. 3: 499 (1844)	1:215
	<i>Eucalyptus haemastoma</i> Sm. sensu Rodway (1903) misapplied to a hybrid involving <b>Eucalyptus sieberi</b>	
e	<b>Eucalyptus johnstonii</b> Maiden, Crit. Rev. Eucalyptus 6: 280 (1922)	1:219
	<i>Eucalyptus linearis</i> Dehnh. = <b>Eucalyptus pulchella</b>	1:224
e	<b>Eucalyptus morrisbyi</b> Brett, Pap. & Proc. Roy. Soc. Tasmania 129: pl. 14-15 (1939)	1:215
	<i>Eucalyptus muelleri</i> T.B.Moore nom. illeg. = <b>Eucalyptus johnstonii</b>	
e	<b>Eucalyptus nebulosa</b> A.M.Gray, Kanunnah 3: 42 (2008)	
i	<b>Eucalyptus nitens</b> (H.Deane & Maiden) Maiden, Crit. Rev. Eucalyptus 2: 272 (1913)	
e	<b>Eucalyptus nitida</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 137, t.29 (1856)	1:225
	<b>Eucalyptus obliqua</b> L'Hér., Sert. Angl. 18, t.20 (1792)	1:220
	<b>Eucalyptus ovata</b> Labill. var. <b>ovata</b> , Nov. Holl. Pl. 2: 13, t.153 (1806)	1:211
	<b>Eucalyptus pauciflora</b> Sieber ex Spreng. subsp. <b>pauciflora</b> , Syst. Veg. (ed. 16) [Sprengel] 4: 195 (1827)	1:222
	<b>Eucalyptus perriniana</b> F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania: 181 (1894)	1:217
e	<b>Eucalyptus pulchella</b> Desf., Cat. Pl. Horti Paris., ed. 3: 284, 408 (1829)	1:224
	<b>Eucalyptus radiata</b> Sieber ex DC. subsp. <b>radiata</b> , Prodr. [A. P. de Candolle] 3: 218 (1828)	1:224
	<i>Eucalyptus radiata</i> Sieber ex DC. subsp. <i>robertsonii</i> sensu Curtis & Morris (1975) = <b>Eucalyptus radiata</b> subsp. <b>radiata</b>	
	<b>Eucalyptus regnans</b> F.Muell., Ann. Rep. Victorian Acclim. Soc. 20 in obs. (1871)	1:220
e	<b>Eucalyptus risdonii</b> Hook.f., London J. Bot. 6: 477 bis (1847)	1:226
	<i>Eucalyptus risdonii</i> Hook.f. var. <i>elata</i> = <b>Eucalyptus delegatensis</b> subsp. <b>tasmaniensis</b>	
e	<b>Eucalyptus rodwayi</b> R.T.Baker & H.G.Sm., Pap. & Proc. Roy. Soc. Tasmania 1912: 191 (1913)	1:211
n	<b>Eucalyptus rubida</b> H.Deane & Maiden subsp. <b>rubida</b> , Proc. Linn. Soc. New South Wales 24: 456, t.xl (1899)	1:214
	<i>Eucalyptus salicifolia</i> Cav. = <b>Eucalyptus amygdalina</b>	1:224
	<b>Eucalyptus sieberi</b> L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 125 (1962)	1:222
	<i>Eucalyptus sieberiana</i> F.Muell. nom. illeg. = <b>Eucalyptus sieberi</b>	

	Eucalyptus simmondsii Maiden = <b>Eucalyptus nitida</b>	1:225
	Eucalyptus stuartiana F.Muell. ex Miq. = <b>Eucalyptus ovata</b> var. <b>ovata</b>	
e	<b>Eucalyptus subcrenulata</b> Maiden & Blakely, Crit. Revis. Eucalyptus 8: 59 (1929)	1:219
	Eucalyptus tasmanica Blakely = <b>Eucalyptus tenuiramis</b>	1:226
e	<b>Eucalyptus tenuiramis</b> Miq., Ned. Kruidk. Arch. 4: 128 (1856)	1:226
e	<b>Eucalyptus urnigera</b> Hook.f., London J. Bot. 6: 477 bis (1847)	1:217
	Eucalyptus urnigera Hook.f. var. elongata Rodway = <b>Eucalyptus urnigera</b>	
e	<b>Eucalyptus vernicosa</b> Hook.f., London J. Bot. 6: 478 bis (1847)	1:219
e	<b>Eucalyptus viminalis</b> Labill. subsp. <b>hentyensis</b> Brooker & Slee, J. Adelaide Bot. Gard. 21: 92 (2007)	
	<b>Eucalyptus viminalis</b> Labill. subsp. <b>viminalis</b> , Nov. Holl. Pl. 2: 12, t.151 (1806)	1:214
	<b>Euryomyrtus parviflora</b> Miq., Ned. Kruidk. Arch. 4(1): 149 (1856)	
	<b>Euryomyrtus ramosissima</b> (A.Cunn.) Trudgen, Nuytsia 13(3): 560 (2001)	1:198
	Euryomyrtus ramosissima (A.Cunn.) Trudgen subsp. prostrata (Hook.f.) Trudgen = <b>Euryomyrtus parviflora</b>	1:198
	Fabricia laevigata Gaertn. = <b>Leptospermum laevigatum</b>	
	<b>Kunzea ambigua</b> (Sm.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 629 (1917)	1:202
	Kunzea corifolia (Vent.) Schauer = <b>Kunzea ambigua</b>	
i	<b>Kunzea ericoides</b> (A.Rich.) Joy Thoms., Telopea 2: 379 (1983)	
	Leptospermum flavescens Sm. sensu Rodway (1903) = <b>Leptospermum glaucescens</b>	
	Leptospermum flavescens Sm. var. commune Benth. = <b>Leptospermum glaucescens</b>	
	Leptospermum flavescens Sm. var. grandiflorum (Lodd., G.Lodd & W.Lodd.) Benth. = <b>Leptospermum grandiflorum</b>	
	Leptospermum flavescens Sm. var. nitidum (Hook.f.) Rodway = <b>Leptospermum nitidum</b>	
e	<b>Leptospermum glaucescens</b> S.Schauer, Linnaea 15: 421 (1841)	1:201
e	<b>Leptospermum grandiflorum</b> Lodd., G.Lodd & W.Lodd., G.Lodd. & W.Lodd., Bot. Cab. 6: t.514 (1821)	1:200
	Leptospermum humifusum A.Cunn. ex S.Schauer sensu Curtis (1956) = <b>Leptospermum rupestre</b> nom. illeg.	
	<b>Leptospermum laevigatum</b> (Gaertn.) F.Muell., Cat. Pl. Melb. Bot. Gard. 22 (1858)	1:200
	<b>Leptospermum lanigerum</b> (Sol. ex Aiton) Sm., Trans. Linn. Soc. London 3: 263 (1797)	1:200
	Leptospermum lanigerum (Sol. ex Aiton) Sm. var. montanum Rodway = <b>Leptospermum lanigerum</b>	
	Leptospermum myrtifolium Sieber ex DC. sensu Rodway (1903) = <b>Leptospermum glaucescens</b>	
e	<b>Leptospermum nitidum</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 139 (1856)	1:201
	Leptospermum pilosum Schauer = <b>Leptospermum lanigerum</b>	
e	<b>Leptospermum riparium</b> D.I.Morris, Records of the Queen Victoria Museum 50: 2 (1974)	1:201
	Leptospermum rodwayanum Summerh. & H.F.Comber = <b>Leptospermum grandiflorum</b>	
e	<b>Leptospermum rupestre</b> Hook.f., Icon. Pl. 4: t.308 (1840)	1:202
	<b>Leptospermum scoparium</b> J.R.Forst. & G.Forst., Char. Gen. Pl. 72, t.36, figs.f-l (1775)	1:200
	Leptospermum scoparium J.R.Forst. & G.Forst. var. eximium B.L.Burtt = <b>Leptospermum scoparium</b>	
	Leptospermum sericeum Labill. sensu Curtis (1956) = <b>Leptospermum glaucescens</b>	

	<b>Melaleuca armillaris</b> (Sol. ex Gaertn.) Sm. subsp. <b>armillaris</b> , Trans. Linn. Soc. London 3: 277 (1797)	1:205
	<b>Melaleuca ericifolia</b> Sm., Trans. Linn. Soc. London 3: 276 (1797)	1:204
	<b>Melaleuca gibbosa</b> Labill., Nov. Holl. Pl. 2: 30, t.172 (1806)	1:204
	<b>Melaleuca pallida</b> (Bonpl.) Craven, Novon 16: 472 (2006)	1:203
e	<b>Melaleuca pustulata</b> Hook.f., London J. Bot. 6: 476 bis (1847)	1:205
	<b>Melaleuca squamea</b> Labill., Nov. Holl. Pl. 2: 28, t.168 (1806)	1:204
	<b>Melaleuca squarrosa</b> Donn ex Sm., Trans. Linn. Soc. London 6: 300 (1802)	1:204
n e	<b>Melaleuca virens</b> Craven, Novon 16: 473 (2006)	1:203
i	<b>Syzygium smithii</b> (Poir.) Nied., Nat. Pflanzenfam. [Engler & Prantl] III. 7: 85 (1893)	
	<b>Thryptomene micrantha</b> Hook.f., Hooker's J. Bot. Kew Gard. Misc. 5: 299, t.8 (1853)	1:198
<b>NYCTAGINACEAE</b>		
i	<b>Mirabilis jalapa</b> L., Sp. Pl. 1: 177 (1753)	
<b>OLEACEAE</b>		
		FTO 113
i	<b>Fraxinus angustifolia</b> Vahl subsp. <b>angustifolia</b> , Enum. Pl. [Vahl] 1: 52 (1804)	
i	<b>Ligustrum vulgare</b> L., Sp. Pl. 1: 7 (1753)	3:470
	<b>Notelaea ligustrina</b> Vent., Choix Pl. 25 (1804)	3:470
<b>ONAGRACEAE</b>		
	<i>Boisduvalia tasmanica</i> (Hook.f.) Munz = <b>Epilobium curtisiae</b>	2:234
	<i>Epilobium adenocaulon</i> Hausskn. = <b>Epilobium ciliatum</b>	2:233
	<i>Epilobium alpinum</i> L. nom. illeg., sensu Rodway (1903) = <b>Epilobium tasmanicum</b> (misapplied in Tasmania)	
	<b>Epilobium billardioreanum</b> Ser. ex DC. subsp. <b>billardioreanum</b> , Prodr. [A. P. de Candolle] 3: 41 (1828)	2:232
	<b>Epilobium billardioreanum</b> Ser. ex DC. subsp. <b>cinereum</b> (A.Rich.) P.H.Raven & Engelhorn, New Zealand J. Bot. 9: 349 (1971)	2:230
	<i>Epilobium billardioreanum</i> Ser. ex DC. subsp. <i>hydrophilum</i> P.H.Raven & Engelhorn = <b>Epilobium billardioreanum</b> subsp. <b>billardioreanum</b>	
	<b>Epilobium billardioreanum</b> Ser. ex DC. subsp. <b>intermedium</b> P.H.Raven & Engelhorn, New Zealand J. Bot. 9: 348 (1971)	
i	<b>Epilobium ciliatum</b> Raf., Med. Repos. 5: 361 (1808)	2:233
	<i>Epilobium confertifolium</i> Hook.f. sensu Bentham (1867) = <b>Epilobium tasmanicum</b> (misapplied in Tasmania)	
	<b>Epilobium curtisiae</b> P.H.Raven, Aliso 5: 249, fig.1 (1963)	2:234
	<i>Epilobium erosum</i> Hausskn. = <b>Epilobium gunnianum</b>	2:230
e	<b>Epilobium fugitivum</b> P.H.Raven & Engelhorn, New Zealand J. Bot. 9: 346 (1971)	
	<i>Epilobium glabellum</i> G.Forst. = <b>Epilobium billardioreanum</b>	
	<b>Epilobium gunnianum</b> Hausskn., Oesterr. Bot. Z. 29: 149 (1879)	2:230
	<b>Epilobium hirtigerum</b> A.Cunn., Ann. Nat. Hist. 3: 33 (1839)	2:231
	<i>Epilobium junceum</i> G.Forst. ex Spreng. sensu Curtis (1963) = <b>Epilobium billardioreanum</b> subsp. <b>cinereum</b>	2:230
n i * t	<i>Epilobium nummulariifolium</i> R.Cunn. ex A.Cunn., Ann. Nat. Hist. 3: 31 (1839)	
i t	<b>Epilobium obscurum</b> Schreb., Spic. Fl. Lips. 147, 155 (1771)	
	<b>Epilobium pallidiflorum</b> A.Cunn., Ann. Nat. Hist. 3: 34 (1839)	2:231

e	<b>Epilobium perpusillum</b> Hausskn., Monogr. Epilobium: 301, t.21, fig. 90 (1884)	2:233
i t	<b>Epilobium rotundifolium</b> G.Forst., Fl. Ins. Austr. 27 (1786)	
	<b>Epilobium sarmentaceum</b> Hausskn., Oesterr. Bot. Z. 29: 149 (1879)	2:229
	<b>Epilobium tasmanicum</b> Hausskn., Monogr. Epilobium: 296, t.20, fig.84 (1884)	2:233
	Epilobium tenuipes Hook.f. = <b>Epilobium tasmanicum</b>	
	Epilobium tetragonum L. sensu Bentham (1867) = <b>Epilobium billardioreanum</b> (misapplied in Tasmania)	
	<b>Epilobium willisii</b> P.H.Raven & Engelhorn, New Zealand J. Bot. 9: 347 (1971)	
i	<b>Fuchsia magellanica</b> Lam., Encycl. (Lamarck) 2: 565 (1788)	2:235
i #	<b>Oenothera biennis</b> L., Sp. Pl. 1: 346 (1753)	
i	<b>Oenothera glazioviana</b> Micheli, Fl. Bras. (Martius) 13: 178 (1875)	
	Oenothera stricta Ledeb. ex Link = <b>Oenothera stricta</b> subsp. <b>stricta</b>	2:235
i	<b>Oenothera stricta</b> Ledeb. ex Link subsp. <b>stricta</b> , Enum. Hort. Berol. Alt. 1: 377 (1821)	2:235
	Oenothera tasmanica Hook.f. = <b>Epilobium curtisiae</b>	
<b>OROBANCHACEAE</b>		
	Orobanche australiana F.Muell. recorded in error	
i	<b>Orobanche minor</b> Sm., Engl. Bot. 6: t.422 (1797)	3:533
<b>OXALIDACEAE</b>		
		FTO 70
i	<b>Oxalis articulata</b> Savigny, Encycl. (Lamarck) 4: 686 (1798)	1:102
i	<b>Oxalis corniculata</b> L. subsp. <b>corniculata</b> , Sp. Pl. 1: 435 (1753)	1:101
	<b>Oxalis exilis</b> A.Cunn., Ann. Nat. Hist. 3: 316 (1839)	
i	<b>Oxalis incarnata</b> L., Sp. Pl. 1: 433 (1753)	1:102
	Oxalis lactea Hook. = <b>Oxalis magellanica</b>	1:101
i	<b>Oxalis latifolia</b> Kunth, Nov. Gen. Sp. [H.B.K.] 5: 237, t.467 (1823)	1:101
	<b>Oxalis magellanica</b> G.Forst., Commentat. Soc. Regiae Sci. Gott. 9: 33 (1789)	1:101
	<b>Oxalis perennans</b> Haw., Misc. Nat. 181 (1803)	
i	<b>Oxalis pes-caprae</b> L., Sp. Pl. 1: 434 (1753)	1:101
i	<b>Oxalis purpurea</b> L., Sp. Pl. 1: 433 (1753)	1:102
	<b>Oxalis radicata</b> A.Rich., Tent. Fl. Abyss. 2: 123 (1847)	
	<b>Oxalis rubens</b> Haw., Misc. Nat. 182 (1803)	
<b>PAPAVERACEAE</b>		
		FTO 44
	Argemone mexicana L. sensu Curtis & Morris (1975) = <b>Argemone ochroleuca</b> subsp. <b>ochroleuca</b>	1:28
i	<b>Argemone ochroleuca</b> Sweet subsp. <b>ochroleuca</b> , Brit. Fl. Gard. [Sweet] 3: t.242 (1828)	1:28
i	<b>Eschscholzia californica</b> Cham., Horae Phys. Berol. [Nees]: 73-74, t.15 (1820)	1:29
i	<b>Glaucium flavum</b> Crantz, Stirp. Austr. Fasc. Sp. 2: 133 (1763)	1:28
i	<b>Papaver aculeatum</b> Thunb., Prodr. Pl. Cap. 92 (1800)	1:27
i	<b>Papaver argemone</b> L., Sp. Pl. 1: 506 (1753)	1:27
i	<b>Papaver dubium</b> L., Sp. Pl. 2: 1196 (1753)	1:27
i	<b>Papaver hybridum</b> L., Sp. Pl. 1: 506 (1753)	1:27
i	<b>Papaver rhoeas</b> L., Sp. Pl. 1: 507 (1753)	1:27
i	<b>Papaver somniferum</b> L. subsp. <b>setigerum</b> (DC.) L.Corb., Nouv. Fl. Normandie: 30 (1894)	1:28
i	<b>Papaver somniferum</b> L. subsp. <b>somniferum</b> , Sp. Pl. 1: 508 (1753)	1:28

**PASSIFLORACEAE**

- n ?i # **Passiflora cinnabarina** Lindl., Gard. Chron. 23: 724 (1855)  
 Passiflora mollissima (Kunth) L.H.Bailey sensu Buchanan (1999) = **Passiflora tarminiana**  
 i **Passiflora tarminiana** Coppens & V.E.Barney, Novon 11: 9 (2001)

**PHYTOLACCACEAE**

- i **Phytolacca octandra** L., Sp. Pl., ed. 2: 631 (1762)

**PITTOSPORACEAE**

- Billardiera alpina (McGill.) E.M.Benn. = **Rhytidosporum inconspicuum**  
 Billardiera fusiformis Labill. sensu Buchanan (2009) = **Billardiera heterophylla** 1:59  
 i **Billardiera heterophylla** (Lindl.) L.W.Cayzer & Crisp, Austral. Syst. Bot. 17: 119 (2004) 1:59  
 e **Billardiera longiflora** Labill., Nov. Holl. Pl. 1: 64, t.89 (1805) 1:58  
 Billardiera longiflora Labill. var. alpina Rodway = **Billardiera longiflora**  
 Billardiera longiflora Labill. var. ovalis (Lindl.) E.M.Benn. = **Billardiera ovalis**  
**Billardiera macrantha** Hook.f., Bot. Antarct. Voy. Ill. (Fl. Tasman.) 1: 37 (1855)  
**Billardiera mutabilis** Salisb., Parad. Lond. 1: t.48 (1806) 1:59  
 e **Billardiera nesophila** L.W.Cayzer & D.L.Jones, Austral. Syst. Bot. 17(1): 105 (2004)  
 e **Billardiera ovalis** Lindl., Edwards's Bot. Reg. 20: t.1719 (1834)  
 Billardiera procumbens (Hook.) E.M.Benn. = **Rhytidosporum procumbens** 1:58  
 Billardiera scandens Sm. sensu Curtis & Morris (1975) = **Billardiera mutabilis** 1:59  
 e **Billardiera viridiflora** L.W.Cayzer & D.L.Jones, Austral. Syst. Bot. 17: 105 (2004)  
**Bursaria spinosa** Cav. subsp. **spinosa**, Icon. [Cavanilles] 4: 30, t.350 (1797) 1:57  
 Marianthus procumbens (Hook.) Benth. = **Rhytidosporum procumbens** 1:58  
**Pittosporum bicolor** Hook., J. Bot. (Hooker) 1: 249 (1834) 1:56  
 i **Pittosporum crassifolium** Banks & Sol. ex A.Cunn., Ann. Nat. Hist. 4: 106 (1840)  
 i **Pittosporum eugenioides** A.Cunn., Ann. Nat. Hist. 4: 106 (1840)  
 i **Pittosporum tenuifolium** Sol. ex Gaertn., Fruct. Sem. Pl. 1: 286 (1788)  
 i **Pittosporum undulatum** Vent., Descr. Pl. Nouv. 76, t.76 (1802)  
 i **Pittosporum undulatum** Vent. subsp. **x emmettii** W.M.Curtis, Records of the Queen  
 Victoria Museum 50: 3( 1974) = **Pittosporum bicolor** x **P. undulatum** 1:57  
 Rhytidosporum alpinum McGill. sensu Buchanan (1999) = **Rhytidosporum inconspicuum**  
**Rhytidosporum inconspicuum** L.W.Cayzer, Crisp & I.Telford, Austral. Syst. Bot. 12:  
 700 (1999)  
**Rhytidosporum procumbens** (Hook.) F.Muell., Pl. Victoria 1: 75 (1862) 1:58  
 Sollya heterophylla Lindl. = **Billardiera heterophylla** 1:59

**PLANTAGINACEAE**

- Plantago antarctica** Decne., Prodr. [A. P. de Candolle] 13: 703 (1852) 3:562  
 Plantago archeri Hook.f. = **Plantago tasmanica** var. **archeri**  
 i **Plantago australis** Lam., Tabl. Encycl. 1: 339 (1792)  
 e **Plantago bellidioides** Decne., Prodr. [A. P. de Candolle] 13: 701 (1852) 3:561  
 Plantago brownii Rapin sensu Rodway (1903) = **Plantago triantha**  
 Plantago carnosa R.Br. = **Plantago triantha**  
 i **Plantago coronopus** L. subsp. **commutata** (Guss.) Pilg., Repert. Spec. Nov. Regni Veg. 28:  
 287 (1930) 3:560  
 i **Plantago coronopus** L. subsp. **coronopus**, Sp. Pl. 1: 115 (1753) 3:560  
 e **Plantago daltonii** Decne., Prodr. [A. P. de Candolle] 13: 703 (1852) 3:563

	<b>Plantago debilis</b> R.Br., Prodr. Fl. Nov. Holland. 425 (1810)	3:560
	<b>Plantago gaudichaudii</b> Barnéoud, Monogr. Plantag. 15 (1845)	
e	<b>Plantago glabrata</b> Hook.f., London J. Bot. 6: 276 (1847)	3:562
	<b>Plantago glacialis</b> B.G.Briggs, Carolin & Pulley, Contr. New South Wales Natl. Herb. 4: 395 (1973)	3:564
e	<b>Plantago gunnii</b> Hook.f., London J. Bot. 5: 446 (1846)	3:564
	<b>Plantago hispida</b> R.Br., Prodr. Fl. Nov. Holland. 425 (1810)	3:561
i	<b>Plantago lanceolata</b> L., Sp. Pl. 1: 113 (1753)	3:560
i	<b>Plantago major</b> L., Sp. Pl. 1: 112 (1753)	3:559
	<i>Plantago muelleri</i> Pilg. sensu Curtis (1967) = <b>Plantago glacialis</b>	3:564
e	<b>Plantago paradoxa</b> Hook.f., London J. Bot. 6: 277 (1847)	3:564
e	<b>Plantago tasmanica</b> Hook.f. var. <b>archeri</b> (Hook.f.) W.M.Curtis, The Student's Flora of Tasmania 3: 650 (1967)	3:563
	<i>Plantago tasmanica</i> Hook.f. var. <i>daltonii</i> (Decne.) Hook.f. = <b>Plantago daltonii</b>	
	<i>Plantago tasmanica</i> Hook.f. var. <i>glabrata</i> (Hook.f.) Hook.f. = <b>Plantago glabrata</b>	
e	<b>Plantago tasmanica</b> Hook.f. var. <b>tasmanica</b> , London J. Bot. 6: 276 (1847)	3:562
t	<b>Plantago triantha</b> Spreng., Syst. Veg. (ed. 16) [Sprengel] 1: 439 (1824)	3:563
	<b>Plantago varia</b> R.Br., Prodr. Fl. Nov. Holland. 425 (1810)	3:561
	<b>PLUMBAGINACEAE</b>	FTO 94
	<b>Limonium australe</b> (R.Br.) Kuntze var. <b>australe</b> , Revis. Gen. Pl. 2: 395 (1891)	3:465
e	<b>Limonium australe</b> (R.Br.) Kuntze var. <b>baudinii</b> (Lincz.) A.M.Gray, Kanunnah 4: 117 (2010)	
	<i>Limonium baudinii</i> Lincz. = <b>Limonium australe</b> var. <b>baudinii</b>	
i #	<b>Limonium sinuatum</b> (L.) Mill., Gard. Dict., ed. 8: 6 (1768)	
	<i>Statice australis</i> (R.Br.) Spreng. = <b>Limonium australe</b>	
	<i>Taxanthea australis</i> R.Br. = <b>Limonium australe</b>	
	<b>POLEMONIACEAE</b>	FTO 102
n	<i>Collomia grandiflora</i> Douglas ex Lindl. – previously listed as naturalised but insufficient evidence exists to support this	3:485
	<i>Collomia linearis</i> (Cav.) Nutt. = <i>Collomia grandiflora</i>	3:485
i	<b>Navarretia squarrosa</b> (Eschsch.) Hook. & Arn., Bot. Beechey Voy. 368 (1839)	3:485
	<b>POLYGALACEAE</b>	FTO 60
	<b>Comesperma calymega</b> Labill., Nov. Holl. Pl. 2: 23, t.162 (1806)	1:62
	<b>Comesperma defoliatum</b> F.Muell., Pl. Victoria 1: 189 (1862)	1:63
	<b>Comesperma ericinum</b> DC., Prodr. [A. P. de Candolle] 1: 334 (1824)	1:62
	<b>Comesperma retusum</b> Labill., Nov. Holl. Pl. 2: 22, t.160 (1806)	1:62
	<b>Comesperma volubile</b> Labill., Nov. Holl. Pl. 2: 24, t.163 (1806)	1:62
i	<b>Polygala myrtifolia</b> L., Sp. Pl. 2: 703 (1753)	1:63
i	<b>Polygala vulgaris</b> L., Sp. Pl. 2: 702 (1753)	
	<b>POLYGONACEAE</b>	FTO 95
i	<b>Acetosa sagittata</b> (Thunb.) L.A.S.Johnson & B.G.Briggs, Contr. New South Wales Natl. Herb. 3: 166 (1962)	
i	<b>Acetosella vulgaris</b> Fourr., Ann. Soc. Linn. Lyon, Ser. 2, 17: 145 (1869)	3:587
i	<b>Emex australis</b> Steinh., Ann. Sci. Nat., Bot., sér. 2, 9: 195 (1838)	3:583
i	<b>Fallopia convolvulus</b> (L.) Á.Löve, Taxon 19: 300 (1970)	3:591



i	<b>Fallopia japonica</b> (Houtt.) Ronse Decr., Bot. J. Linn. Soc. 98: 369 (1988)	
	<b>Muehlenbeckia adpressa</b> (Labill.) Meisn., Pl. Vasc. Gen. [Meisner] 2: 227 (1843)	3:592
	Muehlenbeckia adpressa (Labill.) Meisn. var. hastifolia Meisn. = <b>Muehlenbeckia gunnii</b>	
	Muehlenbeckia adpressa (Labill.) Meisn. var. rotundifolia Benth. nom. illeg. = <b>Muehlenbeckia adpressa</b>	
	Muehlenbeckia australis (G.Forst.) Meisn. sensu Buchanan (2009) = <b>Muehlenbeckia adpressa</b>	
	<b>Muehlenbeckia axillaris</b> (Hook.f.) Endl., Gen. Pl. [Endlicher] Suppl. 4(2): 51 (1848)	3:593
	<b>Muehlenbeckia gunnii</b> (Hook.f.) Endl., Gen. Pl. [Endlicher] Suppl. 4(2): 51 (1848)	3:593
i	<b>Persicaria capitata</b> (Buch.-Ham. ex D.Don) H.Gross, Bot. Jahrb. Syst. 49: 277 (1913)	
	<b>Persicaria decipiens</b> (R.Br.) K.L.Wilson, Telopea 3: 178 (1988)	3:590
	<b>Persicaria hydropiper</b> (L.) Delarbre, Fl. Auvergne (Delarbre) 2: 518 (1800)	3:590
i	<b>Persicaria lapathifolia</b> (L.) Gray, Nat. Arr. Brit. Pl. 2: 270 (1821)	3:591
i	<b>Persicaria maculosa</b> Gray, Nat. Arr. Brit. Pl. 2: 269 (1821)	3:591
	<b>Persicaria praetermissa</b> (Hook.f.) H.Hara, Fl. E. Himalaya: 73 (1966)	3:589
?i	<b>Persicaria prostrata</b> (R.Br.) Soják, Preslia 46: 154 (1974)	3:590
	<b>Persicaria subsessilis</b> (R.Br.) K.L.Wilson, Telopea 3: 180 (1988)	3:591
	Polygonum adpressum Labill. = <b>Muehlenbeckia adpressa</b>	
i	<b>Polygonum arenastrum</b> Jord. ex Boreau, Fl. Centre France, ed. 3 [Boreau], 2: 559 (1857)	3:589
i	<b>Polygonum aviculare</b> L., Sp. Pl. 1: 362 (1753)	3:588
	Polygonum convolvulus L. = <b>Fallopia convolvulus</b>	3:591
	Polygonum decipiens R.Br. = <b>Persicaria decipiens</b>	3:590
	Polygonum hydropiper L. = <b>Persicaria hydropiper</b>	3:590
	Polygonum lapathifolium L. = <b>Persicaria lapathifolia</b>	3:591
	Polygonum minus Huds. sensu Bentham (1870) = <b>Persicaria decipiens</b> (misapplied in Tasmania)	
	Polygonum persicaria L. = <b>Persicaria maculosa</b>	3:591
	Polygonum plebeium R.Br. sensu Curtis (1967) = <b>Polygonum arenastrum</b>	3:589
	Polygonum prostratum R.Br. = <b>Persicaria prostrata</b>	3:590
	Polygonum strigosum R.Br. sensu Curtis (1967) = <b>Persicaria praetermissa</b>	3:589
	Polygonum subsessile R.Br. = <b>Persicaria subsessilis</b>	3:591
	Reynoutria japonica Houtt. = <b>Fallopia japonica</b>	
	Rumex acetosella L. = <b>Acetosella vulgaris</b>	3:587
	Rumex angiocarpus Murb. = <b>Acetosella vulgaris</b>	3:586
	<b>Rumex bidens</b> R.Br., Prodr. Fl. Nov, Holland. 421 (1810)	3:586
	<b>Rumex brownii</b> Campd., Monogr. Rumex 64: 81 (1819)	3:585
i	<b>Rumex conglomeratus</b> Murray, Prodr. Stirp. Gott. 52 (1770)	3:585
i	<b>Rumex crispus</b> L., Sp. Pl. 1: 335 (1753)	3:584
	Rumex dumosiformis Rech. = <b>Rumex dumosus</b>	3:586
	<b>Rumex dumosus</b> A.Cunn. ex Meisn., Prodr. [A. P. de Candolle] 14: 62 (1856)	3:586
	Rumex dumosus A.Cunn. ex Meisn. var. dumosiformis (Rech.f.) Rech.f. = <b>Rumex dumosus</b>	3:586
i n	<b>Rumex obtusifolius</b> L., Sp. Pl. 1: 335 (1753)	3:585
n	Rumex obtusifolius L. subsp. transiens (Simonk.) Rech.f. = <b>Rumex obtusifolius</b>	
i	<b>Rumex pulcher</b> L. subsp. <b>pulcher</b> , Sp. Pl. 1: 336 (1753)	3:585
	Rumex sagittatus Thunb. = <b>Acetosa sagittata</b>	
	Rumex sanguineus L. sensu Rodway (1903) = <b>Rumex conglomeratus</b> (misapplied in Tasmania)	

	<b>PORTULACACEAE</b>	FTO 100
	<b>Calandrinia calyptрата</b> Hook.f., Icon. Pl. 3: t.296 (1840)	1:78
	<i>Calandrinia caulescens</i> Kunth sensu Curtis & Morris (1975) = <b>Calandrinia ciliata</b>	1:79
i	<b>Calandrinia ciliata</b> (Ruiz & Pav.) DC., Prodr. [A. P. de Candolle] 3: 359 (1828)	1:79
	<b>Calandrinia eremaea</b> Ewart, Fl. Victoria: 486 (1931)	
	<b>Calandrinia granulifera</b> Benth., Fl. Austral. 1: 176 (1863)	1:78
	<i>Calandrinia menziesii</i> (Hook.) Torr. & A.Gray sensu Buchanan (1999) = <b>Calandrinia ciliata</b>	1:79
	<i>Calandrinia neesiana</i> H.Eichler = <b>Calandrinia granulifera</b>	1:78
	<i>Calandrinia pygmaea</i> F.Muell. nom. illeg. = <b>Calandrinia granulifera</b>	
	<i>Claytonia australasica</i> Hook.f. = <b>Montia australasica</b>	1:79
n	<i>Claytonia perfoliata</i> Donn ex Willd. subsp. <i>perfoliata</i> – previously listed as naturalised but insufficient evidence exists to support this	
	<b>Montia australasica</b> (Hook.f.) Pax & K.Hoffm., Nat. Pflanzenfam., ed. 2 [Engler & Prantl] 16c: 259 (1934)	1:79
	<b>Montia fontana</b> L. subsp. <b>chondrosperma</b> (Fenzl) Walters, Watsonia 3: 4 (1953)	1:79
	<i>Neopaxia australasica</i> (Hook.f.) O.Nilsson = <b>Montia australasica</b>	1:79
	<b>Portulaca oleracea</b> L., Sp. Pl. 1: 445 (1753)	1:78
	<b>PRIMULACEAE</b>	FTO 103 (SAMOLUS)
	<i>Anagallis arvensis</i> L. = <b>Lysimachia arvensis</b>	3:467
	<i>Anagallis arvensis</i> L. subsp. <i>foemina</i> sensu Curtis (1967) = <b>Lysimachia arvensis</b>	3:468
	<i>Anagallis arvensis</i> L. var. <i>arvensis</i> = <b>Lysimachia arvensis</b>	3:468
	<i>Anagallis arvensis</i> L. var. <i>caerulea</i> Gouan = <b>Lysimachia arvensis</b>	3:468
i	<b>Lysimachia arvensis</b> (L.) U.Manns & Anderb., Willdenowia 39: 51 (2009)	3:467
n i *	<i>Lysimachia minima</i> (L.) U.Manns & Anderb., Willdenowia 39: 52 (2009)	
i	<b>Lysimachia nummularia</b> L., Sp. Pl. 1: 148 (1753)	3:467
	<i>Samolus littoralis</i> R.Br. = <b>Samolus repens</b>	
	<b>Samolus repens</b> (J.R.Forst. & G.Forst.) Pers. var. <b>repens</b> , Syn. Pl. (Persoon) 1: 171 (1805)	3:468
	<i>Sheffieldia incana</i> Labill. = <b>Samolus repens</b>	
	<b>PROTEACEAE</b>	FTO 48
e	<b>Agastachys odorata</b> R.Br., Trans. Linn. Soc. London 10: 158 (1810)	3:601
	<i>Banksia australis</i> R.Br. = <b>Banksia marginata</b>	
	<i>Banksia depressa</i> R.Br. = <b>Banksia marginata</b>	
	<i>Banksia insularis</i> R.Br. = <b>Banksia marginata</b>	
x	<b>Banksia integrifolia</b> L.f. subsp. <b>integrifolia</b> , Suppl. Pl. 127 (1782)	3:616
	<b>Banksia marginata</b> Cav., Anales Hist. Nat. 1(3): 227, t.13 (1800)	3:615
	<i>Banksia media</i> R.Br. sensu Hooker (1860) = <b>Banksia serrata</b>	
	<b>Banksia serrata</b> L.f., Suppl. Pl. 126 (1782)	3:616
e	<b>Bellendena montana</b> R.Br., Trans. Linn. Soc. London 10: 166 (1810)	3:600
e	<b>Cenarrhenes nitida</b> Labill., Nov. Holl. Pl. 1: 36, t.50 (1805)	3:601
e	<b>Conospermum hookeri</b> (Meisn.) E.M.Benn., Fl. Australia 16: 485 (1995)	3:600
	<i>Conospermum taxifolium</i> C.F.Gaertn. = <b>Conospermum hookeri</b> (Tasmanian plants)	3:600
	<i>Embothrium tinctorium</i> Labill. = <b>Lomatia tinctoria</b>	
	<i>Embothrium truncatum</i> Labill. = <b>Telopea truncata</b>	
i	<b>Grevillea</b> hybrids	
	<b>Grevillea australis</b> R.Br., Trans. Linn. Soc. London 10: 171 (1810)	3:606

	<i>Grevillea australis</i> R.Br. var. <i>brevifolia</i> Hook.f. = <b>Grevillea australis</b>	3:606
	<i>Grevillea australis</i> R.Br. var. <i>erecta</i> Hook.f. = <b>Grevillea australis</b>	3:606
	<i>Grevillea australis</i> R.Br. var. <i>linearifolia</i> Hook.f. = <b>Grevillea australis</b>	3:606
	<i>Grevillea australis</i> R.Br. var. <i>montana</i> Hook.f. = <b>Grevillea australis</b>	3:606
	<i>Grevillea australis</i> R.Br. var. <i>planifolia</i> Hook.f. = <b>Grevillea australis</b>	3:606
	<i>Grevillea australis</i> R.Br. var. <i>subulata</i> Hook.f. = <b>Grevillea australis</b>	3:606
	<i>Grevillea australis</i> R.Br. var. <i>tenuifolia</i> (R.Br.) Meisn. = <b>Grevillea australis</b>	3:606
	<i>Grevillea seymouriae</i> Sweet ex Meisn. sensu Meisner (1856) attributed in error	
	<i>Grevillea stuartii</i> Meisn. sensu Hooker (1860) = <b>Grevillea australis</b>	
	<i>Grevillea tenuifolia</i> R.Br. = <b>Grevillea australis</b>	
	<i>Hakea acicularis</i> (Sm. ex Vent.) Knight nom. illeg. sensu Hooker (1860) = <b>Hakea decurrens</b> subsp. <b>platytaenia</b> (misapplied in Tasmania)	
	<i>Hakea acicularis</i> (Sm. ex Vent.) Knight var. <i>lissosperma</i> (R.Br.) Benth. = <b>Hakea lissosperma</b>	
	<b>Hakea decurrens</b> R.Br. subsp. <b>physocarpa</b> W.R.Barker, J. Adelaide Bot. Gard. 17: 193 (1996)	3:610
	<b>Hakea decurrens</b> R.Br. subsp. <b>platytaenia</b> W.R.Barker, J. Adelaide Bot. Gard. 17: 196 (1996)	
e	<b>Hakea epiglottis</b> Labill. subsp. <b>epiglottis</b> , Nov. Holl. Pl. 1: 30, t.40 (1805)	3:609
e	<b>Hakea epiglottis</b> Labill. subsp. <b>milliganii</b> (Meisn.) R.M.Barker, Fl. Australia 17B: 394 (1999)	
n i x	<b>Hakea laurina</b> R.Br., Suppl. Prodr. Fl. Nov. Holl. 29 (1830)	
	<b>Hakea lissosperma</b> R.Br., Trans. Linn. Soc. London 10: 180 (1810)	3:611
e	<b>Hakea megadenia</b> R.M.Barker, Aspects of Tasmanian Botany: 83 (1991)	
	<b>Hakea microcarpa</b> R.Br., Trans. Linn. Soc. London 10: 182 (1810)	3:611
	<b>Hakea nodosa</b> R.Br., Trans. Linn. Soc. London 10: 179 (1810)	3:610
	<i>Hakea pugioniformis</i> R.Br. = <b>Hakea teretifolia</b>	
	<i>Hakea rostrata</i> F.Muell. ex Meisn. sensu Curtis (1967) = <b>Hakea epiglottis</b> & <b>H. megadenia</b>	3:609
	<i>Hakea rugosa</i> R.Br. sensu Curtis (1967) = <b>Hakea epiglottis</b> & <b>H. megadenia</b>	3:609
i	<b>Hakea salicifolia</b> (Vent.) B.L.Burtt subsp. <b>salicifolia</b> , Bull. Misc. Inform. Kew 1: 33 (1941)	
	<i>Hakea sericea</i> Schrad. & J.C.Wendl. sensu Curtis (1967) = <b>Hakea decurrens</b> subsp. <b>physocarpa</b>	3:610
	<b>Hakea teretifolia</b> (Salisb.) Britten subsp. <b>hirsuta</b> (Endl.) R.M.Barker, J. Adelaide Bot. Gard. 13: 105 (1990)	3:609
	<b>Hakea ulicina</b> R.Br., Suppl. Prodr. Fl. Nov. Holl. 29 (1830)	3:611
	<i>Hylogyne australis</i> R.Br. = <b>Telopea truncata</b>	
	<b>Isopogon ceratophyllus</b> R.Br., Trans. Linn. Soc. London 10: 72 (1810)	3:599
i #	<b>Lomatia fraseri</b> R.Br., Suppl. Prodr. Fl. Nov. Holl. 34 (1830)	
e	<b>Lomatia polymorpha</b> R.Br., Trans. Linn. Soc. London 10: 200 (1810)	3:613
e	<b>Lomatia tasmanica</b> W.M.Curtis, The Student's Flora of Tasmania 3: 651 (1967)	3:614
e	<b>Lomatia tinctoria</b> (Labill.) R.Br., Trans. Linn. Soc. London 10: 199 (1810)	3:613
e	<b>Orites acicularis</b> (R.Br.) Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 3: 427 (1818)	3:605
e	<b>Orites diversifolius</b> R.Br., Trans. Linn. Soc. London 10: 190 (1810)	3:604
e	<b>Orites milliganii</b> Meisn., Prodr. [A. P. de Candolle] 14: 424 (1856)	3:604
e	<b>Orites revolutus</b> R.Br., Trans. Linn. Soc. London 10: 190 (1810)	3:605
e	<b>Persoonia gunnii</b> Hook.f., London J. Bot. 6: 283 (1847)	
	<i>Persoonia gunnii</i> Hook.f. var. <i>angustifolia</i> Benth. = <b>Persoonia muelleri</b> subsp. <b>angustifolia</b>	3:603
	<i>Persoonia gunnii</i> Hook.f. var. <i>oblanceolata</i> Orchard = <b>Persoonia gunnii</b>	3:603
	<b>Persoonia juniperina</b> Labill., Nov. Holl. Pl. 1: 33, t.45 (1805)	3:603

	Persoonia juniperina Labill. var. brevifolia Meisn. = <b>Persoonia juniperina</b>	3:603
	Persoonia juniperina Labill. var. mollis Orchard = <b>Persoonia juniperina</b>	3:603
	Persoonia juniperina Labill. var. ulicina Meisn. = <b>Persoonia juniperina</b>	3:603
e	<b>Persoonia moscallii</b> Orchard, Brunonia 6: 238, fig.12 (1984)	
e	<b>Persoonia muelleri</b> (P.Parm.) Orchard subsp. <b>angustifolia</b> (Benth.) L.A.S.Johnson & P.H.Weston, Fl. Australia 16: 472 (1995)	3:603
e	<b>Persoonia muelleri</b> (P.Parm.) Orchard subsp. <b>densifolia</b> (Orchard) L.A.S.Johnson & P.H.Weston, Fl. Australia 16: 472 (1995)	
e	<b>Persoonia muelleri</b> (P.Parm.) Orchard subsp. <b>muelleri</b> , Brunonia 6: 226 (1984)	
	Persoonia muelleri (P.Parm.) Orchard var. angustifolia (Benth.) Orchard = <b>Persoonia muelleri</b> subsp. <b>angustifolia</b>	3:603
	Persoonia muelleri (P.Parm.) Orchard var. densifolia Orchard = <b>Persoonia muelleri</b> subsp. <b>densifolia</b>	
	Persoonia muelleri (P.Parm.) Orchard var. muelleri = <b>Persoonia muelleri</b> subsp. <b>muelleri</b>	
e	<b>Telopea truncata</b> (Labill.) R.Br., Trans. Linn. Soc. London 10: 198 (1810)	3:612
	<b>RANUNCULACEAE</b>	FTO 47
n i *	Adonis microcarpa DC., Syst. Nat. [Candolle] 1: 223 (1817)	
e	<b>Anemone crassifolia</b> Hook., Icon. Pl. 3(6): t.257 (1840)	1:11
n i	<b>Aquilegia vulgaris</b> L., Sp. Pl. 1: 533 (1753)	
	Batrachium trichophyllum (Chaix) F.W.Schultz = <b>Ranunculus trichophyllum</b>	1:14
	Caltha introloba F.Muell. sensu Bentham (1863) = <b>Psychrophila phylloptera</b> (misapplied in Tasmania)	
	Caltha novae-zelandiae Hook. sensu Rodway (1903) = <b>Psychrophila phylloptera</b> (misapplied in Tasmania)	
	Caltha phylloptera A.W.Hill = <b>Psychrophila phylloptera</b>	1:20
	<b>Clematis aristata</b> R.Br. ex Ker Gawl., Bot. Reg. 3: t.238 (1817)	1:9
	Clematis aristata R.Br. ex Ker Gawl. subsp. confertissima Kuntze = <b>Clematis aristata</b>	
	Clematis aristata R.Br. ex Ker Gawl. subsp. gentianoides (DC.) Kuntze = <b>Clematis gentianoides</b>	
	Clematis aristata R.Br. ex Ker Gawl. subsp. gunniana Kuntze = <b>Clematis clitorioides</b>	
	Clematis aristata R.Br. ex Ker Gawl. subsp. procumbens Kuntze = <b>Clematis gentianoides</b>	
	Clematis aristata R.Br. ex Ker Gawl. subsp. pubescens (Hügel ex Endl.) Kuntze = <b>Clematis aristata</b>	
	Clematis aristata R.Br. ex Ker Gawl. subsp. tasmanica Kuntze = <b>Clematis gentianoides</b>	
	Clematis aristata R.Br. ex Ker Gawl. var. blanda (Hook.) Benth. = <b>Clematis clitorioides</b>	
	Clematis aristata R.Br. ex Ker Gawl. var. coriacea (DC.) Benth. = <b>Clematis aristata</b>	
	Clematis aristata R.Br. ex Ker Gawl. var. gentianoides (DC.) F.Muell. = <b>Clematis gentianoides</b>	
	Clematis aristata R.Br. ex Ker Gawl. var. gunniana (Kuntze) Domin = <b>Clematis clitorioides</b>	
	Clematis aristata R.Br. ex Ker Gawl. var. minor Hook.f. = <b>Clematis clitorioides</b>	
	Clematis blanda Hook. = <b>Clematis clitorioides</b>	
	<b>Clematis clitorioides</b> DC., Syst. Nat. [Candolle] 1: 158 (1817)	
	Clematis clitorioides DC. var. decipiens Domin = <b>Clematis clitorioides</b>	
	Clematis coriacea DC. = <b>Clematis aristata</b>	
	<b>Clematis decipiens</b> H.Eichler ex Jeanes, Fl. Australia 2: 461 (2007)	
e	<b>Clematis gentianoides</b> DC., Syst. Nat. [Candolle] 1: 159 (1817)	1:10

	Clematis gentianoides DC. var. normalis Domin nom. inval. = <b>Clematis gentianoides</b>	
	Clematis gentianoides var. procumbens = <b>Clematis gentianoides</b>	
	Clematis gentianoides DC. var. tasmanica (Kuntze) Domin = <b>Clematis gentianoides</b>	
	Clematis hexapetala L.f. subsp. brachystemon Kuntze = <b>Clematis clitoroides</b>	
	Clematis linearifolia Steud. sensu Hooker (1860) = <b>Clematis clitoroides</b> (misapplied in Tasmania)	
	<b>Clematis microphylla</b> DC., Syst. Nat. [Candolle] 1: 147 (1817)	1:10
i	<b>Clematis vitalba</b> L. var. <b>vitalba</b> , Sp. Pl. 1: 544 (1753)	1:10
	<b>Myosurus australis</b> F.Muell., Trans. Philos. Soc. Victoria 1: 6 (1854)	1:11
	Myosurus minimus L. sensu Curtis & Morris (1975) = <b>Myosurus australis</b>	1:11
e	<b>Psychophila phylloptera</b> (A.W.Hill) H.Eichler, Fl. Australia 2: 459 (2007)	1:20
t	<b>Ranunculus acaulis</b> Banks & Sol. ex DC., Syst. Nat. [Candolle] 1: 270 (1817)	1:17
n i	<b>Ranunculus acris</b> L. subsp. <b>acris</b> , Sp. Pl. 1: 554 (1753)	1:18
	<b>Ranunculus amphitrichus</b> Colenso, Trans. & Proc. New Zealand Inst. 17: 237 (1885)	1:17
	Ranunculus aquatilis L. sensu Curtis (1956) = <b>Ranunculus trichophyllus</b>	
n	Ranunculus arvensis L. – previously listed as naturalised but insufficient evidence exists to support this	1:18
e	<b>Ranunculus collicola</b> Menadue, Brunonia 8(2): 373 (1986)	
	<b>Ranunculus collinus</b> R.Br. ex DC., Syst. Nat. [Candolle] 1: 271 (1817)	1:17
	Ranunculus concinnus (Hook.f.) Melville = <b>Ranunculus decurvus</b>	1:15
	Ranunculus cuneatus Hook. nom. illeg. = <b>Ranunculus triplodontus</b>	
e	<b>Ranunculus decurvus</b> (Hook.f.) Melville, Kew Bull. 10: 202 (1955)	1:15
	<b>Ranunculus diminutus</b> B.G.Briggs, Telopea 5(4): 583 (1994)	
n	Ranunculus flammula L. subsp. flammula – previously listed as naturalised but insufficient evidence exists to support this	
	<b>Ranunculus glabrifolius</b> Hook., J. Bot. (Hooker) 1: 243 (1834)	1:17
	<b>Ranunculus gunnianus</b> Hook., J. Bot. (Hooker) 1: 244, t.133 (1834)	1:14
	Ranunculus hirtus Banks & Sol. ex DC. nom. illeg. = <b>Ranunculus pimpinellifolius</b>	
	Ranunculus inconspicuus Hook.f. = <b>Ranunculus collinus</b>	
	Ranunculus inundatus R.Br. ex DC. sensu Curtis & Morris (1975) = <b>Ranunculus amphitrichus</b>	1:17
e	<b>Ranunculus jugosus</b> Menadue, Brunonia 8(2): 377 (1986)	
	<b>Ranunculus lappaceus</b> Sm., Cycl. (Rees) 29(2): no.61 (1815)	1:14
	Ranunculus lappaceus Sm. var. nanus (Hook.) Benth. = <b>Ranunculus nanus</b>	
	Ranunculus lappaceus Sm. var. pimpinellifolius (Hook.) Benth. = <b>Ranunculus pimpinellifolius</b>	
	Ranunculus lappaceus Sm. var. scapigerus (Hook.) Benth. = <b>Ranunculus scapiger</b>	
	Ranunculus lappaceus Sm. var. subsericeus Benth. = <b>Ranunculus pascuinus</b>	
	Ranunculus millanii F.Muell. sensu Rodway (1903) = <b>Ranunculus setaceus</b> (misapplied in Tasmania)	
i	<b>Ranunculus muricatus</b> L., Sp. Pl. 1: 555 (1753)	1:18
e	<b>Ranunculus nanus</b> Hook., J. Bot. (Hooker) 1: 242 (1834)	1:16
i	<b>Ranunculus parviflorus</b> L., Syst. Nat., ed. 10, 2: 1087 (1759)	1:19
e	<b>Ranunculus pascuinus</b> (Hook.f.) Melville, Kew Bull. 10: 198 (1955)	1:15
	Ranunculus philonotis Retz. sensu Bentham (1863) = <b>Ranunculus sardous</b> (misapplied in Tasmania)	
	<b>Ranunculus pimpinellifolius</b> Hook., J. Bot. (Hooker) 1: 243 (1834)	1:14

	Ranunculus plebeius R.Br. ex DC. sensu Curtis (1956) mistakenly attributed to Tasmania	
e	<b>Ranunculus prasinus</b> Menadue, Brunonia 8: 375 (1986)	
	<b>Ranunculus pumilio</b> R.Br. ex DC. var. <b>pumilio</b> , Syst. Nat. [Candolle] 1: 271 (1817)	1:19
i	<b>Ranunculus repens</b> L., Sp. Pl. 1: 554 (1753)	1:18
	Ranunculus rivularis Banks & Sol. ex DC. sensu Curtis & Morris (1975) = <b>Ranunculus amphitrichus</b>	1:17
	Ranunculus rivularis Banks & Sol. ex DC. var. inconspicuus Benth. = <b>Ranunculus collinus</b>	
	Ranunculus rivularis Banks & Sol. ex DC. var. major Benth. = <b>Ranunculus glabrifolius</b>	
	Ranunculus rivularis Banks & Sol. ex DC. var. subfluitans Benth. nom. illeg. = <b>Ranunculus amphitrichus</b>	
i	<b>Ranunculus sardous</b> Crantz, Stirp. Austr. Fasc. 2: 84 (1763)	1:19
	<b>Ranunculus scapiger</b> Hook., J. Bot. (Hooker) 1: 224 (1834)	1:15
n	Ranunculus sceleratus L. subsp. sceleratus – previously listed as naturalised but insufficient evidence exists to support this	1:20
	<b>Ranunculus sessiliflorus</b> R.Br. ex DC. var. <b>sessiliflorus</b> , Syst. Nat. [Candolle] 1: 302 (1817)	1:19
e	<b>Ranunculus setaceus</b> Rodway, Pap. & Proc. Roy. Soc. Tasmania 1900-1901: 107 (1902)	1:14
?i	<b>Ranunculus trichophyllus</b> Chaix, Hist. Pl. Dauphiné (Villars) 1: 335 (1786)	1:14
	Ranunculus trichophyllus Chaix subsp. drouetii (F.W.Schultz ex Godr.) A.R.Clapham = <b>Ranunculus trichophyllus</b>	
n i	<b>Ranunculus trilobus</b> Desf., Fl. Atlant. 1: 437, t.113 (1798)	1:19
e	<b>Ranunculus triplodontus</b> Melville, Kew Bull. 10: 204 (1955)	1:16
	Valvaria gentianoides (DC.) Ser. = <b>Clematis gentianoides</b>	
	<b>RESEDACEAE</b>	FTO 84
i	<b>Reseda alba</b> L., Sp. Pl. 1: 449 (1753)	1:53
i	<b>Reseda lutea</b> L., Sp. Pl. 1: 449 (1753)	1:52
i	<b>Reseda luteola</b> L., Sp. Pl. 1: 448 (1753)	1:52
	Reseda odorata L. = <b>Reseda lutea</b> (misapplied in Tasmania)	
	<b>RHAMNACEAE</b>	
e	<b>Cryptandra alpina</b> Hook.f., Bot. Antarct. Voy. Ill. (Fl. Tasman.) 1: 75, t.12B (1855)	1:121
	<b>Cryptandra amara</b> Sm., Cycl. (Rees) 10: no.2 (1808)	1:120
e	<b>Cryptandra exilis</b> D.I.Morris, Aspects of Tasmanian Botany: 57 (1991)	
	Cryptandra mollis Hook.f. = <b>Spyridium parvifolium</b> var. <b>molle</b>	
	Cryptandra obovata Hook.f. = <b>Spyridium obovatum</b>	
	Cryptandra parvifolia Hook.f. = <b>Spyridium parvifolium</b> var. <b>parvifolium</b>	
	Cryptandra pimelioides Hook.f. = <b>Stenanthemum pimeleoides</b>	
	Cryptandra sieberi Fenzl sensu Hooker (1860) = <b>Cryptandra amara</b>	
	Cryptandra tomentosa Lindl. sensu Buchanan et al. (1989) = <b>Cryptandra exilis</b>	
	Cryptandra vexillifera Hook. = <b>Spyridium vexilliferum</b>	
	Discaria australis Hook. nom. illeg. = <b>Discaria pubescens</b>	
	<b>Discaria pubescens</b> (Brongn.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 620 (1917)	1:121
	Pomaderris affinis N.A.Wakef. sensu Curtis (1956) = <b>Pomaderris intermedia</b>	1:114
	<b>Pomaderris apetala</b> Labill. subsp. <b>apetala</b> , Nov. Holl. Pl. 1: 62, t.87 (1805)	1:115
	<b>Pomaderris apetala</b> Labill. subsp. <b>maritima</b> N.G.Walsh & Coates, Muelleria 10: 34 (1997)	
	<b>Pomaderris aspera</b> Sieber ex DC., Prodr. [A. P. de Candolle] 2: 33 (1825)	
	Pomaderris discolor (Vent.) Poir. sensu Hooker (1860) = <b>Pomaderris intermedia</b>	

	<b>Pomaderris elachophylla</b> F.Muell., Fragm. (Mueller) 2: 131 (1861)	1:116
e	<b>Pomaderris elliptica</b> Labill. var. <b>diemenica</b> N.G.Walsh & Coates, Muelleria 10: 51 (1997)	
	<b>Pomaderris elliptica</b> Labill. var. <b>elliptica</b> , Nov. Holl. Pl. 1: 61, t.86 (1805)	1:114
	<i>Pomaderris ferruginea</i> Sieber ex Fenzl sensu Bentham (1863) recorded in error	
	<b>Pomaderris intermedia</b> Sieber ex DC., Prodr. [A. P. de Candolle] 2: 33 (1825)	1:114
	<i>Pomaderris kumeraho</i> A.Cunn. mistakenly attributed to Tasmania	
	<b>Pomaderris oraria</b> F.Muell. ex Reissek subsp. <b>oraria</b> , Linnaea 29: 268 (1858)	1:116
	<b>Pomaderris paniculosa</b> F.Muell. ex Reissek subsp. <b>paralia</b> N.G.Walsh, Muelleria 7: 274 (1990)	
	<b>Pomaderris phyllicifolia</b> Lodd. ex Link subsp. <b>ericoides</b> (Maiden & Betche) N.G.Walsh & Coates, Muelleria 10: 52 (1997)	
	<b>Pomaderris phyllicifolia</b> Lodd. ex Link subsp. <b>phyllicifolia</b> , Enum. Hort. Berol. Alt. 1: 232 (1821)	1:116
	<b>Pomaderris pilifera</b> N.A.Wakef. subsp. <b>pilifera</b> , Vict. Naturalist 68: 140 (1951)	1:115
e	<b>Pomaderris pilifera</b> N.A.Wakef. subsp. <b>talpicutica</b> A.M.Gray & Wapstra, Muelleria 25: 129 (2007)	
	<b>Pomaderris racemosa</b> Hook., J. Bot. (Hooker) 1: 256 (1834-35)	1:116
i	<b>Rhamnus alaternus</b> L., Sp. Pl. 1: 193 (1753)	
	<b>Spyridium eriocephalum</b> Fenzl var. <b>eriocephalum</b> , Enum. Pl. [Endlicher]: 24 (1837)	1:119
e	<b>Spyridium gunnii</b> (Hook.f.) Benth., Fl. Austral. 1: 429 (1863)	1:118
e	<b>Spyridium lawrencei</b> (Hook.f.) Benth., Fl. Austral. 1: 430 (1863)	1:119
	<i>Spyridium microphyllum</i> (F.Muell. ex Reissek) Druce = <b>Spyridium lawrencei</b>	1:119
e	<b>Spyridium obcordatum</b> (Hook.f.) W.M.Curtis, Vict. Naturalist 87: 251 (1970)	1:117
	<i>Spyridium obovatum</i> (Hook.f.) Benth. var. <i>gunnii</i> (Hook.f.) Rodway = <b>Spyridium gunnii</b>	
e	<b>Spyridium obovatum</b> (Hook.) Benth. var. <b>obovatum</b> , Fl. Austral. 1: 429 (1863)	1:118
e	<b>Spyridium obovatum</b> (Hook.) Benth. var. <b>velutinum</b> (F.Muell. ex Reissek) Benth., Fl. Austral. 1: 429 (1863)	1:118
e	<b>Spyridium parvifolium</b> (Hook.) F.Muell. var. <b>molle</b> (Hook.f.) Benth., Fl. Austral. 1: 428 (1863)	1:118
	<b>Spyridium parvifolium</b> (Hook.) F.Muell. var. <b>parvifolium</b> , Fragm. (Mueller) 3: 79 (1862)	1:118
	<i>Spyridium serpyllaceum</i> F.Muell. nom. illeg. = <b>Spyridium obcordatum</b>	
e	<b>Spyridium ulicinum</b> (Hook.) Benth., Fl. Austral. 1: 434 (1863)	1:119
	<b>Spyridium vexilliferum</b> (Hook.) Reissek var. <b>vexilliferum</b> , Linnaea 29: 285 (1858)	1:119
e	<b>Stenanthemum pimeleoides</b> (Hook.f.) Benth., Fl. Austral. 1: 436 (1863)	1:120
<b>ROSACEAE</b>		
	<i>Acaena agnipila</i> Gand. var. <i>aequispina</i> Orchard = <b>Acaena ovina</b>	1:176
	<i>Acaena agnipila</i> Gand. var. <i>tenuispica</i> (Bitter) Orchard = <b>Acaena ovina</b>	1:176
	<i>Acaena anserinifolia</i> (J.R.Forst. & G.Forst.) Druce sensu Curtis (1956) = <b>Acaena novae-zelandiae</b>	
	<b>Acaena × anserovina</b> Orchard, Trans. Roy. Soc. South Australia 93: 104 (1969)	1:176
	<b>Acaena echinata</b> Nees, Pl. Preiss. [J.G.C.Lehman] 1: 95 (1844)	1:175
	<i>Acaena echinata</i> Nees var. <i>retrorsumpilosa</i> (Bitter) Orchard = <b>Acaena echinata</b>	1:175
	<i>Acaena echinata</i> Nees var. <i>subglabrallyx</i> (Bitter) Orchard = <b>Acaena echinata</b>	1:175
	<i>Acaena echinata</i> Nees var. <i>tylacantha</i> Orchard = <b>Acaena echinata</b>	1:175
e	<b>Acaena montana</b> Hook.f., London J. Bot. 6: 476 bis (1847)	1:175
	<b>Acaena novae-zelandiae</b> Kirk, Trans. Proc. New Zealand Inst. 3: 177 (1871)	1:175
	<b>Acaena ovina</b> A.Cunn., Geog. Mem. New South Wales [Field]: 358 (1825)	1:175
	<i>Acaena ovina</i> A.Cunn. var. <i>velutina</i> Orchard = <b>Acaena ovina</b>	1:175

- Acaena pallida** (Kirk) Allan, Fl. New Zealand 1: 360 (1961)  
*Acaena sanguisorbae* (L.) Vahl sensu Bentham (1864) = **Acaena novae-zelandiae** (misapplied in Tasmania)  
*Acaena sanguisorbae* (L.) Vahl var. *montana* (Hook.f.) Hook.f. = **Acaena montana**  
*Alchemilla arvensis* (L.) Scop. = **Aphanes arvensis**
- i **Aphanes arvensis** L., Sp. Pl. 1: 123 (1753) 1:174  
**Aphanes australiana** (Rothm.) Rothm., Bull. Misc. Inform. Kew 1938: 270 (1938)  
*Aphanes inexpectata* W.Lippert sensu Buchanan (1999) = **Aphanes microcarpa**
- i **Aphanes microcarpa** (Boiss. & Reut.) Rothm., Repert. Spec. Nov. Regni Veg. 42: 172 (1937)  
?i **Argentina anserina** (L.) Rydb., Mem. Dept. Bot. Columbia Coll. 2: 159 (1898) 1:173
- i **Cotoneaster franchetii** Bois, Rev. Hort. [Paris] 74: 379 (1902)  
i **Cotoneaster glaucophyllus** Franch. var. **serotinus** (Hutch.) L.T.Lu & Brach, Novon 12: 495 (2002)  
i **Cotoneaster pannosus** Franch., Pl. Delavay. 223 (1890)  
i **Cotoneaster simonsii** Baker, Refug. Bot. [Saunders] 1: t.55 (1869)  
*Cotoneaster symondsii* T.Moore = **Cotoneaster simonsii**
- i **Crataegus monogyna** Jacq., Fl. Austriac. (Jacquin) 3: 50 (1775) 1:178  
*Geum renifolium* F.Muell. = **Geum talbotianum**
- e **Geum talbotianum** W.M.Curtis, Records of the Queen Victoria Museum 50: 4 (1974) 1:172  
x **Geum urbanum** L. var. **strictum** Hook.f., Bot. Antarct. Voy. III (Fl. Tasman.) 1: 114 (1856) 1:172  
*Malus × domestica* Borkh. = **Malus pumila**
- i **Malus pumila** Mill., Gard. Dict., ed. 8: Malus no 3 (1768)  
i **Potentilla anglica** Laichard., Veg. Eur. 1: 475 (1790) 1:173  
*Potentilla anserina* L. = **Argentina anserina** 1:173
- i **Potentilla recta** L., Sp. Pl. 1: 497 (1753) 1:173  
i **Potentilla reptans** L., Sp. Pl. 1: 499 (1753) 1:173  
*Poterium polygamum* Waldst. & Kit. = **Sanguisorba minor** 1:176  
*Poterium sanguisorba* L. sensu Rodway (1903) = **Sanguisorba minor** 1:176
- i **Prunus domestica** L. subsp. **insititia** (L.) Bonnier & Layens, Fl. France [Rouy & Foucard]: 95 (1894) 1:178  
*Prunus insititia* L. = **Prunus domestica** subsp. **insititia**
- i **Prunus laurocerasus** L., Sp. Pl. 1: 474 (1753)  
i **Prunus spinosa** L., Sp. Pl. 1: 475 (1753) 1:178  
i **Rosa canina** L., Sp. Pl. 1: 491 (1753) 1:177  
i **Rosa rubiginosa** L., Mant. Pl. 2: 564 (1771) 1:177
- i **Rubus sp. Tasmania (J.R.Hosking 1551) SA Herbarium**  
i **Rubus anglocandicans** A.Newton, Watsonia 11: 243 (1977) 1:171  
i **Rubus echinatus** Lindl., Syn. Brit. Fl. 94 (1829)  
i **Rubus erythrops** Edees & A.Newton, Watsonia 12: 135 (1978)  
*Rubus fruticosus* L. a catch-all name for all the blackberry species of **Rubus**
- e **Rubus gunnianus** Hook., Icon. Pl. 3: t.291 (1840) 1:171  
i **Rubus laciniatus** Willd., Hort. Berol. [Willdenow] 2(8): t.82 (1806) 1:171  
i **Rubus leucostachys** Schleich. ex Sm., Engl. Fl. 2: 403 (1824) 1:171  
i **Rubus loganobaccus** L.H.Bailey, Gentes Herb. 1: 155 (1923)  
*Rubus macropodus* Ser. ex DC. sensu Hooker (1860) = **Rubus parvifolius**  
**Rubus parvifolius** L., Sp. Pl. 2: 1197 (1753) 1:171



n i	<b>Rubus philadelphicus</b> Blanch., Torreyia 7: 56 (1907)	
i	<b>Rubus polyanthemus</b> Lindeb., Bot. Not. 1883: 105 (1883)	
n i	<b>Rubus rubritinctus</b> W.C.R.Watson, Watsonia 3: 287 (1956)	
i #	<b>Rubus rugosus</b> Sm., Cycl. (Rees) 30: no 34 (1815)	
i	<b>Rubus vestitus</b> Weihe, Comp. Fl. German. 1: 684 (1825)	
i	<b>Sanguisorba minor</b> Scop., Fl. Carniol., ed. 2, 1: 110 (1771)	1:176
	Sanguisorba minor Scop. subsp. muricata (Spach) Briq. = <b>Sanguisorba minor</b>	1:176
i	<b>Sorbus aucuparia</b> L., Sp. Pl. 1: 477 (1753)	

## RUBIACEAE

	Asperula charophyton Airy Shaw & Turrill sensu Curtis (1963) recorded in error	2:276
	<b>Asperula conferta</b> Hook.f., London J. Bot. 6: 464 bis (1847)	2:275
	Asperula conferta Hook.f. var. abbreviata Airy Shaw & Turrill = <b>Asperula conferta</b>	2:275
	Asperula conferta Hook.f. var. scoparioides Airy Shaw & Turrill = <b>Asperula conferta</b>	2:275
	<b>Asperula gunnii</b> Hook.f., London J. Bot. 6: 464 bis (1847)	2:275
	Asperula gunnii Hook.f. var. curta (Hook.f.) Airy Shaw & Turrill = <b>Asperula gunnii</b>	2:276
	Asperula gunnii Hook.f. var. pusilla (Hook.f.) Benth. = <b>Asperula pusilla</b>	
	<b>Asperula minima</b> Hook.f., London J. Bot. 6: 464 bis (1847)	2:276
	<b>Asperula oblanceolata</b> I.Thomps., Muellera 27: 68 (2009)	
	Asperula oligantha F.Muell. nom. illeg., sensu Rodway (1903) a catch-all name that includes most Tasmanian <b>Asperula</b> species	
	<b>Asperula pusilla</b> Hook.f., London J. Bot. 6: 464 bis (1847)	2:276
	<b>Asperula scoparia</b> Hook.f. subsp. <b>scoparia</b> , London J. Bot. 6: 463 bis (1847)	2:275
	<b>Asperula subsimplex</b> Hook.f., London J. Bot. 6: 463 bis (1847)	2:276
	Canthium quadrifidum Labill. = <b>Coprosma quadrifida</b>	
	Coprosma billardierei Hook.f. nom. illeg. = <b>Coprosma quadrifida</b>	
	<b>Coprosma hirtella</b> Labill., Nov. Holl. Pl. 1: 70, t.95 (1805)	2:268
	<b>Coprosma moorei</b> F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania 1893: 179, t.1 (1894)	2:269
	<b>Coprosma nitida</b> Hook.f., London J. Bot. 6: 465 bis (1847)	2:269
	<b>Coprosma nivalis</b> W.R.B.Oliv., Bull. Bernice P. Bishop Mus. 132: 37 (1935)	
	<b>Coprosma perpusilla</b> Colenso subsp. <b>perpusilla</b> , Trans. & Proc. New Zealand Inst. 22: 466 (1890)	
	<b>Coprosma pumila</b> Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.). 2: 543 (1847)	2:269
	<b>Coprosma quadrifida</b> (Labill.) B.L.Rob., Proc. Amer. Acad. Arts 45: 409 (1910)	2:268
i	<b>Coprosma repens</b> A.Rich., Voy. Astrolabe I: 264 (1832)	
i	<b>Coprosma robusta</b> Raoul, Ann. Sci. Nat., Bot., sér. 3, 2: 121 (1844)	
e	<b>Galium albescens</b> Hook.f., London J. Bot. 6: 462 bis (1847)	2:273
i	<b>Galium aparine</b> L., Sp. Pl. 1: 108 (1753)	2:273
	<b>Galium australe</b> DC., Prodr. [A. P. de Candolle] 4: 608 (1830)	2:273
	<b>Galium binifolium</b> N.A.Wakef. subsp. <b>conforme</b> I.Thomps., Muellera 27: 88 (2009)	
	<b>Galium ciliare</b> Hook.f. subsp. <b>ciliare</b> , London J. Bot. 6: 461 bis (1847)	2:272
	<b>Galium ciliare</b> Hook.f. subsp. <b>terminale</b> I.Thomps., Muellera 27: 82 (2009)	
	<b>Galium compactum</b> Ehrend. & McGill., Telopea 2: 370 (1983)	
	<b>Galium densum</b> Hook.f., London J. Bot. 6: 461 bis (1847)	
i	<b>Galium divaricatum</b> Pourr. ex Lam., Encycl. (Lamarck) 2: 580 (1788)	
	<b>Galium gaudichaudii</b> DC. subsp. <b>parviflorum</b> I.Thomps., Muellera 27: 98 (2009)	2:272
	<b>Galium leiocarpum</b> I.Thomps., Muellera 27: 79 (2009)	

i	<b>Galium murale</b> (L.) All., Fl. Pedem. 1: 8 (1785)	2:273
i	<b>Galium palustre</b> L., Sp. Pl. 1: 105 (1753)	
	<i>Galium squalidum</i> Hook.f. = <b>Galium australe</b>	
n	<i>Galium tricornutum</i> Dandy – previously listed as naturalised but insufficient evidence exists to support this	
	<i>Galium umbrosum</i> G.Forst. nom. illeg., sensu Rodway (1903) a catch-all name that includes all glabrous-fruited Tasmanian <b>Galium</b>	
	<i>Galium vagans</i> Hook.f. = <b>Galium binifolium</b> subsp. <b>conforme</b> (uncertain)	
i # t	<b>Galium verum</b> L., Sp. Pl. 1: 107 (1753)	
	<i>Nertera depressa</i> Banks & Sol. ex Gaertn. = <b>Nertera granadensis</b>	2:270
	<b>Nertera granadensis</b> (Mutis ex L.f.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 637 (1917)	2:270
	<b>Opercularia ovata</b> Hook.f., London J. Bot. 6: 465 bis (1847)	2:270
	<b>Opercularia varia</b> Hook.f., London J. Bot. 6: 466 bis (1847)	2:271
i	<b>Sherardia arvensis</b> L., Sp. Pl. 1: 102 (1753)	2:277
	<b>RUTACEAE</b>	FTO 87
e	<b>Acradenia frankliniae</b> Kippist, Proc. Linn. Soc. London 2: 201 (1853)	1:106
	<i>Boronia anemonifolia</i> A.Cunn. subsp. <i>anemonifolia</i> recorded in error	
	<b>Boronia anemonifolia</b> A.Cunn. subsp. <b>variabilis</b> (Hook.) P.G.Neish, Muelleria 14: 11 (2000)	1:105
	<i>Boronia anemonifolia</i> A.Cunn. var. <i>dentigera</i> (F.Muell.) Benth. = <i>Boronia anemonifolia</i> subsp. <i>anemonifolia</i>	
e	<b>Boronia citriodora</b> Gunn ex Hook.f. subsp. <b>citriodora</b> , Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 68 (1855)	1:105
e	<b>Boronia citriodora</b> Gunn ex Hook.f. subsp. <b>orientalis</b> Duretto, Muelleria 17: 83 (2003)	
e	<b>Boronia citriodora</b> Gunn ex Hook.f. subsp. <b>paulwilsonii</b> Duretto, Muelleria 17: 81 (2003)	
e	<b>Boronia elisabethiae</b> Duretto, Muelleria 17: 88 (2003)	
e	<b>Boronia gunnii</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 68, t.10 (1855)	
e	<b>Boronia hemichiton</b> Duretto, Muelleria 17: 87 (2003)	
e	<b>Boronia hippopala</b> Duretto, Muelleria 17: 84 (2003)	
	<b>Boronia nana</b> Hook. var. <b>hyssoipifolia</b> Melville, Kew Bull. 9: 463 (1954)	1:105
	<b>Boronia nana</b> Hook. var. <b>nana</b> , Icon. Pl. 3: t.270 (1840)	1:105
	<b>Boronia parviflora</b> Sm., Tracts Nat. Hist. 295: t.6 (1798)	1:105
	<i>Boronia pilonema</i> Labill. = <b>Boronia parviflora</b>	
	<b>Boronia pilosa</b> Labill. subsp. <b>pilosa</b> , Nov. Holl. Pl. 1: 97, t.124 (1805)	1:105
e	<b>Boronia pilosa</b> Labill. subsp. <b>tasmanensis</b> Duretto, Muelleria 17: 97 (2003)	
	<i>Boronia pilosa</i> Labill. var. <i>floribunda</i> Hook. = <b>Boronia pilosa</b> subsp. <b>pilosa</b>	1:105
	<i>Boronia pilosa</i> Labill. var. <i>laricifolia</i> (Hook.) Hook.f. = <b>Boronia pilosa</b> subsp. <b>pilosa</b>	1:105
	<i>Boronia pinnata</i> Sm. var. <i>citriodora</i> (Gunn ex Hook.f.) Rodway = <b>Boronia citriodora</b>	
	<i>Boronia pinnata</i> Sm. var. <i>gunnii</i> (Hook.f.) Benth. = <b>Boronia gunnii</b>	
	<i>Boronia polygalifolia</i> Sm. sensu Bentham (1863) = <b>Boronia nana</b> var. <b>nana</b> (misapplied in Tasmania)	
	<i>Boronia polygalifolia</i> Sm. var. <i>trifoliata</i> Benth. = <b>Boronia nana</b> var. <b>nana</b>	
	<b>Boronia rhomboidea</b> Hook., Icon. Pl. 8: t.722 (1845)	1:105
e	<b>Boronia rozefeldsii</b> Duretto, Muelleria 17: 101 (2003)	
	<i>Boronia tetrandra</i> Labill. var. <i>grandiflora</i> Hook. = <b>Boronia gunnii</b>	

	Boronia variabilis Hook. = <b>Boronia anemonifolia</b> subsp. <b>variabilis</b>	
i	<b>Coleonema pulchellum</b> I.Williams, J. S. African Bot. 47: 89 (1981)	
	<b>Correa alba</b> Andrews var. <b>alba</b> , Bot. Repos. 1: t.18 (1798)	1:109
e	<b>Correa alba</b> Andrews var. <b>rotundifolia</b> DC., Prodr. [A. P. de Candolle] 1: 719 (1824)	1:109
	<b>Correa backhouseana</b> Hook. var. <b>backhouseana</b> , J. Bot. (Hooker) 1: 253 (1834)	1:110
e	<b>Correa lawrenceana</b> Hook. var. <b>ferruginea</b> Hook.f., Bot. Antarct. Voy. III (Fl. Tasman.) 1: 62 (1855)	
e	<b>Correa lawrenceana</b> Hook. var. <b>lawrenceana</b> , J. Bot. (Hooker) 1: 254 (1834)	1:111
	Correa lawrenciana Hook. var. <b>glabra</b> Benth. = <b>Correa lawrenceana</b> var. <b>lawrenceana</b>	
e	<b>Correa reflexa</b> (Labill.) Vent. var. <b>nummulariifolia</b> (Hook.f.) Paul G.Wilson, Trans. Roy. Soc. South Australia 85: 30 (1961)	1:110
	<b>Correa reflexa</b> (Labill.) Vent. var. <b>reflexa</b> , Jard. Malmaison 1: t.13 (1803)	1:110
	Correa rufa (Labill.) Vent. sensu Brown (1810) = <b>Correa alba</b>	
	Correa speciosa Donn ex Andrews = <b>Correa reflexa</b>	
	Correa speciosa Donn ex Andrews var. <b>backhouseana</b> (Hook.) Benth. = <b>Correa backhouseana</b>	
	Correa speciosa Donn ex Andrews var. <b>normalis</b> Benth. nom. inval., nom. superfl. = <b>Correa reflexa</b>	
	Eriostemon daviesii (Hook.f.) F.Muell. = <b>Phebalium daviesii</b>	
	Eriostemon hillebrandii F.Muell. nom. illeg., sensu Rodway (1903) = <b>Leionema bilobum</b>	
	Eriostemon montanus (Hook.f.) F.Muell. = <b>Leionema montanum</b>	
	Eriostemon obovalis A.Cunn. sensu Bentham (1863) = <b>Philothea verrucosa</b> (misapplied in Tasmania)	
	Eriostemon oldfieldii F.Muell. = <b>Leionema oldfieldii</b>	
	Eriostemon squameus Labill. = <b>Nematolepis squamea</b>	
	Eriostemon verrucosus A.Rich. = <b>Philothea verrucosa</b>	1:107
	Eriostemon virgatus Hook.f. = <b>Philothea virgata</b>	1:107
e	<b>Leionema bilobum</b> (Lindl.) Paul G.Wilson subsp. <b>truncatum</b> (Hook.f.) Duretto & K.L.Durham, Muelleria 23: 11 (2006)	1:108
e	<b>Leionema montanum</b> (Hook.) Paul G.Wilson, Nuytsia 12: 275 (1998)	1:108
e	<b>Leionema oldfieldii</b> (F.Muell.) Paul G.Wilson, Nuytsia 12: 276 (1998)	1:108
e	<b>Nematolepis squamea</b> (Labill.) Paul G.Wilson subsp. <b>retusa</b> (Hook.) Paul G.Wilson, Nuytsia 12: 279 (1998)	1:109
	<b>Nematolepis squamea</b> (Labill.) Paul G.Wilson subsp. <b>squamea</b> , Nuytsia 12: 279 (1998)	1:109
	Phebalium billardierei A.Juss. = <b>Nematolepis squamea</b>	
	Phebalium bilobum Lindl. = <b>Leionema bilobum</b>	1:108
e	<b>Phebalium daviesii</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 358 (1859)	1:108
	Phebalium glandulosum Hook. var. <b>daviesii</b> (Hook.f.) Benth. = <b>Phebalium daviesii</b>	
	Phebalium montanum Hook. = <b>Leionema montanum</b>	1:108
	Phebalium oldfieldii (F.Muell.) Benth. = <b>Leionema oldfieldii</b>	1:108
	Phebalium squameum (Labill.) Engl. = <b>Nematolepis squamea</b>	1:109
	Phebalium truncatum Hook.f. = <b>Leionema bilobum</b> subsp. <b>truncatum</b>	
e	<b>Philothea freyciana</b> Rozefelds, Muelleria 15: 23 (2001)	
	<b>Philothea verrucosa</b> (A.Rich.) Paul G.Wilson, Nuytsia 12: 260 (1998)	1:107
	<b>Philothea virgata</b> (Hook.f.) Paul G.Wilson, Nuytsia 12: 260 (1998)	1:107
	<b>Zieria arborescens</b> Sims subsp. <b>arborescens</b> , Bot. Mag. 34: t.1395 (1811)	1:103
	Zieria cytisoides Sm. sensu Curtis & Morris (1975) = <b>Zieria littoralis</b>	

	<b>Zieria littoralis</b> J.A.Armstr., Austral. Syst. Bot. 15: 389 (2002)	1:103
	Zieria smithii Jacks. sensu Bentham (1863) = <b>Zieria arborescens</b> subsp. <b>arborescens</b> (misapplied in Tasmania)	
	Zieria smithii Jacks. var. macrophylla (Bonpl.) Benth. = <b>Zieria arborescens</b> subsp. <b>arborescens</b>	
	<b>Zieria veronicea</b> (F.Muell.) Benth. subsp. <b>veronicea</b> , Fl. Austral. 1: 305 (1863)	1:103
	<b>SALICACEAE</b>	FTO 80
i	<b>Populus alba</b> L., Sp. Pl. 2: 1034 (1753)	
n i *	Salix alba L. var. vitellina (L.) Stokes, Bot. Mat. Med. 4: 506 (1812)	
	Salix alba × fragilis sensu Curtis (1967) = <b>Salix fragilis</b> nothovar. <b>fragilis</b>	3:648
	Salix atrocinnerea Brot. sensu Curtis (1967) = <b>Salix reichardtii</b>	3:649
	Salix babylonica L. sensu Curtis (1967) = Salix × pendulina nothovar. pendulina	3:648
n i *	Salix × calodendron Wimm., Sal. Eur. (Wimm.): 187 (1866)	
i	<b>Salix cinerea</b> L. subsp. <b>cinerea</b> , Sp. Pl. 2: 1021 (1753)	3:649
i	<b>Salix cinerea</b> L. subsp. <b>oleifolia</b> (Sm.) Macreight, Man. Brit. Bot. (Macreight): 212 (1837)	3:649
i	<b>Salix × fragilis</b> L. nothovar. <b>fragilis</b> , Sp. Pl. 2: 1017 (1753)	3:648
n i *	Salix matsudana Koidz. cv. tortuosa, Bot. Mag. (Tokyo) 29: 312 (1915)	
	Salix × pendulina Wender. nothovar. pendulina – not considered naturalised	
i #	<b>Salix purpurea</b> L., Sp. Pl. 2: 1017 (1753)	
i	<b>Salix × reichardtii</b> A.Kern., Verh. Zool.-Bot. Ges. Wien 10: 249 (1860)	3:649
n i *	Salix × rubens Schrank, Baier. Fl. 1: 226 (1789)	3:648
n i *	Salix × sepulcralis Simonk. nothovar. chrysocoma (Dode) Meikle, Watsonia 15: 274 (1985)	
	<b>SANTALACEAE</b>	FTO 91
	<b>Exocarpos cupressiformis</b> Labill., Voy. Rech. Perouse 1: 156, t.14 (1800)	3:628
e	<b>Exocarpos humifusus</b> R.Br., Prodr. Fl. Nov. Holland. 356 (1810)	3:629
	<b>Exocarpos nanus</b> Hook.f., London J. Bot. 6: 281 (1847)	3:629
	<b>Exocarpos strictus</b> R.Br., Prodr. Fl. Nov. Holland. 357 (1810)	3:628
	<b>Exocarpos syrticola</b> (F.Muell. ex Miq.) Stauffer, Mitt. Bot. Mus. Univ. Zurich 213: 173 (1959)	3:628
	Leptomeria billardierei R.Br. = <b>Leptomeria drupacea</b>	3:626
	<b>Leptomeria drupacea</b> (Labill.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916: 632 (1917)	3:626
e	<b>Leptomeria glomerata</b> F.Muell. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 370 (1859)	3:627
x	<b>Thesium australe</b> R.Br., Prodr. Fl. Nov. Holland. 353 (1810)	3:626
	Thesium drupaceum Labill. = <b>Leptomeria drupacea</b>	
	<b>SAPINDACEAE</b>	FTO 86
	Dodonaea ericifolia G.Don = <b>Dodonaea filiformis</b>	1:122
e	<b>Dodonaea filiformis</b> Link, Enum. Hort. Berol. Alt. 1: 381 (1821)	1:122
	<b>Dodonaea viscosa</b> Jacq. subsp. <b>spatulata</b> (Sm.) J.G.West, Brunonia 7: 43 (1984)	1:122
	<b>SCROPHULARIACEAE</b>	
i #	<b>Antirrhinum majus</b> L., Sp. Pl. 2: 617 (1753)	
i	<b>Bartsia trixago</b> L., Sp. Pl. 2: 602 (1753)	
	Bellardia trixago (L.) All. = <b>Bartsia trixago</b>	
	Celsia cretica L. = <b>Verbascum creticum</b>	3:513
	Chionohebe ciliolata (Hook.f.) B.G.Briggs & Ehrend. = <b>Veronica ciliolata</b> subsp. <b>fiordensis</b>	
i	<b>Cymbalaria muralis</b> G.Gaertn., B.Mey. & Scherb., Oekon. Fl. Wetterau 2: 397 (1800)	3:514

	Derwentia derwentiana (Andrews) B.G.Briggs & Ehrend. subsp. derwentiana = <b>Veronica derwentiana</b> subsp. <b>derwentiana</b>	3:523
	Derwentia nivea (Lindl.) B.G.Briggs & Ehrend. = <b>Veronica nivea</b>	3:523
i	<b>Digitalis purpurea</b> L., Sp. Pl. 2: 621 (1753)	3:519
i	<b>Erythranthe moschata</b> (Douglas ex Lindl.) G.L.Nesom, Phytoneuron 39: 38 (2012)	
e	<b>Euphrasia</b> sp. <b>Bivouac Bay (W.R.Barker 7626 et al.) W.R.Barker</b>	
	Euphrasia alpina R.Br. nom. illeg. – a name of uncertain application	
	Euphrasia alpina R.Br. var. angustifolia Benth. sensu Hooker (1860) = <b>Euphrasia collina</b> subsp. <b>collina</b>	
	Euphrasia alpina R.Br. var. humilis Benth. sensu Hooker (1860) = <b>Euphrasia striata</b>	
e	<b>Euphrasia amphisysepala</b> W.R.Barker, J. Adelaide Bot. Gard. 10: 204 (1987)	
e	<b>Euphrasia amplidens</b> W.R.Barker, J. Adelaide Bot. Gard. 26: 23 (2013)	
	Euphrasia brownii F.Muell. nom. illeg., sensu Rodway (1903) a catch-all name that includes several <b>Euphrasia</b> species	
	Euphrasia brownii F.Muell. var. striata (R.Br.) Rodway = <b>Euphrasia striata</b>	3:530
	<b>Euphrasia collina</b> R.Br. subsp. <b>collina</b> , Prodr. Fl. Nov. Holland. 436 (1810)	3:527
e	<b>Euphrasia collina</b> R.Br. subsp. <b>deflexifolia</b> (Gand.) W.R.Barker, The Endemic Flora of Tasmania 6: 477 (1978)	
e	<b>Euphrasia collina</b> R.Br. subsp. <b>diemenica</b> (Spreng.) W.R.Barker, The Endemic Flora of Tasmania 6: 477 (1978)	3:528
e	<b>Euphrasia collina</b> R.Br. subsp. <b>Dukes Marshes (A.Moscal 861) Tas Herbarium</b>	
e	<b>Euphrasia collina</b> R.Br. subsp. <b>gunnii</b> (Du Rietz) W.R.Barker, J. Adelaide Bot. Gard. 5: 199 (1982)	3:528
e	<b>Euphrasia collina</b> R.Br. subsp. <b>Northwest Tasmania (M.Visoiu 216) Tas Herbarium</b>	
e	<b>Euphrasia collina</b> R.Br. subsp. <b>tasmanica</b> (Gand.) W.R.Barker, J. Adelaide Bot. Gard. 21: 93 (2007)	
	<b>Euphrasia collina</b> R.Br. subsp. <b>tetragona</b> (R.Br.) W.R.Barker, J. Adelaide Bot. Gard. 5: 189 (1982)	
	Euphrasia cuspidata Hook.f. = <b>Euphrasia hookeri</b>	3:532
	Euphrasia diemenica Spreng. = <b>Euphrasia collina</b> subsp. <b>diemenica</b>	3:528
	Euphrasia sp. fabula W.C.Potts & W.R.Barker MS = <b>Euphrasia Bivouac Bay (W.R.Barker 7626 et al.) W.R.Barker</b>	
e	<b>Euphrasia fragosa</b> W.R.Barker, J. Adelaide Bot. Gard. 17: 217 (1996)	
	Euphrasia gibbsiae Du Rietz subsp. aff. subglabrifolia (Du Rietz) W.R.Barker = <b>Euphrasia gibbsiae</b> subsp. <b>Mt Field (W.R.Barker 1187) Tas Herbarium</b>	
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>comberi</b> (Du Rietz) W.R.Barker, J. Adelaide Bot. Gard. 5: 122 (1982)	3:531
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>discolor</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 128 (1982)	
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>gibbsiae</b> , Svensk Bot. Tidskr. 42: 104 (1948)	3:531
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>kingii</b> (W.M.Curtis) W.R.Barker, The Endemic Flora of Tasmania 6: 477 (1978)	3:530
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>microdonta</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 126 (1982)	
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>Mt Anne (A.M.Buchanan 5109) Tas Herbarium</b>	
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>Mt Field (W.R.Barker 1187) Tas Herbarium</b>	
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>psilantherea</b> (F.Muell.) W.R.Barker, J. Adelaide Bot. Gard. 5: 119 (1982)	
e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>pulvinestris</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 131 (1982)	

e	<b>Euphrasia gibbsiae</b> Du Rietz subsp. <b>wellingtonensis</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 120 (1982)	
	Euphrasia gunnii Du Rietz = <b>Euphrasia collina</b> subsp. <b>gunnii</b>	3:528
e	<b>Euphrasia hookeri</b> Wettst., Monogr. Euphrasia: 268 (1896)	3:532
	Euphrasia kingii W.M.Curtis = <b>Euphrasia gibbsiae</b> subsp. <b>kingii</b>	3:530
	Euphrasia multicaulis Benth. sensu Hooker (1860) = <b>Euphrasia collina</b> subsp. <b>tetragona</b>	
e	<b>Euphrasia phragmostoma</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 103 (1982)	
	<b>Euphrasia scabra</b> R.Br., Prodr. Fl. Nov. Holland. 437 (1810)	3:531
e	<b>Euphrasia semipicta</b> W.R.Barker, J. Adelaide Bot. Gard. 5: 139 (1982)	
e	<b>Euphrasia striata</b> R.Br., Prodr. Fl. Nov. Holland. 436 (1810)	3:530
	<b>Glossostigma elatinoides</b> (Benth.) Benth. ex Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 189 (1853)	3:518
	Gratiola latifolia R.Br. = <b>Gratiola peruviana</b>	3:517
	<b>Gratiola nana</b> Benth., Prodr. [A. P. de Candolle] 10: 404 (1846)	3:517
	<b>Gratiola peruviana</b> L., Sp. Pl. 1: 17 (1753)	3:517
	<b>Gratiola pubescens</b> R.Br., Prodr. Fl. Nov. Holland. 435 (1810)	3:517
	<b>Gratiola pumilo</b> F.Muell., Linnaea 25: 431 (1853)	
	Hebe elliptica (G.Forst.) Pennell = <b>Veronica elliptica</b>	
i	<b>Kickxia elatine</b> (L.) Dumort. subsp. <b>elatine</b> , Fl. Belg. (Dumortier): 35 (1827)	3:514
n i *	Kickxia spuria (L.) Dumort. subsp. <b>integrifolia</b> (Brot.) R.Fern., Bot. J. Linn. Soc. 64: 74 (1971)	3:514
	Limosella aquatica L. sensu Rodway (1903) = <b>Limosella australis</b>	3:519
	<b>Limosella australis</b> R.Br., Prodr. Fl. Nov. Holland. 443 (1810)	3:519
	Limosella lineata Glück sensu Curtis (1967) = <b>Limosella australis</b>	3:519
	Linaria spuria (L.) Mill. = <b>Kickxia spuria</b>	
i	<b>Linaria vulgaris</b> Mill., Gard. Dict., ed. 8, LIN I (1768)	3:513
	<b>Mazus pumilio</b> R.Br., Prodr. Fl. Nov. Holland. 439 (1810)	3:516
	Mimulus moschatus Douglas ex Lindl. = <b>Erythranthe moschata</b>	3:516
	Mimulus repens R.Br. = <b>Thyridia repens</b>	3:515
e	<b>Ourisia integrifolia</b> R.Br., Prodr. Fl. Nov. Holland. 439 (1810)	3:520
	Parahebe derwentiana (Andrews) B.G.Briggs & Ehrend. = <b>Veronica derwentiana</b>	3:523
i	<b>Parentucellia latifolia</b> (L.) Caruel, Fl. Ital. (Parlatore) 6: 480 (1885)	3:532
i	<b>Parentucellia viscosa</b> (L.) Caruel, Fl. Ital. (Parlatore) 6: 482 (1885)	3:533
	Pygmea ciliolata Hook.f. = <b>Veronica ciliolata</b> subsp. <b>fiordensis</b>	
	Scrophularia aquatica L. sensu Curtis (1967) = <b>Scrophularia auriculata</b>	3:515
i	<b>Scrophularia auriculata</b> L., Sp. Pl. 2: 620 (1753)	3:515
	<b>Thyridia repens</b> (R.Br.) W.R.Barker & Beardsley, Phytoneuron 39: 20 (2012)	
i	<b>Verbascum blattaria</b> L., Sp. Pl. 1: 178 (1753)	3:512
i	<b>Verbascum creticum</b> (L.) Cav., Elench. Pl. Horti Matr. 39 (1803)	3:513
i	<b>Verbascum thapsus</b> L., Sp. Pl. 1: 177 (1753)	3:512
i	<b>Verbascum virgatum</b> Stokes, Bot. Arr. Brit. Pl., ed. 2: 227 (1787)	3:512
	Veronica arguta R.Br. sensu Hooker (1860) – a name of uncertain application	
i	<b>Veronica arvensis</b> L., Sp. Pl. 1: 13 (1753)	3:525
	<b>Veronica calycina</b> R.Br., Prodr. Fl. Nov. Holland. 435 (1810)	3:524
t	<b>Veronica ciliolata</b> (Hook.f.) Cheeseman subsp. <b>fiordensis</b> (Ashwin) Meudt, Austral. Syst. Bot. 21: 413 (2008)	

e	<b>Veronica continua</b> B.G.Briggs, <i>Telopea</i> 11: 278 (2006)	
	<b>Veronica derwentiana</b> Andrews subsp. <b>derwentiana</b> , <i>Bot. Repos</i> 8: t. 531 (1808)	3:523
	<i>Veronica distans</i> R.Br. var. <i>pubescens</i> Benth. = <b>Veronica novae-hollandiae</b>	3:523
i t	<b>Veronica elliptica</b> G.Forst., <i>Fl. Ins. Austr.</i> 3 (1786)	
e	<b>Veronica formosa</b> R.Br., <i>Prodr. Fl. Nov. Holland.</i> 434 (1810)	3:522
	<b>Veronica gracilis</b> R.Br., <i>Prodr. Fl. Nov. Holland.</i> 435 (1810)	3:523
i	<b>Veronica hederifolia</b> L., <i>Sp. Pl.</i> 1: 13 (1753)	3:525
	<i>Veronica labiata</i> R.Br. nom. illeg. = <b>Veronica derwentiana</b>	
	<b>Veronica nivea</b> Lindl., <i>Edwards's Bot. Reg.</i> 28: 42 (1842)	3:523
?i # x	<b>Veronica notabilis</b> F.Muell. ex Benth., <i>Fl. Austral.</i> 4: 511 (1868)	3:524
e	<b>Veronica novae-hollandiae</b> Poir., <i>Encycl. (Lamarck)</i> 8: 526 (1808)	3:523
n	<i>Veronica peregrina</i> L. – previously listed as naturalised but insufficient evidence exists to support this	3:525
i	<b>Veronica persica</b> Poir., <i>Encycl. (Lamarck)</i> 8: 542 (1808)	3:526
	<b>Veronica plebeia</b> R.Br., <i>Prodr. Fl. Nov. Holland.</i> 435 (1810)	3:524
i t	<b>Veronica scutellata</b> L., <i>Sp. Pl.</i> 1: 12 (1753)	
i	<b>Veronica serpyllifolia</b> L., <i>Sp. Pl.</i> 1: 12 (1753)	3:525
<b>SOLANACEAE</b>		FTO 112
	<i>Anthocercis tasmanica</i> (Miers) Hook.f. = <b>Cyphanthera tasmanica</b>	3:509
e	<b>Cyphanthera tasmanica</b> Miers, <i>Ann. Mag. Nat. Hist., Ser. 2</i> , 11: 377 (1853)	3:509
i	<b>Datura ferox</b> L. d, <i>Amoen. Acad., Linnaeus ed.</i> 3: 403 (1756)	3:508
i	<b>Datura stramonium</b> L. d, <i>Sp. Pl.</i> 1: 179 (1753)	3:508
n	<i>Hyoscyamus albus</i> L. – previously listed as naturalised but insufficient evidence exists to support this	3:509
	<i>Hyoscyamus niger</i> L. sensu Curtis (1967) = <i>Hyoscyamus albus</i>	3:509
i	<b>Lycium barbarum</b> L., <i>Sp. Pl.</i> 1: 192 (1753)	
i	<b>Lycium ferocissimum</b> Miers, <i>Ann. Mag. Nat. Hist., Ser.2</i> , 14: 187 (1854)	3:507
i t	<b>Nicotiana rustica</b> L., <i>Sp. Pl.</i> 1: 180 (1753)	
n	<i>Nicotiana sylvestris</i> Speg. & Comes – previously listed as naturalised but insufficient evidence exists to support this	
n i	<b>Physalis peruviana</b> L., <i>Sp. Pl., ed. 2</i> , 2: 1670 (1763)	
i	<b>Salpichroa organifolia</b> (Lam.) Thell., <i>Mem. Soc. Sci. Nat. Math. Cherbourg</i> 38: 452 (1912)	3:506
	<i>Solanum americanum</i> Mill. = <b>Solanum nodiflorum</b> (misapplied in Tasmania)	
	<i>Solanum aviculare</i> G.Forst. sensu Bentham (1868) = <b>Solanum laciniatum</b> (misapplied in Tasmania)	
i t	<b>Solanum dulcamara</b> L., <i>Sp. Pl.</i> 1: 185 (1753)	3:504
i	<b>Solanum furcatum</b> Dunal ex Poir., <i>Encycl. (Lamarck) Suppl.</i> 3: 750 (1814)	3:504
	<b>Solanum laciniatum</b> Aiton, <i>Hortus Kew. (W.Aiton)</i> 1: 247 (1789)	3:505
i	<b>Solanum marginatum</b> L.f., <i>Suppl. Pl.</i> 147 (1782)	3:506
i	<b>Solanum mauritianum</b> Scop., <i>Delic. Fl. Faun. Insubr.</i> 3: 16, t. 8 (1788)	
i	<b>Solanum nigrum</b> L., <i>Sp. Pl.</i> 1: 186 (1753)	3:504
	<i>Solanum nitidibaccatum</i> Bitter = <b>Solanum physalifolium</b> var. <b>nitidibaccatum</b>	3:505
n i *	<i>Solanum nodiflorum</i> Jacq., <i>Collectanea [Jacquin]</i> 2: 288 (1789)	
	<b>Solanum opacum</b> A.Braun & Bouché, <i>Index Seminum Hort. Berol.</i> 8: 18 no.38 (1853)	

i	<b>Solanum physalifolium</b> Rusby var. <b>nitidibaccatum</b> (Bitter) Edmonds, Bot. J. Linn. Soc. 92: 27 (1986)	3:505
i	<b>Solanum pseudocapsicum</b> L., Sp. Pl. 1: 184 (1753) Solanum sarrachoides Sendtn. = <b>Solanum physalifolium</b> var. <b>nitidibaccatum</b>	3:506
n i	<b>Solanum triflorum</b> Nutt., Gen. N. Amer. Pl. [Nuttall] 1: 128 (1818)	
	<b>Solanum vescum</b> F.Muell., Trans. & Proc. Victorian Inst. Advancem. Sci. 1: 69 (1855) Solanum vescum F.Muell. var. <i>dauidii</i> Geras. (a cultivar name.)	3:506
<b>STACKHOUSIACEAE</b>		
	Stackhousia flava Hook. = <b>Stackhousia viminea</b>	1:112
	Stackhousia gunnii Hook.f. sensu Curtis (1956), Curtis and Morris (1975) = <b>Stackhousia subterranea</b>	1:112
	Stackhousia lineariifolia A.Cunn. = <b>Stackhousia monogyna</b>	
	<b>Stackhousia monogyna</b> Labill., Nov. Holl. Pl. 1: 77 t.104 (1805)	1:112
	<b>Stackhousia pulvinaris</b> F.Muell., Trans. Philos. Soc. Victoria 1: 101 (1855)	1:112
	<b>Stackhousia spathulata</b> Sieber ex Spreng., Syst. Veg. (ed. 16) [Sprengel] 4: 124 (1827)	1:112
	<b>Stackhousia subterranea</b> W.R.Barker, J. Adelaide Bot. Gard. 21: 90 (2007)	1:112
	<b>Stackhousia viminea</b> Sm., Cycl. (Rees): 33 (1816)	1:112
<b>STERCULIACEAE</b>		
	<b>Lasiopetalum baueri</b> Steetz, Pl. Preiss. [J.G.C.Lehman] 2: 339 (1848)	1:90
	Lasiopetalum dasyphyllum Sieber ex Steetz = <b>Lasiopetalum macrophyllum</b>	1:89
	<b>Lasiopetalum discolor</b> Hook., Companion Bot. Mag. 1: 276 (1836)	1:89
	<b>Lasiopetalum macrophyllum</b> Graham, Edinburg New Philos. J. 31: 390 (1841)	1:89
e	<b>Lasiopetalum micranthum</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 51 (1855)	1:90
<b>STYLIDIACEAE</b> FTO 127		
	Candollea armeria Labill. = <b>Stylidium armeria</b>	
	Candollea serrulata Labill. = <b>Stylidium graminifolium</b>	
e	<b>Forstera bellidifolia</b> Hook., Icon. Pl. 9: t.851 (1851)	2:395
x	<b>Levenhookia dubia</b> Sond., Pl. Preiss. [J.G.C.Lehman] 1: 392 (1845)	2:395
t	<b>Phyllachne colensoi</b> (Hook.f.) Berggr., Minneskr. Kongl. Fysiogr. Sällsk. Lund 8: 11, t.3 figs.1-27 (1877)	2:396
	<b>Stylidium armeria</b> (Labill.) Labill. subsp. <i>armeria</i> , Nov. Holl. Pl. 2: 66 (1806)	
	<b>Stylidium beaugleholei</b> J.H.Willis, Muellera 1: 153 (1967)	2:394
	Stylidium brachyphyllum Sond. sensu Curtis (1963) = <b>Stylidium beaugleholei</b>	2:394
	<b>Stylidium despectum</b> R.Br., Prodr. Fl. Nov. Holland. 571 (1810)	2:394
	Stylidium dilatatum W.D.Jacks. & R.J.E.Wiltshire = <b>Stylidium armeria</b> subsp. <i>armeria</i>	
	<b>Stylidium graminifolium</b> Sw., Sp. Pl., ed. 4 [Willdenow], 4: 146 (1805)	2:393
	Stylidium inundatum R.Br. sensu Buchanan (2005) = <b>Stylidium beaugleholei</b>	2:394
	Stylidium melastachys R.Br. = <b>Stylidium armeria</b>	
	<b>Stylidium perpusillum</b> Hook.f., London J. Bot. 6: 266 (1847)	2:394
	Stylidium umbellatum (Labill.) Labill. sensu Brown (1810) = <b>Stylidium armeria</b> – a name of uncertain application	
<b>THYMELAEACEAE</b> FTO 89		
i t	<b>Daphne laureola</b> L., Sp. Pl. 1: 356 (1753)	
	Drapetes tasmanicus Hook.f. = <b>Kelleria dieffenbachii</b>	3:625



	<b>Kelleria dieffenbachii</b> (Hook.) Endl., Gen. Pl. [Endlicher] Suppl. 4(2): 61 (1848)	3:625
	<b>Pimelea axiflora</b> F.Muell. ex Meisn. subsp. <b>axiflora</b> , Linnaea 26: 345 (1854)	3:622
	<i>Pimelea cernua</i> R.Br. = <b>Pimelea linifolia</b>	
e	<b>Pimelea cinerea</b> R.Br., Prodr. Fl. Nov. Holland. 361 (1810)	3:619
	<b>Pimelea curviflora</b> R.Br., Prodr. Fl. Nov. Holland. 362 (1810)	3:624
	<i>Pimelea curviflora</i> R.Br. var. <i>gracilis</i> (R.Br.) Threlfall = <b>Pimelea curviflora</b>	3:624
	<i>Pimelea curviflora</i> R.Br. var. <i>sericea</i> Benth. = <b>Pimelea curviflora</b>	3:624
	<b>Pimelea drupacea</b> Labill., Nov. Holl. Pl. 1: 10 (1805)	3:622
e	<b>Pimelea filiformis</b> Hook.f., London J. Bot. 6: 280 (1847)	3:622
	<b>Pimelea flava</b> R.Br. subsp. <b>flava</b> , Prodr. Fl. Nov. Holland. 361 (1810)	3:623
n	<i>Pimelea</i> sp. Freycinet (A.M.Buchanan 15902) Tas Herbarium = <b>Pimelea leiophylla</b>	
	<b>Pimelea glauca</b> R.Br., Prodr. Fl. Nov. Holland. 360 (1810)	3:619
	<i>Pimelea gracilis</i> R.Br. sensu Hooker (1860) = <b>Pimelea curviflora</b>	
	<i>Pimelea gunnii</i> Hook.f. = <b>Pimelea cinerea</b>	
	<b>Pimelea humilis</b> R.Br., Prodr. Fl. Nov. Holland. 361 (1810)	3:620
	<i>Pimelea incana</i> R.Br. = <b>Pimelea nivea</b>	
n e	<b>Pimelea leiophylla</b> A.M.Gray & M.Baker, Muelleria 35: 16 (2016)	
	<b>Pimelea ligustrina</b> Labill. subsp. <b>ligustrina</b> , Nov. Holl. Pl. 1: 9, f.3 (1805)	
	<i>Pimelea lindleyana</i> Meisn. = <b>Pimelea linifolia</b>	3:620
	<b>Pimelea linifolia</b> Sm., Spec. Bot. New Holland 31, t.11 (1793)	3:620
	<i>Pimelea linifolia</i> Sm. subsp. <i>linifolia</i> = <b>Pimelea linifolia</b>	
	<i>Pimelea linifolia</i> Sm. subsp. <i>linoides</i> (A.Cunn.) Threlfall = <b>Pimelea linifolia</b>	
	<b>Pimelea micrantha</b> F.Muell. ex Meisn., Linnaea 26: 351 (1854)	3:624
e	<b>Pimelea milliganii</b> Meisn., Prodr. [A. P. de Candolle] 14: 509 (1857)	3:619
e	<b>Pimelea nivea</b> Labill., Nov. Holl. Pl. 1: 10 (1805)	3:621
	<b>Pimelea pauciflora</b> R.Br., Prodr. Fl. Nov. Holland. 360 (1810)	3:622
	<i>Pimelea phycoides</i> Meisn. sensu Rye (1990) probably recorded in error	
e	<b>Pimelea pygmaea</b> F.Muell. & C.Stuart ex Meisn., Linnaea 26: 346 (1853)	3:623
e	<b>Pimelea sericea</b> R.Br., Prodr. Fl. Nov. Holland. 361 (1810)	3:621
	<b>Pimelea serpyllifolia</b> R.Br. subsp. <b>serpyllifolia</b> , Prodr. Fl. Nov. Holland. 360 (1810)	3:623
	<i>Pimelea spatulata</i> Labill. = <b>Pimelea linifolia</b>	
	<i>Pimelea stricta</i> Meisn. sensu Curtis (1967) = <b>Pimelea linifolia</b>	3:624
	<b>Pimelea sp. Tunbridge (A.Moscal 9026) Tas Herbarium</b>	
<b>TREMANDRACEAE</b>		FTO 71
	<b>Tetratheca ciliata</b> Lindl., Three Exped. Australia [Mitchell] 2: 205 (1838)	1:60
e	<b>Tetratheca sp. Flinders Is. (T.Rudman HO510551) Tas Herbarium</b>	
e	<b>Tetratheca sp. Freycinet Pen. (A.C.Rozefelds 323) Tas Herbarium</b>	
	<i>Tetratheca glandulosa</i> Labill. = <b>Tetratheca labillardierei</b>	1:60
e	<b>Tetratheca gunnii</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.). 1: 36, t.7B (1855)	
	<b>Tetratheca labillardierei</b> Joy Thomps., Telopea 1: 189 (1976)	1:60
	<b>Tetratheca pilosa</b> Labill. subsp. <b>latifolia</b> Joy Thomps., Telopea 1: 213 (1976)	
	<b>Tetratheca pilosa</b> Labill. subsp. <b>pilosa</b> , Nov. Holl. Pl. 1: 95, t.122 (1805)	1:60
	<i>Tetratheca pilosa</i> Labill. var. <i>procumbens</i> = <b>Tetratheca procumbens</b>	
	<b>Tetratheca procumbens</b> Gunn ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 35, t.7A (1855)	1:61

	<b>TROPAEOLACEAE</b>	FTO 82
i	<b>Tropaeolum majus</b> L., Sp. Pl. 1: 345 (1753)	
	<b>ULMACEAE</b>	FTO 63
i	<b>Ulmus × hollandica</b> Mill., Gard. Dict., ed. 8, 5 (1768)	3:642
	<b>URTICACEAE</b>	FTO 64
	Australina muelleri Wedd. = <b>Australina pusilla</b> subsp. <b>muelleri</b>	3:641
	<b>Australina pusilla</b> (Poir.) Gaudich. subsp. <b>muelleri</b> (Wedd.) Friis & Wilmot-Dear, Nordic. J. Bot. 7: 126 (1987)	3:641
	<b>Australina pusilla</b> (Poir.) Gaudich. subsp. <b>pusilla</b> , Voy. Bonite, Bot. 3: t. 114A (1852)	3:641
	Helxine soleirolii Req. sensu Curtis (1967) = <b>Soleirolia soleirolii</b>	3:640
	<b>Parietaria cardiostegia</b> Greuter, Fl. Australia 3: 190 (1989)	
	<b>Parietaria debilis</b> G.Forst., Fl. Ins. Austr. 73 (1786)	3:640
n i *	<b>Parietaria judaica</b> L., Sp. Pl., ed. 2, 2: 1492 (1763)	
i	<b>Soleirolia soleirolii</b> (Req.) Dandy, Feddes Repert. 70: 4 (1965)	3:640
	<b>Urtica incisa</b> Poir., Encycl. (Lamarck) Suppl. 4: 224 (1816)	3:639
i	<b>Urtica urens</b> L., Sp. Pl. 2: 984 (1753)	3:639
	<b>VALERIANACEAE</b>	FTO 135
i	<b>Centranthus ruber</b> (L.) DC., Fl. Franc. (DC. & Lamarck), ed. 3, 4(2): 239 (1805)	2:278
i	<b>Valerianella eriocarpa</b> Desv., J. Bot. (Desvaux) 2: 314, t. 11 fig. 2 (1809)	2:278
i	<b>Valerianella locusta</b> (L.) Laterr., Fl. Bordel., ed. 2, 93 (1821)	
i t	<b>Valerianella rimosa</b> Bastard, J. Bot. Agric. 3: 20 (1814)	
	<b>VERBENACEAE</b>	
i	<b>Verbena bonariensis</b> L., Sp. Pl. 1: 20 (1753)	
	Verbena incompta P.W.Michael = <b>Verbena bonariensis</b>	
i	<b>Verbena officinalis</b> L., Sp. Pl. 1: 20 (1753)	3:541
	<b>VIOLACEAE</b>	FTO 79
	Hymenanchera angustifolia R.Br. = <b>Melicytus angustifolius</b>	
	Hymenanchera banksii F.Muell. nom. illeg., nom. superfl. = <b>Melicytus dentatus</b>	
	Hymenanchera dentata R.Br. ex DC. sensu Curtis & Morris (1975) = <b>Melicytus dentatus</b> and <b>M. angustifolius</b>	1:55
e	<b>Melicytus angustifolius</b> (DC.) Garn.-Jones subsp. <b>angustifolius</b> , New Zealand J. Bot. 25: 127 (1987)	
	<b>Melicytus angustifolius</b> (DC.) Garn.-Jones subsp. <b>divaricatus</b> Stajsic & R.Douglas, Austral. Syst. Bot. 27: 320 (2014)	
	<b>Melicytus dentatus</b> (R.Br. ex DC.) Molloy & Mabb., Curtis's Bot. Mag. 17: 234 (2000)	1:55
i	<b>Viola arvensis</b> Murray, Prodr. Stirp. Gott. 73 (1770)	1:55
	<b>Viola betonicifolia</b> Sm. subsp. <b>betonicifolia</b> , Cycl. (Rees) 37: no.7 (1817)	1:54
	<b>Viola calejana</b> G.Don, Gen. Hist. 1: 329 (1831)	1:55
	<b>Viola cleistogamoides</b> (L.G.Adams) Seppelt, Fl. S. Australia, ed. 4, 2: 870 (1986)	
t	<b>Viola cunninghamii</b> Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 16 (1852)	1:54
	<b>Viola fuscoviolacea</b> (L.G.Adams) T.A.James, Muelleria 9: 35 (1996)	
	Viola hederacea Labill. subsp. <b>cleistogamoides</b> L.G.Adams = <b>Viola cleistogamoides</b>	
e	<b>Viola hederacea</b> Labill. subsp. <b>curtisiae</b> L.G.Adams, Fl. Australia 8: 386 (1982)	
	Viola hederacea Labill. subsp. <b>fuscoviolacea</b> L.G.Adams = <b>Viola fuscoviolacea</b>	
	<b>Viola hederacea</b> Labill. subsp. <b>hederacea</b> , Nov. Holl. Pl. 1: 66, t.91 (1805)	1:54

i	<b>Viola odorata</b> L., Sp. Pl. 2: 934 (1753)	
	<b>Viola sieberiana</b> Spreng., Syst. Veg. (ed. 16) [Sprengel] 4(2): 96 (1827)	1:54
<b>WINTERACEAE</b>		FTO 2
	<i>Drimys aromatica</i> (R.Br. ex DC.) F.Muell. = <b>Tasmannia lanceolata</b>	
	<i>Drimys lanceolata</i> (Poir.) Baill. = <b>Tasmannia lanceolata</b>	1:25
	<i>Tasmannia aromatica</i> R.Br. ex DC. = <b>Tasmannia lanceolata</b>	
	<b>Tasmannia lanceolata</b> (Poir.) A.C.Sm., Taxon 18: 287 (1969)	1:25
<b>ZYGOPHYLLACEAE</b>		FTO 58
	<i>Roepera billardierei</i> A.Juss. sensu Hooker (1860) = <b>Zygophyllum billardierei</b>	
	<i>Roepera latifolia</i> Hook.f. sensu Hooker (1860) recorded in error	
	<i>Zygophyllum apiculatum</i> F.Muell. sensu Curtis (1975) recorded in error	1:93
	<b>Zygophyllum billardierei</b> DC., Prodr. [A. P. de Candolle] 1: 705 (1824)	1:93

## MONOCOTYLEDONEAE

### AGAVACEAE

i	<b>Agave americana</b> L., Sp. Pl. 1: 323 (1753)	
i	<b>Cordyline australis</b> (G.Forst.) Endl., Prodr. Fl. Norfolk. 29 (1833)	
i	<b>Phormium tenax</b> J.R.Forst. & G.Forst., Char. Gen. Pl. 48, t.24 (1775)	4b:418

### ALISMATACEAE

i	<b>Alisma lanceolatum</b> With., Arr. Brit. Pl., ed. 3, 2: 362 (1796)	
i	<b>Alisma plantago-aquatica</b> L., Sp. Pl. 1: 342 (1753)	4b:4
	<i>Damasonium australe</i> Salisb. = <b>Damasonium minus</b>	4b:6
	<b>Damasonium minus</b> (R.Br.) Buchenau, Abh. Naturwiss. Vereins Bremen 2: 20 (1869)	4b:6

### ALOEACEAE

i	<b>Aloe maculata</b> All., Auct. Syn. Meth. Stirp. Hort. Regii Taur. 13 (1773)	
	<i>Aloe saponaria</i> (Aiton) Haw. = <b>Aloe maculata</b>	
n i	<b>Kniphofia uvaria</b> (L.) Oken, Allg. Naturgesch. 3: 566 (1841)	

### APONOGETONACEAE

i	<b>Aponogeton distachyos</b> L.f., Suppl. Pl. 215 (1782)	4b:13
---	--	-------

### ARACEAE

i	<b>Zantedeschia aethiopica</b> (L.) Spreng., Sys. Veg. (ed. 16) [Sprengel] 3: 765 (1826)	4b:30
---	--	-------

### BURMANNIACEAE

	<b>Thismia rodwayi</b> F.Muell., Vict. Naturalist 7: 115 (1890)	4b:423
--	---	--------

### CENTROLEPIDACEAE

	<i>Alepyrum monogynum</i> Hook.f. = <b>Centrolepis monogyna</b>	
	<i>Alepyrum muelleri</i> Hook.f. = <b>Centrolepis glabra</b>	
	<i>Alepyrum muscoides</i> Hook.f. = <b>Centrolepis muscoides</b>	
	<i>Alepyrum polygynum</i> R.Br. sensu Hooker (1860) = <b>Centrolepis polygyna</b>	
	<b>Aphelia gracilis</b> Sond., Linnaea 28: 227 (1856)	4b:49
	<i>Aphelia gunnii</i> Hook.f. = <b>Aphelia gracilis</b>	

	<b>Aphelia pumilio</b> F.Muell. ex Sond., Linnaea 28: 226 (1856)	4b:49
	<b>Centrolepis aristata</b> (R.Br.) Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 1: 44 (1817)	4b:54
	<b>Centrolepis fascicularis</b> Labill., Nov. Holl. Pl. 1: 7, t.1 (1804)	4b:55
	<b>Centrolepis glabra</b> (F.Muell. ex Sond.) Hieron., Abh. Naturf. Ges. Halle 12: 209 (1873)	4b:52
e	<b>Centrolepis monogyna</b> (Hook.f.) Benth., Fl. Austral. 7: 205 (1878)	4b:53
	Centrolepis monogyna (Hook.f.) Benth. subsp. paludicola (W.M.Curtis) D.A.Cooke =	
	<b>Centrolepis monogyna</b>	4b:53
e	<b>Centrolepis muscoides</b> (Hook.f.) Hieron., Abh. Naturf. Ges. Halle 12: 209 (1873)	4b:53
	Centrolepis paludicola W.M.Curtis = <b>Centrolepis monogyna</b>	4b:53
e	<b>Centrolepis pedderensis</b> W.M.Curtis, Brunonia 7: 299 (1984)	4b:53
	<b>Centrolepis polygyna</b> (R.Br.) Hieron., Abh. Naturf. Ges. Halle 12: 210 (1873)	4b:51
	Centrolepis pulvinata (R.Br.) Roem. & Schult. = <b>Centrolepis strigosa</b> subsp. <b>pulvinata</b>	4b:55
e	<b>Centrolepis strigosa</b> (R.Br.) Roem. & Schult. subsp. <b>pulvinata</b> (R.Br.) D.A.Cooke, J. Adelaide Bot. Gard. 15: 28 (1992)	4b:55
	<b>Centrolepis strigosa</b> (R.Br.) Roem. & Schult. subsp. <b>strigosa</b> , Syst. Veg., ed. 15 bis [Roemer & Schultes] 1: 43 (1817)	4b:55
	Centrolepis tenuior (R.Br.) Roem. & Schult. sensu Hooker (1860) = <b>Centrolepis strigosa</b> subsp. <b>strigosa</b>	
	Devauxia billardierei R.Br. = <b>Centrolepis fascicularis</b>	
	Devauxia pulvinata R.Br. = <b>Centrolepis strigosa</b> subsp. <b>pulvinata</b>	
	Devauxia tenuior R.Br. = <b>Centrolepis strigosa</b> subsp. <b>strigosa</b>	
e	<b>Gaimardia amblyphylla</b> W.M.Curtis, Brunonia 7: 299 (1984)	4b:57
e	<b>Gaimardia fitzgeraldii</b> F.Muell. & Rodway, Pap. & Proc. Roy. Soc. Tasmania 1894-5: 55 (1896)	4b:57
t	<b>Gaimardia setacea</b> Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 267 (1853)	4b:57
<b>COMMELINACEAE</b>		
	Tradescantia albiflora Kunth = <b>Tradescantia fluminensis</b>	4b:36
i	<b>Tradescantia fluminensis</b> Vell., Fl. Flumin. 140 (1829)	4b:36
<b>CYMODOCEACEAE</b>		
	<b>Amphibolis antarctica</b> (Labill.) Sond. & Asch. ex Asch., Linnaea 35: 164 (1867)	4b:27
	Cymodocea antarctica (Labill.) Endl. ex Kunth = <b>Amphibolis antarctica</b>	4b:27
<b>CYPERACEAE</b>		
	<b>Baumea acuta</b> (Labill.) Palla, Allg. Bot. Z. Syst. 8: 69 (1902)	4b:131
	<b>Baumea arthropphylla</b> (Nees) Boeckeler, Linnaea 38: 242 (1874)	4b:129
	<b>Baumea articulata</b> (R.Br.) S.T.Blake, Contr. Queensland Herb. 8: 28 (1969)	4b:128
	<b>Baumea gunnii</b> (Hook.f.) S.T.Blake, Contr. Queensland Herb. 8: 27 (1969)	4b:130
	<b>Baumea juncea</b> (R.Br.) Palla, Allg. Bot. Z. Syst. 15: 113 (1909)	4b:129
?i	<b>Baumea planifolia</b> (Benth.) K.L.Wilson, Telopea 5: 589 (1994)	
	<b>Baumea rubiginosa</b> (Spreng.) Boeckeler, Linnaea 38: 241 (1874)	4b:129
	<b>Baumea tetragona</b> (Labill.) S.T.Blake, Contr. Queensland Herb. 8: 30 (1969)	4b:131
	<b>Bolboschoenus caldwellii</b> (V.J.Cook) Soják, Cas. Nár. Mus., Odd. Prír. 141: 62 (1972)	4b:92
	<b>Bolboschoenus medianus</b> (V.J.Cook) Soják, Cas. Nár. Mus., Odd. Prír. 141: 63 (1972)	4b:93
	Carex acicularis Boott sensu Rodway (1903) = <b>Carex archeri</b>	4b:148
	Carex albula Allan sensu Curtis & Morris (1994) recorded in error	4b:164

e	<b>Carex sp. Algonkian Rivulet (S.J.Jarman HOI 10282) Tas Herbarium</b>	4b:163
	<b>Carex appressa</b> R.Br., Prodr. Fl. Nov. Holland. 242 (1810)	4b:152
	<i>Carex appressa</i> R.Br. f. minor Kük. = <b>Carex appressa</b>	4b:152
	<i>Carex appressa</i> R.Br. var. <i>virgata</i> (Boott) Kük. = <b>Carex appressa</b>	4b:153
	<b>Carex archeri</b> Boott, Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 98 (1858)	4b:148
e	<b>Carex barbata</b> Boott, Ill. Gen. Carex 1: 68, t.187 (1858)	4b:160
	<b>Carex bichenoviana</b> Boott, Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 101 (1858)	4b:167
	<b>Carex breviculmis</b> R.Br., Prodr. Fl. Nov. Holland. 242 (1810)	4b:158
n i *	<i>Carex buxbaumii</i> Wahlenb., Kongl. Vetensk. Acad. Nya Handl. 24: 163 (1803)	
	<i>Carex caespitosa</i> L. sensu Brown (1810) = <b>Carex gaudichaudiana</b>	
	<i>Carex canescens</i> L. = <b>Carex curta</b>	4b:155
	<b>Carex capillacea</b> Boott, Ill. Gen. Carex 1: 44, t.110 (1858)	4b:148
e	<b>Carex cataractae</b> R.Br., Prodr. Fl. Nov. Holland. 242 (1810)	4b:162
	<b>Carex cephalotes</b> F.Muell., Trans. Philos. Soc. Victoria 1: 110 (1855)	4b:149
	<b>Carex chlorantha</b> R.Br., Prodr. Fl. Nov. Holland. 242 (1810)	4b:149
	<i>Carex</i> aff. <i>cirrhosa</i> Berggr. sensu Curtis & Morris (1994) = <b>Carex sp. Algonkian Rivulet (S.J.Jarman HOI 10282) Tas Herbarium</b>	4b:163
	<b>Carex curta</b> Gooden., Trans. Linn. Soc. London 2: 145 (1794)	4b:155
i t	<b>Carex demissa</b> Hornem., Fl. Dan., 7(21): 4, t.1342 (1806)	4b:161
	<i>Carex</i> aff. <i>diandra</i> Schrank sensu Curtis & Morris (1994) = <b>Carex sp. Western Tasmania (A.Moscal 7489) Tas Herbarium</b>	4b:150
i	<b>Carex divisa</b> Huds., Fl. Angl. (Hudson): 348 (1762)	4b:149
i	<b>Carex divulsa</b> Stokes, Bot. Arr. Brit. Pl., ed. 2. 2: 1035 (1787)	4b:150
	<b>Carex fascicularis</b> Sol. ex Boott, Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 283 (1853)	4b:166
i	<b>Carex flacca</b> Schreb., Spicil. Fl. Lips. 178 (1771)	4b:157
i t	<b>Carex flagellifera</b> Colenso, Trans. & Proc. New Zealand Inst. 16: 342 (1884)	4b:163
	<i>Carex flava</i> L. sensu Rodway (1903) = <b>Carex cataractae</b> , <b>C. demissa</b> , <b>C. flaviformis</b> & <b>C. lepidocarpa</b>	
t	<b>Carex flaviformis</b> Nelves, Kew Bull. 10: 84 (1955)	4b:162
	<b>Carex gaudichaudiana</b> Kunth, Enum. Pl. [Kunth] 2: 417 (1837)	4b:156
	<b>Carex gunniana</b> Boott, Proc. Linn. Soc. London 1: 258 (1845)	4b:160
	<b>Carex hypandra</b> F.Muell. ex Benth., Fl. Austral. 7: 439 (1878)	4b:157
	<b>Carex inversa</b> R.Br., Prodr. Fl. Nov. Holland. 242 (1810)	4b:153
	<b>Carex iynx</b> Nelves, Proc. Linn. Soc. London 155: 279 (1944)	4b:159
i t	<b>Carex lepidocarpa</b> Tausch, Flora 17: 179 (1834)	4b:161
i	<b>Carex leporina</b> L., Sp. Pl. 2: 973 (1753)	4b:155
	<i>Carex littorea</i> Labill. = <b>Carex pumila</b>	
	<b>Carex longebrachiata</b> Boeckeler, Linnaea 41: 282 (1877)	4b:158
	<i>Carex longifolia</i> R.Br. sensu Rodway (1903) = <b>Carex iynx</b> & <b>C. longebrachiata</b>	
	<i>Carex ovalis</i> Gooden. = <b>Carex leporina</b>	
	<i>Carex paniculata</i> L. sensu Rodway (1903) = <b>Carex appressa</b>	4b:152
n i * t	<i>Carex pilulifera</i> L., Sp. Pl. 2: 976 (1753)	
	<b>Carex polyantha</b> F.Muell., Trans. Philos. Soc. Victoria 1: 110 (1855)	4b:156
	<i>Carex pseudocyperus</i> L. sensu Rodway (1903) = <b>Carex fascicularis</b>	4b:166
	<b>Carex pumila</b> Thunb. ex Murray, Syst. Veg., ed. 14 (J. A. Murray). 846 (1784)	4b:166

	<b>Carex raleighii</b> Nelves, Bull. Misc. Inform. Kew 1939: 310 (1939)	4b:154
n i *	<i>Carex scoparia</i> Schkuhr ex Willd., Sp. Pl., ed. 4 [Willdenow], 4: 230 (1805)	4b:154
	<b>Carex tasmanica</b> Kük., Bull. Herb. Boissier 4: 59 (1904)	4b:165
	<b>Carex tereticaulis</b> F.Muell., Fragm. (Mueller) 8: 256 (1874)	4b:151
n	<i>Carex testacea</i> Sol. ex Boott – previously listed as naturalised but insufficient evidence exists to support this	4b:164
	<i>Carex virgata</i> Sol. ex Boott = <b>Carex appressa</b>	4b:153
	<i>Carex vulgaris</i> Fr. sensu Rodway (1903) = <b>Carex gaudichaudiana</b>	4b:156
e	<b>Carex sp. Western Tasmania (A.Moscal 7489) Tas Herbarium</b>	4b:150
	<b>Carpha alpina</b> R.Br., Prodr. Fl. Nov. Holland. 230 (1810)	4b:87
e	<b>Carpha curvata</b> W.M.Curtis, Brunonia 7: 303 (1985)	4b:88
e	<b>Carpha rodwayi</b> W.M.Curtis, Brunonia 7: 302 (1985)	4b:88
	<b>Caustis pentandra</b> R.Br., Prodr. Fl. Nov. Holland. 240 (1810)	4b:137
	<i>Chaetospora axillaris</i> R.Br. sensu Hooker (1860) = <b>Schoenus maschalinus</b>	
	<i>Chaetospora capillacea</i> Hook.f. nom. illeg. = <b>Tetraria capillaris</b>	
	<i>Chaetospora imberbis</i> R.Br. sensu Hooker (1860) = <b>Schoenus apogon</b>	
	<i>Chaetospora nitens</i> R.Br. = <b>Schoenus nitens</b>	
	<i>Chaetospora tenuissima</i> Hook.f. nom. illeg. = <b>Schoenus lepidosperma</b>	
	<b>Chorizandra australis</b> K.L.Wilson, Telopea 5: 594 (1994)	4b:138
	<i>Chorizandra cymbaria</i> R.Br. sensu Rodway (1903) = <b>Chorizandra australis</b>	4b:138
	<b>Chorizandra enodis</b> Nees, Pl. Preiss. [J.G.C.Lehman] 2: 73 (1846)	4b:138
	<i>Cladium acutum</i> (Labill.) Poir. = <b>Baumea acuta</b>	
	<i>Cladium filum</i> (Labill.) R.Br. = <b>Gahnia filum</b>	4b:136
	<i>Cladium glomeratum</i> R.Br. sensu Rodway (1903) = <b>Baumea rubiginosa</b>	4b:129
	<i>Cladium gunnii</i> Hook.f. = <b>Baumea gunnii</b>	4b:130
	<i>Cladium junceum</i> R.Br. = <b>Baumea juncea</b>	4b:129
	<i>Cladium laxiflorum</i> Hook.f. = <b>Baumea gunnii</b>	
	<i>Cladium mariscus</i> (L.) R.Br. recorded in error	
n ?i #	<b>Cladium procerum</b> S.T.Blake, Trans. Roy. Soc. South Australia 67: 57 (1943)	4b:428
	<i>Cladium schoenoides</i> R.Br. = <b>Baumea acuta</b>	4b:131
	<i>Cladium tetraquetrum</i> Hook.f. = <b>Baumea tetragona</b>	4b:131
i	<b>Cyperus congestus</b> Vahl, Enum. Pl. [Vahl] 2: 358 (1805)	4b:91
i	<b>Cyperus eragrostis</b> Lam., Tabl. Encycl. 1: 146 (1791)	4b:90
	<b>Cyperus gunnii</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 80, t.140, fig.A (1858)	4b:91
	<b>Cyperus lucidus</b> R.Br., Prodr. Fl. Nov. Holland. 218 (1810)	4b:91
	<i>Cyperus sanguineo-fuscus</i> Nees sensu Hooker (1860) = <b>Cyperus lucidus</b>	
?i #	<b>Cyperus sanguinolentus</b> Vahl, Enum. Pl. [Vahl] 2: 351 (1805)	4b:90
	<i>Cyperus tenellus</i> L.f. = <b>Isolepis levynsiana</b>	4b:90
	<i>Eleocharis acicularis</i> (L.) Roem. & Schult. sensu Rodway (1903) = <b>Eleocharis pusilla</b>	4b:108
	<b>Eleocharis acuta</b> R.Br., Prodr. Fl. Nov. Holland. 224 (1810)	4b:107
	<b>Eleocharis gracilis</b> R.Br., Prodr. Fl. Nov. Holland. 224 (1810)	4b:107
	<b>Eleocharis pusilla</b> R.Br., Prodr. Fl. Nov. Holland. 225 (1810)	4b:108
	<b>Eleocharis sphacelata</b> R.Br., Prodr. Fl. Nov. Holland. 224 (1810)	4b:107
	<i>Elynanthus capillaceus</i> Benth. = <b>Tetraria capillaris</b>	4b:132
	<b>Ficinia nodosa</b> (Rottb.) Goetgh., Muasya & D.A.Simpson, Novon 10: 133 (2000)	4b:105

	<b>Gahnia filum</b> (Labill.) F.Muell., Key to the System of Victorian Plants 1: 456 (1888)	4b:136
	Gahnia fitzgeraldii Rodway = <b>Gahnia microstachya</b>	4b:134
	Gahnia graminifolia Rodway = <b>Gahnia rodwayi</b>	4b:134
	<b>Gahnia grandis</b> (Labill.) S.T.Blake, Contr. Queensland Herb. 8: 33 (1969)	4b:135
	Gahnia melanocarpa R.Br. sensu Buchanan et al. (1989) = <b>Gahnia grandis</b> (misapplied in Tasmania)	
	<b>Gahnia microstachya</b> Benth., Fl. Austral. 7: 414 (1878)	4b:134
	Gahnia psittacorum Labill. = <b>Gahnia grandis</b>	4b:135
	<b>Gahnia radula</b> (R.Br.) Benth., Fl. Austral. 7: 417 (1878)	4b:135
e	<b>Gahnia rodwayi</b> F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania 1892: 93 (1893)	4b:134
	<b>Gahnia sieberiana</b> Kunth, Enum. Pl. [Kunth] 2: 332 (1837)	4b:135
	<b>Gahnia trifida</b> Labill., Nov. Holl. Pl. 1: 89, t.116 (1805)	4b:136
	<b>Gymnoschoenus sphaerocephalus</b> (R.Br.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 83, t.142 (1858)	4b:119
	<b>Isolepis alpina</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 86, t.143, fig.B (1858)	4b:103
	<b>Isolepis aucklandica</b> Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 88 (1844), pl.50 (1845)	4b:102
	Isolepis cartilaginea R.Br. = <b>Isolepis marginata</b>	
	<b>Isolepis cernua</b> (Vahl) Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 2: 106 (1817)	4b:99
	<b>Isolepis crassiuscula</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 86, t.143, fig.A (1858)	4b:98
	<b>Isolepis fluitans</b> (L.) R.Br., Prodr. Fl. Nov. Holland. 221 (1810)	4b:98
	<b>Isolepis habra</b> (Edgar) Soják, Casopsis Narodniho Muzea v Praze 148: 194 (1980)	4b:101
	<b>Isolepis hookeriana</b> Boeckeler, Flora 41: 418 (1858)	4b:104
n i	<b>Isolepis hystrix</b> (Thunb.) Nees, Linnaea 7: 496 (1832)	
	<b>Isolepis inundata</b> R.Br., Prodr. Fl. Nov. Holland. 222 (1810)	4b:100
	Isolepis lenticularis R.Br. sensu Hooker (1860) = <b>Isolepis fluitans</b>	
?i	<b>Isolepis levynsiana</b> Muasya & D.A.Simpson, Novon 17: 59 (2007)	4b:90
e	<b>Isolepis limbata</b> W.M.Curtis, Brunonia 7: 300 (1985)	4b:103
	<b>Isolepis marginata</b> (Thunb.) A.Dietr., Sp. Pl., ed. 6. 1(2): 110 (1832)	4b:104
	<b>Isolepis montivaga</b> (S.T.Blake) K.L.Wilson, Telopea 2: 168 (1981)	4b:103
	Isolepis nodosa (Rottb.) R.Br. = <b>Ficinia nodosa</b>	4b:105
	<b>Isolepis platycarpa</b> (S.T.Blake) Soják, Cas. Nár. Muz. Praze, Rada Prír. 148: 194 (1980)	4b:99
	<b>Isolepis producta</b> (C.B.Clarke) K.L.Wilson, Telopea 2: 168 (1981)	4b:98
i	<b>Isolepis prolifera</b> (Rottb.) R.Br., Prodr. Fl. Nov. Holland. 223 (1810)	
	Isolepis riparia R.Br. sensu Hooker (1860) = <b>Isolepis cernua</b>	
	Isolepis saviana Schult. nom. illeg. sensu Hooker (1860) – a name of uncertain application	
i t	<b>Isolepis setacea</b> (L.) R.Br., Prodr. Fl. Nov. Holland. 222 (1810)	4b:105
	<b>Isolepis stellata</b> (C.B.Clarke) K.L.Wilson, Telopea 2: 169 (1981)	4b:104
	<b>Isolepis subtilissima</b> Boeckeler, Flora 41: 416 (1858)	4b:101
e	<b>Isolepis tasmanica</b> (S.T.Blake) K.L.Wilson, Telopea 2: 170 (1981)	4b:102
	<b>Isolepis wakefieldiana</b> (S.T.Blake) K.L.Wilson, Telopea 2: 170 (1981)	4b:100
	Lampocarya hexandra R.Br. = <b>Gahnia trifida</b>	
	Lepidosperma angustifolia Hook.f. sensu Hooker (1860) = <b>Lepidosperma laterale</b>	
	<b>Lepidosperma concavum</b> R.Br., Prodr. Fl. Nov. Holland. 234 (1810)	4b:126
	<b>Lepidosperma curtisiae</b> K.L.Wilson & D.I.Morris, Telopea 5: 423 (1993)	4b:123
	Lepidosperma elatior Labill. orth. var. sensu Rodway (1903) = <b>Lepidosperma elatius</b>	4b:124

	<b>Lepidosperma elatius</b> Labill., Nov. Holl. Pl. 1: 15, t.11 (1805)	4b:124
	Lepidosperma elatius Labill. var. ensiforme Rodway = <b>Lepidosperma ensiforme</b>	4b:125
	Lepidosperma elatius Labill. var. oldfieldii (Hook.f.) Rodway = <b>Lepidosperma oldfieldii</b>	4b:125
	<b>Lepidosperma ensiforme</b> (Rodway) D.I.Morris, The Student's Flora of Tasmania 4b: 425 (1994)	4b:125
	Lepidosperma falcatum Rodway = <b>Lepidosperma tortuosum</b>	
	<b>Lepidosperma filiforme</b> Labill., Nov. Holl. Pl. 1: 17, t.15 (1805)	4b:121
	<b>Lepidosperma forsythii</b> A.A.Ham., Proc. Linn. Soc. New South Wales 35: 411 (1910)	4b:122
	<b>Lepidosperma gladiatum</b> Labill., Nov. Holl. Pl. 1: 15, t.12 (1805)	4b:125
e	<b>Lepidosperma globosum</b> Labill., Nov. Holl. Pl. 1: 16, t.14 (1805)	4b:123
	<b>Lepidosperma gunnii</b> Boeckeler, Linnaea 38: 325 (1874)	4b:123
e	<b>Lepidosperma inops</b> F.Muell. ex Rodway, Pap. & Proc. Roy. Soc. Tasmania 1892: 93 (1893)	4b:122
	<b>Lepidosperma laterale</b> R.Br., Prodr. Fl. Nov. Holland. 234 (1810)	4b:124
	Lepidosperma lineare R.Br. sensu Rodway (1903) = <b>Lepidosperma gunnii</b>	4b:123
	Lepidosperma lineare R.Br. var. inops (F.Muell. ex Rodway) Rodway = <b>Lepidosperma inops</b>	4b:122
	<b>Lepidosperma longitudinale</b> Labill., Nov. Holl. Pl. 1: 16, t.13 (1805)	4b:124
e	<b>Lepidosperma monticola</b> G.T.Plunkett & J.J.Bruhl, Phytokeys 28: 26 (2013)	
	<b>Lepidosperma neesii</b> Kunth, Enum. Pl. [Kunth] 2: 319 (1837)	4b:127
e	<b>Lepidosperma oldfieldii</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 91, t.146A (1858)	4b:125
	Lepidosperma squamata Labill. sensu Rodway (1903) = <b>Lepidosperma concavum</b>	4b:126
	Lepidosperma tetragonum Labill. sensu Brown (1810) = <b>Baumea tetragona</b>	
	<b>Lepidosperma tortuosum</b> F.Muell., Fragm. (Mueller) 9: 23 (1875)	4b:121
	<b>Lepidosperma viscidum</b> R.Br., Prodr. Fl. Nov. Holland. 234 (1810)	4b:126
	Mesomelaena sphaerocephala (R.Br.) Benth. = <b>Gymnoschoenus sphaerocephalus</b>	4b:119
e	<b>Oreobolus acutifolius</b> S.T.Blake, Contr. Queensland Herb. 8: 39 (1969)	4b:109
	<b>Oreobolus distichus</b> F.Muell., Trans. Philos. Soc. Victoria 1: 109 (1855)	4b:109
e	<b>Oreobolus oligocephalus</b> W.M.Curtis, Brunonia 7: 301 (1985)	4b:110
e	<b>Oreobolus oxycarpus</b> S.T.Blake subsp. <b>brownii</b> Seberg, Bot. J. Linn. Soc. 96: 178 (1988)	4b:110
	<b>Oreobolus pumilio</b> R.Br. subsp. <b>pumilio</b> , Prodr. Fl. Nov. Holland. 236 (1810)	4b:109
e	<b>Oreobolus tholicarpus</b> D.I.Morris, Muelleria 15: 28 (2001)	
	Schoenoides oligocephalus (W.M.Curtis) Seberg = <b>Oreobolus oligocephalus</b>	4b:110
	<b>Schoenoplectus pungens</b> (Vahl) Palla, Bot. Jahrb. Syst. 10: 299 (1888)	4b:94
	<b>Schoenoplectus tabernaemontani</b> (C.C.Gmel.) Palla, Bot. Jahrb. Syst. 10: 299 (1888)	4b:95
	Schoenoplectus validus (Vahl) Á.Löve & D.Löve = <b>Schoenoplectus tabernaemontani</b>	4b:95
e	<b>Schoenus absconditus</b> Kük., Repert. Spec. Nov. Regni Veg. 44: 99 (1938)	4b:117
	Schoenus acutus Labill. = <b>Baumea acuta</b>	
	<b>Schoenus apogon</b> Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 2: 77 (1817)	4b:114
	Schoenus axillaris (R.Br.) Poir. = <b>Schoenus maschalinus</b>	4b:116
e	<b>Schoenus biglumis</b> Kük., Repert. Spec. Nov. Regni Veg. 44: 164 (1938)	4b:118
	<b>Schoenus brevifolius</b> R.Br., Prodr. Fl. Nov. Holland. 231 (1810)	4b:113
	Schoenus brownii Hook.f. = <b>Schoenus apogon</b>	4b:114
	<b>Schoenus calyptratus</b> Kük., Repert. Spec. Nov. Regni Veg. 48: 248 (1940)	4b:117
	<b>Schoenus carsei</b> Cheeseman, Man. New Zealand Fl. 781 (1906)	4b:113
	Schoenus filum Labill. = <b>Gahnia filum</b>	
	<b>Schoenus fluitans</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 81 (1858)	4b:114



	<b>Schoenus latelaminatus</b> Kük., Repert. Spec. Nov. Regni Veg. 44: 88 (1938)	4b:116
	<b>Schoenus lepidosperma</b> (F.Muell.) K.L.Wilson subsp. <b>lepidosperma</b> , Telopea 5: 619 (1994)	4b:114
	<b>Schoenus maschalinus</b> Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 2: 77 (1817)	4b:116
	<b>Schoenus nitens</b> (R.Br.) Poir., Encycl. (Lamarck) Suppl. 2: 252 (1812)	4b:118
e	<b>Schoenus pygmaeus</b> S.T.Blake, Contr. Queensland Herb. 8: 41 (1969)	4b:116
	<i>Schoenus tenuissimus</i> (Hook.f.) Benth. sensu Rodway (1903), Curtis & Morris (1994) =	
	<b>Schoenus lepidosperma</b> subsp. <b>lepidosperma</b>	4b:114
	<b>Schoenus tesquorum</b> J.M.Black, Fl. S. Austral. [J.M. Black] 1: 90 (1922)	4b:115
	<b>Schoenus turbinatus</b> (R.Br.) Poir., Encycl. (Lamarck) Suppl. 2: 251 (1811)	4b:118
	<i>Scirpus americanus</i> Pers. sensu Willis (1970) = <b>Schoenoplectus pungens</b>	4b:94
	<i>Scirpus antarcticus</i> L. sensu Willis (1970) = <b>Isolepis marginata</b>	4b:104
	<i>Scirpus aucklandicus</i> (Hook.f.) Boeckeler = <b>Isolepis aucklandica</b>	4b:102
	<i>Scirpus caldwelii</i> V.J.Cook = <b>Bolboschoenus caldwelii</b>	4b:92
	<i>Scirpus cartilagineus</i> (R.Br.) Poir. = <b>Isolepis marginata</b>	4b:104
	<i>Scirpus cartilagineus</i> (R.Br.) Spreng. var. <i>propinqua</i> (Nees) Benth. = <b>Isolepis inundata</b>	
	<i>Scirpus cernuus</i> Vahl = <b>Isolepis cernua</b>	4b:99
	<i>Scirpus crassiusculus</i> (Hook.f.) Benth. = <b>Isolepis crassiuscula</b>	4b:98
	<i>Scirpus fluitans</i> L. = <b>Isolepis fluitans</b>	4b:98
	<i>Scirpus fluviatilis</i> (Torr.) A.Gray sensu Willis (1970) = <b>Bolboschoenus medianus</b>	4b:93
	<i>Scirpus hookerianus</i> (Boeckeler) S.T.Blake = <b>Isolepis hookeriana</b>	4b:104
	<i>Scirpus inundatus</i> (R.Br.) Poir. = <b>Isolepis inundata</b>	4b:100
	<i>Scirpus lacustris</i> L. sensu Rodway (1903) = <b>Schoenoplectus tabernaemontani</b>	4b:95
	<i>Scirpus lenticularis</i> (R.Br.) Poir. sensu Rodway (1903) = <b>Isolepis fluitans</b>	4b:98
	<i>Scirpus maritimus</i> L. sensu Rodway (1903) = <b>Bolboschoenus caldwelii</b>	4b:92
	<i>Scirpus montivagus</i> S.T.Blake = <b>Isolepis montivaga</b>	4b:103
	<i>Scirpus nodosus</i> Rottb. = <b>Ficinia nodosa</b>	4b:105
	<i>Scirpus platycarpus</i> S.T.Blake = <b>Isolepis platycarpa</b>	4b:99
	<i>Scirpus productus</i> C.B.Clarke = <b>Isolepis producta</b>	4b:98
	<i>Scirpus pungens</i> Vahl = <b>Schoenoplectus pungens</b>	4b:94
	<i>Scirpus riparius</i> (R.Br.) Poir. sensu Rodway (1903) = <b>Isolepis cernua</b>	4b:99
	<i>Scirpus setaceus</i> L. = <b>Isolepis setacea</b>	4b:105
	<i>Scirpus stellatus</i> C.B.Clarke = <b>Isolepis stellata</b>	4b:104
	<i>Scirpus subtilissimus</i> (Boeckeler) S.T.Blake = <b>Isolepis subtilissima</b>	4b:101
	<i>Scirpus tasmanicus</i> S.T.Blake = <b>Isolepis tasmanica</b>	4b:102
	<i>Scirpus triqueter</i> L. sensu Brown (1810) = <b>Schoenoplectus pungens</b>	
	<i>Scirpus validus</i> Vahl = <b>Schoenoplectus tabernaemontani</b>	4b:95
	<i>Scirpus wakefieldianus</i> S.T.Blake = <b>Isolepis wakefieldiana</b>	4b:100
	<b>Tetraria capillaris</b> (F.Muell.) J.M.Black, Trans. Roy. Soc. South Australia 58: 169 (1934)	4b:132
	<b>Tricostularia pauciflora</b> (F.Muell.) Benth., Fl. Austral. 7: 383 (1878)	4b:127
	<b>Uncinia compacta</b> R.Br., Prodr. Fl. Nov. Holland. 241 (1810)	4b:142
t	<b>Uncinia elegans</b> (Kük.) Hamlin, Rec. Domin. Mus. 19: 11 (1959)	4b:140
	<b>Uncinia flaccida</b> S.T.Blake, Proc. Roy. Soc. Queensland 51: 49 (1940)	4b:143
t	<b>Uncinia nervosa</b> Boott, Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 102 (1858)	4b:142
	<b>Uncinia riparia</b> R.Br., Prodr. Fl. Nov. Holland. 241 (1810)	4b:140

	<b>Uncinia tenella</b> R.Br., Prodr. Fl. Nov. Holland. 241 (1810)	4b:143
<b>HAEMODORACEAE</b>		
i #	<b>Anigozanthos flavidus</b> DC., Liliac. (Redouté) 3: t. 176 (1807)	
e	<b>Haemodorum distichophyllum</b> Hook., Icon. Pl. 9: t.866 (1851)	4b:364
<b>HYDATELLACEAE</b>		
	Hydatella filamentosa (Rodway) W.M.Curtis = <b>Trithuria filamentosa</b>	4b:361
e	<b>Trithuria filamentosa</b> Rodway, Pap. & Proc. Roy. Soc. Tasmania 1897: 48 (1898)	4b:361
	<b>Trithuria submersa</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 79 (1858)	4b:360
<b>HYDROCHARITACEAE</b>		
i	<b>Egeria densa</b> Planch., Ann. Sci. Nat., Bot., sér. 3, 11: 80 (1849)	4b:8
i	<b>Elodea canadensis</b> Michx., Fl. Bor.-Amer. (Michaux) 1: 20 (1803)	4b:8
	Elodea densa (Planch.) Casp. = <b>Egeria densa</b>	4b:8
	<b>Halophila australis</b> Doty & B.C.Stone, Brittonia 18: 306, fig.2 (1966)	4b:13
	Halophila ovalis (R.Br.) Hook.f. sensu Rodway (1903) = <b>Halophila australis</b>	4b:13
n	Lagarosiphon major (Ridl.) Moss – previously listed as naturalised but insufficient evidence exists to support this	4b:9
	Vallisneria americana Michx. sensu Buchanan (2007) = <b>Vallisneria australis</b>	4b:11
	<b>Vallisneria australis</b> S.W.L.Jacobs & Les, Syst. Bot. 33: 62 (2008)	4b:11
	Vallisneria gigantea Graebn. sensu Curtis & Morris (1994) = <b>Vallisneria australis</b>	4b:11
	Vallisneria spiralis L. sensu Rodway (1903) = <b>Vallisneria australis</b>	4b:11
<b>IRIDACEAE</b>		
i	<b>Chasmanthe floribunda</b> (Salisb.) N.E.Br., Trans. Roy. Soc. South Africa 20: 274 (1932)	
i	<b>Crocoshmia × crocosmiiflora</b> (Lemoine ex E.Morren) N.E.Br., Trans. Roy. Soc. South Africa 20: 264 (1932)	4b:417
e	<b>Diplarrena latifolia</b> Benth., Fl. Austral. 6: 400 (1873)	4b:405
	<b>Diplarrena moraea</b> Labill., Voy. Rech. Pérouse 1: 157 (1800)	4b:405
	Diplarrena moraea Labill. var. alpina Hook.f. = <b>Diplarrena moraea</b>	
i	<b>Freesia</b> a complex of garden hybrids involving <i>F. alba</i> and <i>F. leichtlinii</i>	4b:416
	Genosiris fragilis Labill. = <b>Patersonia fragilis</b>	
i	<b>Gladiolus × colvillii</b> Sweet, Brit. Fl. Gard. [Sweet] 2: t.155 (1826)	
i	<b>Gladiolus communis</b> L. subsp. <b>byzantinus</b> (Mill.) A.P.Ham., Bot. J. Linn. Soc. 76: 358 (1978)	4b:413
	Gladiolus cuspidatus Jacq. = <b>Gladiolus undulatus</b>	4b:413
i	<b>Gladiolus tristis</b> L., Sp. Pl., ed. 2: 53 (1762)	4b:411
i	<b>Gladiolus undulatus</b> L., Mant. Pl. 1: 27 (1767)	4b:413
i t	<b>Hesperantha coccinea</b> (Backh. & Harv.) Goldblatt & J.C.Manning, Novon 6: 263 (1996)	4b:417
	Hewardia tasmanica Hook. = <b>Isophysis tasmanica</b>	4b:401
	Homeria breyniana (L.) G.J.Lewis sensu Willis (1973) = <b>Moraea flaccida</b>	4b:408
	Homeria flaccida Sweet = <b>Moraea flaccida</b>	4b:408
i	<b>Iris foetidissima</b> L., Sp. Pl. 1: 39 (1753)	4b:406
i	<b>Iris germanica</b> L., Sp. Pl. 1: 38 (1753)	4b:406
e	<b>Isophysis tasmanica</b> (Hook.) T.Moore, Proc. Linn. Soc. London 2: 212 (1853)	4b:401
i	<b>Ixia flexuosa</b> L., Sp. Pl., ed. 2: 51 (1762)	4b:414
i	<b>Ixia maculata</b> L., Sp. Pl., ed. 2: 1664 (1763)	4b:414
i	<b>Ixia polystachya</b> L., Sp. Pl., ed. 2: 51 (1762)	4b:414

	Libertia laurencei Hook.f. = <b>Libertia pulchella</b>	
	<b>Libertia pulchella</b> (R.Br.) Spreng. var. <b>pulchella</b> , Syst. Veg. (ed. 16) [Sprengel] 1: 69 (1824)	4b:402
e	<b>Libertia pulchella</b> (R.Br.) Spreng. var. <b>pygmaea</b> D.I.Morris, The Student's Flora of Tasmania 4b: 425 (1994)	4b:402
i	<b>Moraea flaccida</b> (Sweet) Steud., Nomencl. Bot. [Steudel], ed. 2. 2: 160 (1841)	4b:408
	<b>Patersonia fragilis</b> (Labill.) Asch. & Graebn., Syn. Mitteleur. Fl. [Ascherson & Graebner]. 3: 532 (1906)	4b:403
	Patersonia glauca R.Br. = <b>Patersonia fragilis</b>	4b:403
	Patersonia longiscapa Sweet = <b>Patersonia occidentalis</b>	4b:403
	<b>Patersonia occidentalis</b> R.Br. var. <b>occidentalis</b> , Prodr. Fl. Nov. Holland. 304 (1810)	4b:403
	Romulea longifolia (Salisb.) Baker = <b>Romulea rosea</b> var. <b>australis</b>	4b:409
	Romulea minutiflora Klatt sensu Buchanan (2009) probably <b>Romulea rosea</b>	
i	<b>Romulea rosea</b> (L.) Eckl. var. <b>australis</b> (Ewart) M.P.de Vos, J. S. African Bot. Suppl. 9: 254 (1972)	4b:409
	Schizostylis coccinea Backh. & Harv. = <b>Hesperantha coccinea</b>	4b:417
i	<b>Sisyrinchium iridifolium</b> Kunth, Nov. Gen. Sp. [H.B.K.] 1: 324 (1816)	4b:402
i	<b>Sparaxis bulbifera</b> (L.) Ker Gawl., Ann. Bot. [König & Sims] 1: 226 (1804)	4b:415
	Trichonema roseum Ker Gawl. sensu Rodway (1903) = <b>Romulea rosea</b>	4b:409
i #	<b>Tritonia gladiolaris</b> (Lam.) Goldblatt & J.C.Manning, Bothalia 36: 57 (2006)	4b:416
	Tritonia lineata (Salisb.) Ker Gawl. = <b>Tritonia gladiolaris</b>	4b:416
	Watsonia bulbillifera J.W.Mathews & L.Bolus = <b>Watsonia meriana</b> var. <b>bulbillifera</b>	4b:408
i	<b>Watsonia meriana</b> (L.) Mill. var. <b>bulbillifera</b> (J.W.Mathews & L.Bolus) D.A.Cooke, J. Adelaide Bot. Gard. 18: 6 (1998)	4b:408
i	<b>Watsonia meriana</b> (L.) Mill. var. <b>meriana</b> , Gard. Dict., ed. 8, no.1 (1768)	4b:409
i	<b>Watsonia versfeldii</b> J.W.Mathews & L.Bolus, Ann. Bolus Herb. 3: 140 (1922)	4b:409
	Watsonia versfeldii J.W.Mathews & L.Bolus var. <b>alba</b> J.W.Mathews & L.Bolus = <b>Watsonia versfeldii</b>	4b:409
<b>JUNCACEAE</b>		
i	<b>Juncus acuminatus</b> Michx., Fl. Bor.-Amer. (Michaux) 1: 192 (1803)	4b:75
i	<b>Juncus acutiflorus</b> Ehrh. ex Hoffm., Deutschl. Fl. (Hoffm.) 1: 125 (1791)	4b:77
i	<b>Juncus acutus</b> L., Sp. Pl. 1: 325 (1753)	
	<b>Juncus amabilis</b> Edgar, New Zealand J. Bot. 2: 186, figs.7, 21 (1964)	4b:67
	<b>Juncus antarcticus</b> Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 79, pl.46 (1844)	4b:71
i	<b>Juncus articulatus</b> L., Sp. Pl. 1: 327 (1753)	4b:77
e	<b>Juncus astreptus</b> L.A.S.Johnson, Aspects of Tasmanian Botany: 41 (1991)	4b:66
	<b>Juncus australis</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 66, t.134a (1858)	4b:67
	<b>Juncus bassianus</b> L.A.S.Johnson, Aspects of Tasmanian Botany: 40 (1991)	4b:66
	<b>Juncus bufonius</b> L., Sp. Pl. 1: 328 (1753)	4b:63
i	<b>Juncus bulbosus</b> L., Sp. Pl. 1: 327 (1753)	4b:73
	<b>Juncus caespiticius</b> E.Mey., Pl. Preiss. [J.G.C.Lehman] 2: 47 (1846)	4b:72
	Juncus capillaceus Hook.f. sensu Rodway (1903) = <b>Juncus sandwithii</b>	4b:74
i	<b>Juncus capitatus</b> Weigel, Observ. Bot. (Weigel): 28 (1772)	4b:70
	Juncus communis E.Mey. sensu Rodway (1903) = <b>Juncus australis</b>	4b:67
i	<b>Juncus conglomeratus</b> L., Sp. Pl. 1: 326 (1753)	4b:69
e	<b>Juncus curtisiae</b> L.A.S.Johnson, Aspects of Tasmanian Botany: 44 (1991)	4b:74

i	<b>Juncus effusus</b> L., Sp. Pl. 1: 326 (1753)	4b:69
	<b>Juncus falcatus</b> E.Mey. subsp. <b>falcatus</b> , Syn. Luzul. 34 (1823)	4b:71
	<b>Juncus filicaulis</b> Buchenau, Proc. Linn. Soc. New South Wales 28: 912 (1904)	4b:68
	<b>Juncus fockei</b> Buchenau, Bot. Jahrb. Syst. 12: 358 (1890)	4b:75
i	<b>Juncus gerardii</b> Loisel., J. Bot. (Desvaux) 2: 284 (1809)	
	<b>Juncus gregiflorus</b> L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 243 (1963)	4b:66
	<i>Juncus gunnii</i> Hook.f. = <b>Juncus pauciflorus</b>	
	<b>Juncus holoschoenus</b> R.Br., Prodr. Fl. Nov. Holland. 259 (1810)	4b:75
i	<b>Juncus indescrptus</b> Steud., Syn. Pl. Glum. 2: 304 (1855)	4b:72
	<b>Juncus kraussii</b> Hochst. subsp. <b>australiensis</b> (Buchenau) Snogerup, Willdenowia 23: 61 (1993)	4b:70
	<i>Juncus maritimus</i> Lam. sensu Rodway (1903) = <b>Juncus kraussii</b> subsp. <b>australiensis</b>	4b:70
i #	<b>Juncus microcephalus</b> Kunth, Bonpland & Kunth, Nov. Gen. Sp. [H.B.K.] 1: 237 (1816)	4b:75
	<b>Juncus pallidus</b> R.Br., Prodr. Fl. Nov. Holland. 258 (1810)	4b:64
	<b>Juncus pauciflorus</b> R.Br., Prodr. Fl. Nov. Holland. 259 (1810)	4b:65
	<b>Juncus planifolius</b> R.Br., Prodr. Fl. Nov. Holland. 259 (1810)	4b:71
	<b>Juncus prismatocarpus</b> R.Br., Prodr. Fl. Nov. Holland. 259 (1810)	4b:73
	<b>Juncus procerus</b> E.Mey., Linnaea 3: 367 (1828)	4b:65
	<i>Juncus pusillus</i> Buchenau sensu Willis (1973) = <b>Juncus sandwithii</b>	4b:74
e	<b>Juncus ratkowskyanus</b> L.A.S.Johnson, Aspects of Tasmanian Botany: 43 (1991)	4b:74
	<b>Juncus revolutus</b> R.Br., Prodr. Fl. Nov. Holland. 259 (1810)	4b:63
	<b>Juncus sandwithii</b> Lourteig, Comite National Francais des Recherches Antarctiques 23: 44 (1968)	4b:74
	<b>Juncus sarophorus</b> L.A.S.Johnson, Contr. New South Wales Natl. Herb. 3: 242 (1963)	4b:68
i t	<b>Juncus squarrosus</b> L., Sp. Pl. 1: 327 (1753)	4b:62
	<b>Juncus subsecundus</b> N.A.Wakef., Vict. Naturalist 73: 211 (1957)	4b:69
i	<b>Juncus tenuis</b> Willd., Sp. Pl., ed. 4 [Willdenow] 2: 214 (1799)	4b:62
	<b>Juncus vaginatus</b> R.Br., Prodr. Fl. Nov. Holland. 258 (1810)	4b:65
	<b>Luzula acutifolia</b> H.Nordensk. subsp. <b>acutifolia</b> , Bot. Not. 122: 85 (1969)	4b:84
	<b>Luzula atrata</b> Edgar, New Zealand J. Bot. 13: 794 (1975)	4b:83
e	<b>Luzula australasica</b> Steud. subsp. <b>australasica</b> , Syn. Pl. Glumac. 2: 294 (1855)	4b:82
i t	<b>Luzula campestris</b> (L.) DC., Fl. Franc. (DC. & Lamarck), ed. 3. 3: 161 (1805)	4b:81
i t	<b>Luzula congesta</b> (Thuill.) Lej., Fl. Spa 1: 168 (1811)	4b:81
	<b>Luzula densiflora</b> (H.Nordensk.) Edgar, New Zealand J. Bot. 13: 786 (1975)	4b:80
	<b>Luzula flaccida</b> (Buchenau) Edgar, New Zealand J. Bot. 13: 786 (1975)	4b:80
	<b>Luzula meridionalis</b> H.Nordensk., Bot. Not. 122: 71 (1969)	4b:78
	<b>Luzula modesta</b> Buchenau, Pflanzenr. (Engler) 25: 80, fig.51 (1906)	4b:82
i t	<b>Luzula multiflora</b> (Ehrh.) Lej., Fl. Spa 1: 169 (1811)	4b:81
	<b>Luzula novae-cambriae</b> Gand., Bull. Soc. Bot. France 46: 392 (1900)	4b:82
	<i>Luzula oldfieldii</i> Hook.f. = <b>Luzula australasica</b> subsp. <b>australasica</b>	4b:82
e	<b>Luzula poimena</b> W.M.Curtis, Brunonia 7: 297 (1985)	4b:83

## JUNCAGINACEAE

	<b>Cycnogeton alcockiae</b> (Aston) Mering & Kadereit, Diversity, Phylogeny, and Evolution in the Monocotyledons: 73 (2010)	4b:427
	<b>Cycnogeton procerum</b> (R.Br.) Buchenau, Abh. Naturwiss. Vereins Bremen 224 (1867)	4b:14
	<b>Cycnogeton rheophilum</b> (Aston) Mering & Kadereit, Diversity, Phylogeny, and Evolution in the Monocotyledons: 73 (2010)	4b:427

	Triglochin alcockiae Aston = <b>Cycnogeton alcockiae</b>	4b:427
	Triglochin centrocarpum Hook. = <b>Triglochin nana</b> (Tasmanian plants)	4b:16
	Triglochin decipiens R.Br. = <b>Triglochin striata</b>	
	<b>Triglochin minutissima</b> F.Muell., Fragm. (Mueller) 6: 82 (1867)	4b:16
	<b>Triglochin mucronata</b> R.Br., Prodr. Fl. Nov. Holland. 343 (1810)	4b:16
	<b>Triglochin nana</b> F.Muell., Fragm. (Mueller) 6: 82 (1867)	4b:16
	Triglochin procera R.Br. = <b>Cycnogeton procerum</b>	4b:14
	Triglochin rheophila Aston = <b>Cycnogeton rheophilum</b>	4b:427
	<b>Triglochin striata</b> Ruiz & Pav., Fl. Peruv. 3: 72 (1802)	4b:16
	Triglochin triandra Michx. sensu Hooker (1860) = <b>Triglochin striata</b>	
<b>LEMNACEAE</b>		
	<b>Landoltia punctata</b> (G.Mey.) Les & D.J.Crawford, Novon 9: 532 (1999)	
	<b>Lemna disperma</b> Hegelm., Bot. Zeitung (Berlin) 29: 655 (1871)	4b:32
	Lemna minor L. sensu Rodway (1903) = <b>Lemna disperma</b>	4b:32
	<b>Lemna trisulca</b> L., Sp. Pl. 2: 970 (1753)	4b:32
	Spirodela polyrhiza (L.) Schleid. probably misapplied to <b>Landoltia punctata</b>	
	Wolffia arrhiza (L.) Horkel ex Wimm. sensu Willis (1973) = <b>Wolffia australiana</b>	4b:32
	<b>Wolffia australiana</b> (Benth.) Hartog & Plas, Blumea 20: 151 (1972)	4b:32
<b>LILIACEAE</b>		
i	<b>Agapanthus praecox</b> Willd. subsp. <b>orientalis</b> (F.M.Leight.) F.M.Leight., J. S. African Bot. 4: 21 (1965)	
	Aletris punicea Labill. = <b>Blandfordia punicea</b>	
i	<b>Allium ampeloprasum</b> L., Sp. Pl. 1: 294 (1753)	
i	<b>Allium neapolitanum</b> Cirillo, Pl. Rar. Neap. 1: 13, t.4 (1788)	4b:392
i	<b>Allium triquetrum</b> L., Sp. Pl. 1: 300 (1753)	4b:392
i	<b>Allium vineale</b> L., Sp. Pl. 1: 299 (1753)	4b:392
i #	<b>Alstroemeria aurea</b> Graham, Edinburgh New Philos. J. 15: 181 (1833)	
	Anguillaria dioica R.Br. pro spec. Tasm. = <b>Wurmbea uniflora</b> , <b>W. dioica</b> & <b>W. latifolia</b>	
	Anguillaria uniflora R.Br. = <b>Wurmbea uniflora</b>	
	Anthericum semibarbatum R.Br. = <b>Bulbine semibarbata</b>	
	Arthropodium laxum Hook.f. nom. illeg. = <b>Arthropodium strictum</b>	
	<b>Arthropodium milleflorum</b> (DC.) J.F.Macbr., Contr. Gray Herb. 3(56): 2 (1918)	4b:384
	<b>Arthropodium minus</b> R.Br., Prodr. Fl. Nov. Holland. 276 (1810)	4b:385
	Arthropodium paniculatum (Haw.) R.Br. = <b>Arthropodium milleflorum</b>	4b:384
e	<b>Arthropodium pendulum</b> DC., Cat. Pl. Horti Monsp. 80 (1813)	
	<b>Arthropodium strictum</b> R.Br., Prodr. Fl. Nov. Holland. 276 (1810)	4b:388
i	<b>Asparagus asparagoides</b> (L.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 3: 414 (1914)	
i	<b>Asparagus officinalis</b> L., Sp. Pl. 1: 313 (1753)	4b:370
i	<b>Asparagus scandens</b> Thunb., Prod. Pl. Cap. 66 (1794)	
i	<b>Asphodelus fistulosus</b> L., Sp. Pl. 1: 309 (1753)	4b:380
e	<b>Astelia alpina</b> R.Br. var. <b>alpina</b> , Prodr. Fl. Nov. Holland. 291 (1810)	4b:370
e	<b>Astelia sp. Quartzite (A.M.Buchanan 16828) Tas Herbarium</b>	
	Astelia stylosa F.Muell. ex Hook.f. = <b>Milligania stylosa</b>	
	Blandfordia grandiflora Herb. var. <b>backhousii</b> (Lindl.) Hook.f. = <b>Blandfordia punicea</b>	
	Blandfordia marginata Herb. = <b>Blandfordia punicea</b>	4b:374
e	<b>Blandfordia punicea</b> (Labill.) Sweet, Hort. Brit. [Sweet], ed 2: 517 (1830)	4b:374

	<b>Bulbine bulbosa</b> (R.Br.) Haw., Revis. Pl. Succ. 33 (1821)	4b:383
	<b>Bulbine crassa</b> D.I.Morris & Duretto, Muelleria 22: 93 (2006)	
	<b>Bulbine glauca</b> (Raf.) E.M.Watson, Fl. Australia 45: 469 (1987)	4b:382
	<b>Bulbine semibarbata</b> (R.Br.) Haw., Revis. Pl. Succ. 33 (1821)	4b:382
	<b>Burchardia umbellata</b> R.Br., Prodr. Fl. Nov. Holland. 273 (1810)	4b:396
	<b>Caesia alpina</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 373 (1859)	4b:386
	<b>Caesia calliantha</b> R.J.F.Hend., Fl. Australia 45: 471 (1987)	4b:387
	<i>Caesia corymbosa</i> R.Br. = <b>Chamaescilla corymbosa</b>	
	<b>Caesia parviflora</b> R.Br. var. <b>minor</b> R.J.F.Hend., Fl. Australia 45: 471 (1987)	4b:386
	<b>Caesia parviflora</b> R.Br. var. <b>parviflora</b> , Prodr. Fl. Nov. Holland. 277 (1810)	4b:386
	<b>Caesia parviflora</b> R.Br. var. <b>vittata</b> (R.Br.) R.J.F.Hend., Fl. Australia 45: 472 (1987)	4b:386
	<i>Caesia vittata</i> R.Br. sensu Rodway (1903) = <b>Caesia calliantha</b>	4b:387
	<i>Caesia vittata</i> R.Br. = <b>Caesia parviflora</b> var. <b>vittata</b>	
e	<b>Campynema lineare</b> Labill., Nov. Holl. Pl. 1: 93, t.121 (1805)	4b:395
	<i>Campynema pygmaeum</i> F.Muell. ex Benth. = <b>Campynema lineare</b>	
	<b>Chamaescilla corymbosa</b> (R.Br.) F.Muell. ex Benth. var. <b>corymbosa</b> , Fl. Austral. 7: 48 (1878)	4b:387
	<i>Chlorophytum alpinum</i> (Hook.f.) Benth. ex Baker = <b>Caesia alpina</b>	4b:386
	<b>Dianella amoena</b> G.W.Carr & P.F.Horsfall, Muelleria 8: 369 (1995)	4b:377
	<i>Dianella archeri</i> Hook.f. = <b>Dianella tasmanica</b>	
	<b>Dianella brevicaulis</b> (Ostenf.) G.W.Carr & P.F.Horsfall, Muelleria 8: 375 (1995)	4b:378
	<i>Dianella caerulea</i> Sims sensu Henderson (1987) = <b>Dianella amoena</b> (Tasmanian plants)	
	<i>Dianella laevis</i> R.Br. = <b>Dianella amoena</b>	4b:377
	<i>Dianella longifolia</i> R.Br. sensu Curtis & Morris (1994) = <b>Dianella amoena</b>	4b:377
	<i>Dianella longifolia</i> R.Br. var. <i>aspera</i> Rodway – a name of uncertain application	
	<i>Dianella revoluta</i> R.Br. var. <b>brevicaulis</b> Ostenf. = <b>Dianella brevicaulis</b>	4b:378
	<b>Dianella revoluta</b> R.Br. var. <b>revoluta</b> , Prodr. Fl. Nov. Holland. 280 (1810)	4b:378
	<b>Dianella tasmanica</b> Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 57, t.133A (1858)	4b:378
	<i>Dichopogon strictus</i> (R.Br.) Baker = <b>Arthropodium strictum</b>	4b:388
	<b>Drymophila cyanocarpa</b> R.Br., Prodr. Fl. Nov. Holland. 292 (1810)	4b:369
	<b>Herpolirion novae-zelandiae</b> Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 258 (1853)	4b:383
	<i>Herpolirion tasmaniae</i> Hook.f. = <b>Herpolirion novae-zelandiae</b>	
	<i>Hypoxis glabella</i> R.Br. var. <i>glabella</i> = <b>Pauridia glabella</b> var. <b>glabella</b>	
	<b>Hypoxis hygrometrica</b> Labill. var. <b>hygrometrica</b> , Nov. Holl. Pl. 1: 82 t.108 (1805)	4b:376
	<b>Hypoxis hygrometrica</b> Labill. var. <b>villosisepala</b> R.J.F.Hend., Fl. Australia 45: 488 (1987)	4b:377
	<i>Hypoxis pusilla</i> Hook.f. = <b>Pauridia glabella</b> var. <b>glabella</b>	4b:375
	<i>Hypoxis vaginata</i> Schltld. var. <b>brevistigmata</b> R.J.F.Hend. = <b>Pauridia vaginata</b> var. <b>brevistigmata</b>	4b:376
	<i>Hypoxis vaginata</i> Schltld. var. <i>vaginata</i> = <b>Pauridia vaginata</b> var. <b>vaginata</b>	4b:376
	<i>Laxmannia minor</i> R.Br. sensu Hooker (1860) = <b>Laxmannia orientalis</b>	
	<i>Laxmannia morrisii</i> Keighery = <i>Laxmannia squarrosa</i>	
	<b>Laxmannia orientalis</b> Keighery, Fl. Australia 45: 492 (1987)	4b:388
	<i>Laxmannia sessiliflora</i> Decne. sensu Rodway (1903) = <b>Laxmannia orientalis</b>	4b:388
	<i>Laxmannia squarrosa</i> Lindl. recorded in error	
i	<b>Leucjum aestivum</b> L., Syst. Nat., ed. 10. 2: 975 (1759)	
e	<b>Milligania densiflora</b> Hook.f., Hooker's J. Bot. Kew Gard. Misc. 5: 298 (1853)	4b:374

e	<b>Milligania johnstonii</b> F.Muell. ex Benth., Fl. Austral. 7: 26 (1878)	4b:371
e	<b>Milligania lindoniana</b> Rodway ex W.M.Curtis, Records of the Queen Victoria Museum 45: 1 (1972)	4b:372
e	<b>Milligania longifolia</b> Hook.f., Hooker's J. Bot. Kew Gard. Misc. 5: 297 (1853)	4b:373
e	<b>Milligania stylosa</b> (F.Muell. ex Hook.f.) F.Muell. ex Benth., Fl. Austral. 7: 27 (1878)	4b:373
i	<b>Muscari armeniacum</b> Leichtlin ex Baker, Gard. Chron. 798 (1878) Myrsiphyllum asparagoides (L.) Willd. = <b>Asparagus asparagoides</b> Myrsiphyllum scandens (Thunb.) Oberm. = <b>Asparagus scandens</b>	
i	<b>Narcissus pseudonarcissus</b> L., Sp. Pl. 1: 289 (1753)	4b:394
i	<b>Narcissus tazetta</b> L., Sp. Pl. 1: 290 (1753)	
i	<b>Nothoscordum borbonicum</b> Kunth, Enum. Pl. [Kunth] 4: 463 (1843) Nothoscordum gracile (Aiton) Stearn sensu Curtis & Morris (1994) = <b>Nothoscordum borbonicum</b>	4b:394 4b:394
	Ornithogalum angustifolium Boreau sensu Buchanan (1999) = <b>Ornithogalum umbellatum</b>	
i	<b>Ornithogalum umbellatum</b> L., Sp. Pl. 1: 307 (1753)	4b:369
	<b>Pauridia glabella</b> (R.Br.) Snijman & Kocyan var. <b>glabella</b> , Phytotaxa 116: 27 (2013)	4b:375
	<b>Pauridia vaginata</b> (Schltdl.) Snijman & Kocyan var. <b>brevistigmata</b> (R.J.F.Hend.) Snijman & Kocyan, Phytotaxa 116: 31 (2013)	4b:376
	<b>Pauridia vaginata</b> (Schltdl.) Snijman & Kocyan var. <b>vaginata</b> (Schltdl.) Snijman & Kocyan, Phytotaxa 116: 31 (2013)	4b:376
i #	<b>Scilla peruviana</b> L., Sp. Pl. 1: 309 (1753)	4b:391
	<b>Sowerbaea juncea</b> Andrews, Bot. Repos. 2: t.81 (1800)	4b:389
	Stypantra caespitosa R.Br. = <b>Thelionema caespitosum</b>	4b:379
	Stypantra umbellata R.Br. = <b>Thelionema umbellatum</b>	
	<b>Thelionema caespitosum</b> (R.Br.) R.J.F.Hend., Austrobaileya 2: 109 (1985)	4b:379
	<b>Thelionema umbellatum</b> (R.Br.) R.J.F.Hend., Austrobaileya 2: 109 (1985)	4b:380
	<b>Thysanotus patersonii</b> R.Br., Prodr. Fl. Nov. Holland. 284 (1810)	4b:389
	<b>Tricoryne elatior</b> R.Br., Prodr. Fl. Nov. Holland. 278 (1810)	4b:391
	<b>Wurmbea biglandulosa</b> (R.Br.) T.D.Macfarl. subsp. <b>biglandulosa</b> , Brunonia 3: 191 (1980)	
	<b>Wurmbea dioica</b> (R.Br.) F.Muell. subsp. <b>dioica</b> , Fragm. (Mueller) 10: 119 (1877)	4b:396
	<b>Wurmbea latifolia</b> T.D.Macfarl. subsp. <b>vanessae</b> R.J.Bates, J. Adelaide Bot. Gard. 16: 48 (1995)	4b:398
	<b>Wurmbea uniflora</b> (R.Br.) T.D.Macfarl., Brunonia 3: 194 (1980)	4b:398

## ORCHIDACEAE

	Acianthus bifolius R.Br. = <b>Chiloglottis reflexa</b>	
	<b>Acianthus caudatus</b> R.Br., Prodr. Fl. Nov. Holland. 321 (1810)	4a:117
	Acianthus exsertus R.Br. sensu Curtis (1979) = <b>Acianthus pusillus</b>	4a:117
	<b>Acianthus pusillus</b> D.L.Jones, Austral. Orchid Res. 2: 7 (1991)	4a:117
	Acianthus reniformis (R.Br.) Schltr. = <b>Cyrtostylis reniformis</b>	4a:119
	Acianthus viridis Hook.f. = <b>Townsonia viridis</b>	4a:118
	Anzybas fordhamii (Rupp) D.L.Jones & M.A.Clem. = <b>Corybas fordhamii</b>	4a:124
	Anzybas unguiculatus (R.Br.) D.L.Jones & M.A.Clem. = <b>Corybas unguiculatus</b>	4a:124
	Arachnorchis anthracina (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia anthracina</b>	
	Arachnorchis australis (G.W.Carr) D.L.Jones & M.A.Clem. = <b>Caladenia australis</b>	
	Arachnorchis brachyscapa (G.W.Carr) D.L.Jones & M.A.Clem. = <b>Caladenia brachyscapa</b>	
	Arachnorchis cardiochila (Tate) D.L.Jones & M.A.Clem. = <b>Caladenia cardiochila</b>	

	Arachnorchis caudata (Nicholls) D.L.Jones & M.A.Clem. = <b>Caladenia caudata</b>	4a:101
	Arachnorchis clavigera (A.Cunn. ex Lindl.) D.L.Jones & M.A.Clem. = <b>Caladenia clavigera</b>	4a:105
	Arachnorchis dienema (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia dienema</b>	
	Arachnorchis dilatata (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia dilatata</b>	4a:100
	Arachnorchis echidnachila (Nicholls) D.L.Jones & M.A.Clem. = <b>Caladenia echidnachila</b>	4a:101
	Arachnorchis helvina (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia helvina</b>	
	Arachnorchis lindleyana (Rchb.f.) D.L.Jones & M.A.Clem. = <b>Caladenia lindleyana</b>	
	Arachnorchis pallida (Lindl.) D.L.Jones & M.A.Clem. = <b>Caladenia pallida</b>	4a:104
	Arachnorchis patersonii (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia patersonii</b>	4a:100
	Arachnorchis saggicola (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia saggicola</b>	
	Arthrochilus huntianus (F.Muell.) Blaxell subsp. huntianus = <b>Thynninorchis huntiana</b>	4a:89
	Arthrochilus huntianus (F.Muell.) Blaxell subsp. nothofagicola D.L.Jones = <b>Thynninorchis nothofagicola</b>	
	Bunochilus melagrammus (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis melagramma</b>	4a:27
	Bunochilus stenochilus (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis stenochila</b>	
	Bunochilus tunstallii (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem. = <b>Pterostylis tunstallii</b>	
	Bunochilus williamsonii (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis williamsonii</b>	
	<b>Burnettia cuneata</b> Lindl., Gen. Sp. Orchid. Pl. 518 (1840)	4a:92
	<b>Caladenia alata</b> R.Br., Prodr. Fl. Nov. Holland. 324 (1810)	4a:107
	Caladenia alba R.Br. recorded in error	
	<b>Caladenia alpina</b> R.S.Rogers, Trans. Roy. Soc. South Australia 51: 12 (1927)	4a:111
e	<b>Caladenia angustata</b> Lindl., Gen. Sp. Orchid. Pl. 420 (1840)	4a:108
e	<b>Caladenia anthracina</b> D.L.Jones, Austral. Orchid Res. 3: 21 (1998)	
	Caladenia atkinsonii Rodway = <b>Caladenia carnea</b>	4a:107
e	<b>Caladenia atrata</b> D.L.Jones, Muelleria 8: 178 (1994)	4a:111
e	<b>Caladenia atrochila</b> D.L.Jones, Austral. Orchid Res. 3: 23 (1998)	
	<b>Caladenia aurantiaca</b> (R.S.Rogers) Rupp, Proc. Linn. Soc. New South Wales 71: 280 (1947)	
	<b>Caladenia australis</b> G.W.Carr, Indigenous Flora and Fauna Association Miscellaneous Paper 1: 2 (1991)	
	Caladenia barbata Lindl. sensu Hooker (1860) = <b>Pheladenia deformis</b>	
	<b>Caladenia brachyscapa</b> G.W.Carr, Muelleria 6: 439 (1988)	
	Caladenia caerulea R.Br. sensu Bentham (1873) = <i>Cyanicula caerulea</i>	
e	<b>Caladenia campbellii</b> D.L.Jones, Austral. Orchid Res. 3: 25 (1998)	
x	<b>Caladenia cardiochila</b> Tate, Trans. Roy. Soc. South Australia 9: 60, t.2 (1887)	
	<b>Caladenia carnea</b> R.Br., Prodr. Fl. Nov. Holland. 324 (1810)	4a:106
	Caladenia carnea R.Br. var. pygmaea R.S.Rogers sensu Curtis (1953) = <b>Caladenia pusilla</b>	
	Caladenia carnea R.Br. var. quadriseriata Benth. = <b>Caladenia gracilis</b>	
	Caladenia catenata (Sm.) Druce var. catenata sensu Curtis (1979) = <b>Caladenia carnea &amp; C. sylvicola</b>	4a:106
	Caladenia catenata (Sm.) Druce var. exigua sensu Curtis (1979) = <b>Caladenia alata</b>	4a:107
	Caladenia catenata (Sm.) Druce var. gigantea sensu Curtis (1979) = <b>Caladenia carnea</b>	4a:106
	Caladenia catenata (Sm.) Druce var. minor sensu Curtis (1979) = <b>Caladenia mentiens</b>	4a:106
e	<b>Caladenia caudata</b> Nicholls, Vict. Naturalist 64: 231 (1948)	4a:101
	<b>Caladenia clavigera</b> A.Cunn. ex Lindl., Gen. Sp. Orchid. Pl. 422 (1840)	4a:105
	<b>Caladenia congesta</b> R.Br., Prodr. Fl. Nov. Holland. 324 (1810)	4a:112



	Caladenia cordiformis R.S.Rogers sensu Nicholls & Olsen 1941 = <b>Caladenia clavigera</b>	
	Caladenia corynepetala D.L.Jones = <b>Caladenia dilatata</b>	4a:100
e	<b>Caladenia cracens</b> D.L.Jones, Muelleria 9: 46 (1996)	4a:111
	Caladenia cucullata Fitzg. sensu Curtis (1979) = <b>Caladenia atrata &amp; C. gracilis</b>	4a:111
	Caladenia deformis R.Br. = <b>Pheladenia deformis</b>	4a:113
e	<b>Caladenia dienema</b> D.L.Jones, Austral. Orchid Res. 3: 28 (1998)	
	<b>Caladenia dilatata</b> R.Br., Prodr. Fl. Nov. Holland. 325 (1810)	4a:100
	Caladenia dimorpha Fitzg. sensu Curtis (1979) = <b>Caladenia gracilis</b> (?)	4a:109
e	<b>Caladenia echidnachila</b> Nicholls, Pap. & Proc. Roy. Soc. Tasmania 1932: 13, t.6 (1933)	4a:101
	<b>Caladenia filamentosa</b> R.Br., Prodr. Fl. Nov. Holland. 324 (1810)	4a:99
	<b>Caladenia fuscata</b> (Rchb.f.) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 25 (1989)	
	<b>Caladenia gracilis</b> R.Br., Prodr. Fl. Nov. Holland. 324 (1810)	4a:110
e	<b>Caladenia helvina</b> D.L.Jones, Austral. Orchid Res. 2: 26, f.30 (1991)	
	Caladenia iridescens R.S.Rogers sensu Curtis (1979) = <b>Caladenia transitoria</b>	4a:112
	<b>Caladenia latifolia</b> R.Br., Prodr. Fl. Nov. Holland. 324 (1810)	4a:105
e	<b>Caladenia lindleyana</b> (Rchb.f.) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 27 (1989)	
	Caladenia longii R.S.Rogers = <b>Caladenia angustata</b>	4a:108
	Caladenia lyallii Hook.f. sensu Curtis (1979) = <b>Caladenia cracens &amp; C. alpina</b>	4a:111
	<b>Caladenia mentiens</b> D.L.Jones, Austral. Orchid Res. 3: 35 (1998)	4a:106
	Caladenia menziesii R.Br. = <b>Leptoceras menziesii</b>	4a:99
e	<b>Caladenia pallida</b> Lindl., Gen. Sp. Orchid. Pl. 421 (1840)	4a:104
	<b>Caladenia patersonii</b> R.Br., Prodr. Fl. Nov. Holland. 324 (1810)	4a:100
	Caladenia patersonii R.Br. var. dilatata (R.Br.) Benth. = <b>Caladenia dilatata</b>	
	Caladenia praecox Nicholls sensu Curtis (1979) = <b>Caladenia angustata</b> (?)	4a:109
	<b>Caladenia prolata</b> D.L.Jones, Austral. Orchid Res. 2: 30 (1991)	
	<b>Caladenia pusilla</b> W.M.Curtis, The Student's Flora of Tasmania 4a: 133 (1980)	4a:107
	Caladenia reticulata Fitzg. sensu Curtis (1979) = <b>Caladenia lindleyana</b>	
e	<b>Caladenia saggicola</b> D.L.Jones, Austral. Orchid Res. 3: 39 (1998)	
	Caladenia simulans G.W.Carr = <b>Caladenia dilatata</b>	
	Caladenia suaveolens (R.Br.) Rchb.f. = <b>Lyperanthus suaveolens</b>	4a:94
e	<b>Caladenia sylvicola</b> D.L.Jones, Austral. Orchid Res. 3: 40 (1998)	
	Caladenia testacea R.Br. sensu Rodway (1903) = <b>Caladenia atrata</b>	
e	<b>Caladenia tonellii</b> D.L.Jones, Austral. Orchid Res. 3: 41 (1998)	
	<b>Caladenia transitoria</b> D.L.Jones, Austral. Orchid Res. 3: 42 (1998)	4a:112
	Caladenia tutelata R.S.Rogers = <b>×Calassodia tutelata</b>	4a:113
	Caladenia venusta G.W.Carr sensu Buchanan (1995) = <b>Caladenia saggicola</b>	
	<b>Caladenia vulgaris</b> D.L.Jones, Austral. Orchid Res. 2: 34 (1991)	
	<b>×Calassodia tutelata</b> (R.S.Rogers) M.A.Clem., Austral. Orchid Res. 1: 33 (1989)	4a:113
	<b>Caleana major</b> R.Br., Prodr. Fl. Nov. Holland. 329 (1810)	4a:87
	Caleana minor R.Br. = <b>Paracaleana minor</b>	4a:88
	<b>Calochilus campestris</b> R.Br., Prodr. Fl. Nov. Holland. 320 (1810)	4a:53
e	<b>Calochilus herbaceus</b> Lindl., Gen. Sp. Orchid. Pl. 459 (1840)	4a:55
	<b>Calochilus imberbis</b> R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 51: 4 (1927)	4a:53
	<b>Calochilus paludosus</b> R.Br., Prodr. Fl. Nov. Holland. 320 (1810)	4a:56
	<b>Calochilus platychilus</b> D.L.Jones, Orchadian 15: 547 (2008)	4a:55

	Calochilus robertsonii Benth. sensu Curtis (1979) = <b>Calochilus platychilus</b>	4a:55
	Calochilus saprophyticus R.S.Rogers sensu Curtis (1953) = <b>Calochilus campestris</b>	
	Calonema filamentosum (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia filamentosa</b>	4a:99
	Calonemorchie filamentosa (R.Br.) Szlach. = <b>Caladenia filamentosa</b>	4a:99
	<b>Chiloglottis cornuta</b> Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 69 (1844)	4a:85
	Chiloglottis diphylla R.Br. sensu Rodway (1903) = <b>Chiloglottis reflexa</b>	4a:86
e	<b>Chiloglottis grammata</b> G.W.Carr, Indigenous Flora and Fauna Association Miscellaneous Paper 1: 20 (1991)	4a:86
e	<b>Chiloglottis gunnii</b> Lindl., Gen. Sp. Orchid. Pl. 387 (1840)	4a:84
	Chiloglottis × pescottiana R.S.Rogers sensu Curtis (1953, 1979) = <b>Chiloglottis grammata</b>	4a:86
	Chiloglottis platychila G.W.Carr = <b>Chiloglottis gunnii</b>	
	<b>Chiloglottis reflexa</b> (Labill.) Druce, Rep. Bot. Soc. Exch. Club Brit. Isles 1916, Suppl. 2: 614 (1917)	4a:86
	<b>Chiloglottis trapeziformis</b> Fitzg., Austral. Orch. 1(3): t.9 (1877)	4a:86
e	<b>Chiloglottis triceratops</b> D.L.Jones, Austral. Orchid Res. 3: 66 (1998)	
	Chiloglottis trilabra Fitzg. = <b>Chiloglottis reflexa</b> (Tasmanian plants)	
	<b>Chiloglottis valida</b> D.L.Jones, Austral. Orchid Res. 2: 43 (1991)	
n e	<b>Chiloglottis sp. Wielangta (M.Wapstra 934) Tas Herbarium</b>	
	<b>Corunastylis archeri</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 460 (2002)	4a:82
e	<b>Corunastylis brachystachya</b> (Lindl.) D.L.Jones & M.A.Clem., Orchadian 13: 460 (2002)	4a:79
	<b>Corunastylis despectans</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 460 (2002)	4a:76
e	<b>Corunastylis firthii</b> (Cady) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002)	4a:80
	<b>Corunastylis morrisii</b> (Nicholls) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002)	4a:83
	<b>Corunastylis nuda</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002)	4a:82
e	<b>Corunastylis nudiscapa</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002)	4a:79
	<b>Corunastylis pumila</b> (Hook.f.) D.L.Jones & M.A.Clem., Orchadian 13: 461 (2002)	4a:80
e	<b>Corunastylis tasmanica</b> (D.L.Jones) D.L.Jones & M.A.Clem., Orchadian 13: 462 (2002)	4a:77
	<b>Corybas aconitiflorus</b> Salisb., Parad. Lond. 1(2): t.83 (1806)	4a:123
	<b>Corybas diemenicus</b> (Lindl.) Rchb.f., Beitr. Syst. Pflanzenk. 42 (1871)	4a:121
	Corybas diemenicus (Lindl.) Rupp sensu Curtis (1979) = <b>Corybas incurvus</b>	4a:121
	Corybas dilatatus (Rupp & Nicholls) Rupp & Nicholls ex Rupp = <b>Corybas diemenicus</b>	4a:121
	<b>Corybas fimbriatus</b> (R.Br.) Rchb.f., Beitr. Syst. Pflanzenk. 42 (1871)	4a:121
	<b>Corybas fordhamii</b> (Rupp) Rupp, Vict. Naturalist 59: 61 (1942)	4a:124
	<b>Corybas incurvus</b> D.L.Jones & M.A.Clem., Kew Bull. 43: 135 (1988)	4a:121
	<b>Corybas unguiculatus</b> (R.Br.) Rchb.f., Beitr. Syst. Pflanzenk. 43 (1871)	4a:124
	Corysanthes bicalcarata R.Br. = <b>Corybas aconitiflorus</b>	4a:123
	Corysanthes diemenica Lindl. = <b>Corybas diemenicus</b>	4a:121
	Corysanthes dilatata Rupp & Nicholls sensu Nicholls & Olsen 1941 = <b>Corybas diemenicus</b>	
	Corysanthes fimbriata R.Br. = <b>Corybas fimbriatus</b>	4a:121
	Corysanthes incurva (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem. = <b>Corybas incurvus</b>	4a:121
	Corysanthes pruinosa R.Cunn. sensu Rodway (1903) = <b>Corybas incurvus</b>	4a:121
	Corysanthes unguiculata R.Br. = <b>Corybas unguiculatus</b>	
	Crangonorchis pedoglossa (Fitzg.) D.L.Jones & M.A.Clem. = <b>Pterostylis pedoglossa</b>	4a:13
	<b>Cryptostylis leptochila</b> F.Muell. ex Benth., Fl. Austral. 6: 334 (1873)	4a:126
	Cryptostylis longifolia R.Br. = <b>Cryptostylis subulata</b>	4a:125

	<b>Cryptostylis subulata</b> (Labill.) Rchb.f., Beitr. Syst. Pflanzenk. 15 (1871)	4a:125
	Cyanicula caerulea (R.Br.) Hopper & A.P.Br. recorded in error	
	<b>Cyrtostylis reniformis</b> R.Br., Prodr. Fl. Nov. Holland. 322 (1810)	4a:119
	<b>Cyrtostylis robusta</b> D.L.Jones & M.A.Clem., Lindleyana 2: 157 (1987)	
	Dendrobium striolatum Rchb.f. = <b>Dockrillia striolata</b>	4a:129
	Diplodium alatum (Labill.) D.L.Jones & M.A.Clem. = <b>Pterostylis alata</b>	4a:20
	Diplodium atrans (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis atrans</b>	4a:22
	Diplodium decurvum (R.S.Rogers) D.L.Jones & M.A.Clem. = <b>Pterostylis decurva</b>	4a:22
	Diplodium grandiflorum (R.Br.) D.L.Jones & M.A.Clem. = <b>Pterostylis grandiflora</b>	4a:20
	Dipodium punctatum (Sm.) R.Br. sensu Curtis (1979) = <b>Dipodium roseum</b>	4a:131
	<b>Dipodium roseum</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 2: 51 (1991)	4a:131
i	<b>Disa bracteata</b> Sw., Kongl. Vetensk. Acad. Nya Handl. 21: 211 (1800)	
	Disperis alata Labill. = <b>Pterostylis alata</b>	
	<b>Diuris chryseopsis</b> D.L.Jones, Austral. Orchid Res. 3: 74 (1998)	4a:34
	Diuris corymbosa Lindl. sensu Curtis (1979) = <b>Diuris orientis</b>	4a:36
	Diuris flavopurpurea Messmer sensu Curtis (1979) = <b>Diuris pardina</b>	4a:35
e	<b>Diuris lanceolata</b> Lindl., Gen. Sp. Orchid. Pl. 508 (1840)	4a:34
	Diuris longifolia R.Br. sensu Curtis (1979) = <b>Diuris orientis</b>	4a:36
	Diuris maculata Sm. sensu Curtis (1979) = <b>Diuris pardina</b>	4a:34
	<b>Diuris monticola</b> D.L.Jones, Austral. Orchid Res. 3: 76 (1998)	
	<b>Diuris orientis</b> D.L.Jones, Austral. Orchid Res. 3: 77 (1998)	4a:36
	Diuris palachila R.S.Rogers sensu Curtis (1979) = <b>Diuris chryseopsis</b> × <b>D. pardina</b>	4a:35
	<b>Diuris palustris</b> Lindl., Gen. Sp. Orchid. Pl. 507 (1840)	4a:32
	<b>Diuris pardina</b> Lindl., Gen. Sp. Orchid. Pl. 507 (1840)	4a:34
	Diuris pedunculata R.Br. sensu Curtis (1979) = <b>Diuris chryseopsis</b> p.p.maj.	4a:34
	<b>Diuris sulphurea</b> R.Br., Prodr. Fl. Nov. Holland. 316 (1810)	4a:36
e	<b>Dockrillia striolata</b> (Rchb.f.) Rauschert subsp. <b>chrysantha</b> D.L.Jones, Austral. Orchid Res. 3: 9 (1998)	
	<b>Dockrillia striolata</b> (Rchb.f.) Rauschert subsp. <b>striolata</b> , Feddes Repert. 94(7-8): 447 (1983)	
	Epipactis cucullata Labill. = <b>Eriochilus cucullatus</b>	
	Epipactis reflexa Labill. = <b>Chiloglottis reflexa</b>	
	Eriochilus autumnalis R.Br. sensu Rodway (1903) = <b>Eriochilus cucullatus</b>	4a:91
	<b>Eriochilus cucullatus</b> (Labill.) Rchb.f., Beitr. Syst. Pflanzenk. 27 (1871)	4a:91
	Eriochilus magenteus D.L.Jones sensu Baker & Duretto (2011) = <b>Eriochilus cucullatus</b> possibly reported in error	
	<b>Gastrodia procera</b> G.W.Carr, Indigenous Flora and Fauna Association Miscellaneous Paper 1: 22 (1991)	
	<b>Gastrodia sesamoides</b> R.Br., Prodr. Fl. Nov. Holland. 330 (1810)	4a:127
	<b>Gastrodia surcula</b> D.L.Jones, Orchadian 15: 554 (2008)	
	Genoplesium archeri (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis archeri</b>	4a:82
	Genoplesium brachystachyum (Lindl.) D.L.Jones & M.A.Clem. = <b>Corunastylis brachystachya</b>	4a:79
	Genoplesium despectans (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis despectans</b>	4a:76
	Genoplesium firthii (Cady) D.L.Jones = <b>Corunastylis firthii</b>	4a:80
	Genoplesium morrisii (Nicholls) D.L.Jones & M.A.Clem. = <b>Corunastylis morrisii</b>	4a:83
	Genoplesium nudiscapum (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis nudiscapa</b>	4a:79

Genoplesium nudum (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis nuda</b>	4a:82
Genoplesium pumilum (Hook.f.) D.L.Jones & M.A.Clem. = <b>Corunastylis pumila</b>	4a:80
Genoplesium tasmanicum D.L.Jones = <b>Corunastylis tasmanica</b>	4a:77
×Glossadenia tutelata (R.S.Rogers) Kavulak nom. illeg. = × <b>Calassodia tutelata</b>	
<b>Glossodia major</b> R.Br., Prodr. Fl. Nov. Holland. 326 (1810)	4a:90
Gunnia australis Lindl. sensu Hooker (1860) = <b>Sarcochilus australis</b>	
<b>Hydrorchis orbicularis</b> (R.S.Rogers) D.L.Jones & M.A.Clem., Orchadian 13: 462 (2002)	4a:60
Hymenochilus cycnocephalus (Fitzg.) D.L.Jones & M.A.Clem. sensu Buchanan (2005) = <b>Pterostylis ziegeleri</b>	4a:26
Hymenochilus muticus (R.Br.) D.L.Jones & M.A.Clem. = <b>Pterostylis mutica</b>	4a:25
Hymenochilus pratensis (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis pratensis</b>	
Hymenochilus rubenachii (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis rubenachii</b>	
Hymenochilus wapstrarum (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis wapstrarum</b>	
Hymenochilus ziegeleri (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis ziegeleri</b>	
Jonesiopsis filamentosa (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia filamentosa</b>	4a:99
<b>Leptoceras menziesii</b> (R.Br.) Lindl., Gen. Sp. Orchid. Pl. 416 (1840)	4a:99
Linguella nana (R.Br.) D.L.Jones & M.A.Clem. = <b>Pterostylis nana</b>	4a:14
Lyperanthus nigricans R.Br. = <b>Pyrorchis nigricans</b>	4a:93
<b>Lyperanthus suaveolens</b> R.Br., Prodr. Fl. Nov. Holland. 325 (1810)	4a:94
Malaxis subulata Labill. = <b>Cryptostylis subulata</b>	
<b>Microtidium atratum</b> (Lindl.) D.L.Jones & M.A.Clem., Orchadian 13: 463 (2002)	4a:60
<b>Microtis arenaria</b> Lindl., Gen. Sp. Orchid. Pl. 396 (1840)	4a:58
Microtis atrata Lindl. = <b>Microtidium atratum</b>	4a:60
Microtis biloba Nicholls = <b>Microtis arenaria</b>	4a:58
<b>Microtis oblonga</b> R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 47: 339 (1923)	4a:59
Microtis orbicularis R.S.Rogers = <b>Hydrorchis orbicularis</b>	4a:60
<b>Microtis parviflora</b> R.Br., Prodr. Fl. Nov. Holland. 321 (1810)	4a:59
Microtis porrifolia (Sw.) R.Br. ex Spreng. var. parviflora (R.Br.) Rodway nom. illeg., sensu Rodway (1903) = <b>Microtis parviflora</b>	
Microtis pulchella R.Br. sensu Hooker (1860), misapplied to <b>Microtis unifolia</b> and <b>M. arenaria</b>	
Microtis rara R.Br. sensu Curtis (1979) = <b>Microtis oblonga</b>	4a:59
<b>Microtis unifolia</b> (G.Forst.) Rchb.f., Beitr. Syst. Pflanzenk. 62 (1871)	4a:58
Nemacianthus caudatus (R.Br.) D.L.Jones & M.A.Clem. = <b>Acianthus caudatus</b>	4a:117
Oligochaetochilus commutatus (D.L.Jones) Szlach. = <b>Pterostylis commutata</b>	
Oligochaetochilus pratensis (D.L.Jones) Szlach. = <b>Pterostylis pratensis</b>	
Oligochaetochilus rubenachii (D.L.Jones) Szlach. = <b>Pterostylis rubenachii</b>	
Oligochaetochilus squamatus (R.Br.) Szlach. = <b>Pterostylis squamata</b>	4a:27
Oligochaetochilus stenochilus (D.L.Jones) Szlach. = <b>Pterostylis stenochila</b>	
Oligochaetochilus wapstrarum (D.L.Jones) Szlach. = <b>Pterostylis wapstrarum</b>	
Oligochaetochilus williamsonii (D.L.Jones) Szlach. = <b>Pterostylis williamsonii</b>	
Oligochaetochilus ziegeleri (D.L.Jones) Szlach. = <b>Pterostylis ziegeleri</b>	
<b>Orthoceras strictum</b> R.Br., Prodr. Fl. Nov. Holland. 317 (1810)	4a:30
<b>Paracaleana minor</b> (R.Br.) Blaxell, Contr. New South Wales Natl. Herb. 4: 281 (1972)	4a:88
Petalochilus alatus (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia alata</b>	
Petalochilus atrochilus (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia atrochila</b>	

	Petalochilus aurantiacus (R.S.Rogers) D.L.Jones & M.A.Clem. = <b>Caladenia aurantiaca</b>	
	Petalochilus campbellii (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia campbellii</b>	
	Petalochilus carneus (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia carnea</b>	4a:106
	Petalochilus fuscatus (Rchb.f.) D.L.Jones & M.A.Clem. = <b>Caladenia fuscata</b>	
	Petalochilus mentiensi (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia mentiensi</b>	
	Petalochilus prolatus (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia prolata</b>	
	Petalochilus pusillus (W.M.Curtis) D.L.Jones & M.A.Clem. = <b>Caladenia pusilla</b>	4a:107
	Petalochilus sylvicola (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia sylvicola</b>	
	Petalochilus tonellii (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia tonellii</b>	
	Petalochilus vulgaris (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia vulgaris</b>	
	<b>Pheladenia deformis</b> (R.Br.) D.L.Jones & M.A.Clem., Orchadian 13: 411 (2001)	4a:113
	Plumatichilos plumosum (Cady) Szlach. = <b>Pterostylis plumosa</b>	4a:24
	Plumatichilos tasmanicum (D.L.Jones) Szlach. = <b>Pterostylis tasmanica</b>	
	Prasopphyllum album R.S.Rogers sensu Curtis (1979) identity unresolved	4a:69
	Prasopphyllum alpestre D.L.Jones = <b>Prasopphyllum mimulum</b> (misapplied in Tasmania)	
e	<b>Prasopphyllum alpinum</b> R.Br., Prodr. Fl. Nov. Holland. 318 (1810)	4a:75
e	<b>Prasopphyllum amoenum</b> D.L.Jones, Austral. Orchid Res. 3: 99 (1998)	
e	<b>Prasopphyllum apoxychilum</b> D.L.Jones, Austral. Orchid Res. 3: 100 (1998)	
	Prasopphyllum archeri Hook.f. = <b>Corunastylis archeri</b>	4a:82
e	<b>Prasopphyllum sp. Arthurs Lake (R.Smith DLJ11363) Tas Herbarium</b>	
e	<b>Prasopphyllum atratum</b> D.L.Jones, Austral. Orchid Res. 5: 144 (2006)	
	<b>Prasopphyllum australe</b> R.Br., Prodr. Fl. Nov. Holland. 318 (1810)	4a:66
	Prasopphyllum beaugleholei Nicholls = <b>Corunastylis nuda</b>	4a:82
	Prasopphyllum brachystachyum Lindl. = <b>Corunastylis brachystachya</b>	4a:79
	Prasopphyllum brainei R.S.Rogers = <b>Prasopphyllum lindleyanum</b>	4a:74
	<b>Prasopphyllum brevilabre</b> (Lindl.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 11, t. 110A (1858)	4a:68
	Prasopphyllum buftonianum J.H.Willis = <b>Corunastylis pumila</b>	4a:80
e	<b>Prasopphyllum castaneum</b> D.L.Jones, Austral. Orchid Res. 3: 102 (1998)	
e	<b>Prasopphyllum concinnum</b> Nicholls, Vict. Naturalist 64: 232 (1948)	4a:73
	Prasopphyllum correctum D.L.Jones = <b>Prasopphyllum incorrectum</b> (misapplied in Tasmania)	
e	<b>Prasopphyllum crebriflorum</b> D.L.Jones, Muelleria 18: 103 (2003)	
	Prasopphyllum despectans Hook.f. = <b>Corunastylis despectans</b>	4a:76
	<b>Prasopphyllum elatum</b> R.Br., Prodr. Fl. Nov. Holland. 318 (1810)	4a:67
e	<b>Prasopphyllum favonium</b> D.L.Jones, Austral. Orchid Res. 3: 104 (1998)	
	Prasopphyllum firthii Cady = <b>Corunastylis firthii</b>	4a:80
	Prasopphyllum fitzgeraldii R.S.Rogers & Maiden recorded in error	
	<b>Prasopphyllum flavum</b> R.Br., Prodr. Fl. Nov. Holland. 318 (1810)	4a:66
	Prasopphyllum frenchii F.Muell. sensu Curtis (1979) identity unresolved	4a:73
	Prasopphyllum fuscum R.Br. sensu Curtis (1979) identity unresolved	4a:75
	Prasopphyllum gracile Lindl. sensu Curtis (1979) identity unresolved	4a:72
e	<b>Prasopphyllum incorrectum</b> D.L.Jones, Muelleria 18: 107 (2003)	
e	<b>Prasopphyllum incurvum</b> D.L.Jones, Austral. Orchid Res. 3: 106 (1998)	
	Prasopphyllum intricatum C.Stuart ex Benth. = <b>Corunastylis archeri</b>	4a:82
e	<b>Prasopphyllum limnetes</b> D.L.Jones, Austral. Orchid Res. 5: 151 (2006)	

	<b>Prasophyllum lindleyanum</b> Rchb.f., Beitr. Syst. Pflanzenk. 3: 58 (1871)	4a:74
	Prasophyllum lutescens Lindl. sensu Hooker (1860) = <b>Prasophyllum australe</b>	
	Prasophyllum lutescens Lindl. var. brevilabre Lindl. = <b>Prasophyllum brevilabre</b>	
e	<b>Prasophyllum milfordense</b> D.L.Jones, Austral. Orchid Res. 3: 107 (1998)	
e	<b>Prasophyllum mimulum</b> D.L.Jones, Orchadian 14: 372 (2004)	4a:69
	Prasophyllum aff. montanum R.J.Bates & D.L.Jones sensu Buchanan (2009) = <b>Prasophyllum sp.</b>	
	<b>Arthurs Lake (R.Smith DLJI 1363) Tas Herbarium</b>	
	Prasophyllum morrisii Nicholls = <b>Corunastylis morrisii</b>	4a:83
	Prasophyllum nigricans R.Br. sensu Curtis (1979) = <b>Corunastylis tasmanica</b>	4a:77
	Prasophyllum nudiscapum Hook.f. = <b>Corunastylis nudiscapa</b>	4a:79
	Prasophyllum nudum Hook.f. = <b>Corunastylis nuda</b>	
	Prasophyllum odoratum R.S.Rogers sensu Curtis (1979) identity unresolved	4a:68
e	<b>Prasophyllum olidum</b> D.L.Jones, Austral. Orchid Res. 3: 108 (1998)	
	Prasophyllum parviflorum (R.S.Rogers) Nicholls sensu Curtis (1979) identity unresolved	4a:74
	Prasophyllum patens R.Br. sensu Rodway (1903) = <b>Prasophyllum truncatum</b>	4a:71
	Prasophyllum patens R.Br. var. robustum Nicholls sensu Curtis (1953) = <b>Prasophyllum robustum</b>	
e	<b>Prasophyllum perangustum</b> D.L.Jones, Austral. Orchid Res. 3: 109 (1998)	4a:72
e	<b>Prasophyllum pulchellum</b> D.L.Jones, Austral. Orchid Res. 3: 110 (1998)	
	Prasophyllum pyriforme E.Coleman sensu Buchanan (2005) = <b>Prasophyllum rostratum</b>	
e	<b>Prasophyllum robustum</b> (Nicholls) M.A.Clem. & D.L.Jones, Austral. Orchid Res. 1: 117 (1989)	
	Prasophyllum rogersii Rupp sensu Curtis (1979) = <b>Prasophyllum perangustum</b>	4a:72
e	<b>Prasophyllum rostratum</b> Lindl., Gen. Sp. Orchid. Pl. 516 (1840)	
	Prasophyllum rufum R.Br. sensu Curtis (1979) = <b>Corunastylis tasmanica</b>	4a:79
e	<b>Prasophyllum secutum</b> D.L.Jones, Austral. Orchid Res. 3: 113 (1998)	
	<b>Prasophyllum sphacelatum</b> D.L.Jones, Muellera 9: 59 (1996)	
e	<b>Prasophyllum stellatum</b> D.L.Jones, Austral. Orchid Res. 3: 115 (1998)	
	Prasophyllum suttonii R.S.Rogers & B.Rees sensu Curtis (1979) = <b>Prasophyllum mimulum</b>	4a:69
	<b>Prasophyllum tadgellianum</b> (R.S.Rogers) R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 47: 338 (1923)	
e	<b>Prasophyllum taphanyx</b> D.L.Jones, Orchadian 14: 373 (2004)	
e	<b>Prasophyllum truncatum</b> Lindl., Gen. Sp. Orchid. Pl. 513 (1840)	4a:71
e	<b>Prasophyllum tunbridgense</b> D.L.Jones, Austral. Orchid Res. 3: 117 (1998)	
	Prasophyllum uroglossum Rupp sensu Curtis (1979) identity unresolved	4a:76
	Pterostylis acuminata R.Br. sensu Curtis (1953) = <b>Pterostylis ingens</b>	
e	<b>Pterostylis alata</b> (Labill.) Rchb.f., Beitr. Syst. Pflanzenk. 70 (1871)	4a:20
	Pterostylis alpina R.S.Rogers sensu Curtis (1979) = <b>Pterostylis scabrida</b>	4a:19
e	<b>Pterostylis aphylla</b> Lindl., Gen. Sp. Orchid. Pl. 392 (1840)	4a:24
	<b>Pterostylis atrans</b> D.L.Jones, Muellera 8: 185 (1994)	4a:22
e	<b>Pterostylis atriola</b> D.L.Jones, Austral. Orchid Res. 3: 140 (1998)	
	Pterostylis barbata Lindl. sensu Rodway (1903) = <b>Pterostylis plumosa</b>	4a:24
e	<b>Pterostylis commutata</b> D.L.Jones, Muellera 8: 186 (1994)	
	<b>Pterostylis concinna</b> R.Br., Prodr. Fl. Nov. Holland. 326 (1810)	4a:11
	<b>Pterostylis cucullata</b> R.Br. subsp. <b>cucullata</b> , Prodr. Fl. Nov. Holland. 327 (1810)	
	<b>Pterostylis curta</b> R.Br., Prodr. Fl. Nov. Holland. 326 (1810)	4a:12

	<i>Pterostylis cycnocephala</i> Fitzg. sensu Curtis (1979) = <b>Pterostylis ziegeleri</b>	4a:26
	<b>Pterostylis decurva</b> R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 47: 339, t.27 (1923)	4a:22
e	<b>Pterostylis dubia</b> R.Br., Prodr. Fl. Nov. Holland. 328 (1810)	4a:16
	<b>Pterostylis falcata</b> R.S.Rogers, Proc. Roy. Soc. Victoria 28: 106, t.9 (1915)	4a:17
	<b>Pterostylis foliata</b> Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 249 (1853)	4a:19
e	<b>Pterostylis furcata</b> Lindl., Gen. Sp. Orchid. Pl. 390 (1840)	4a:17
	<b>Pterostylis grandiflora</b> R.Br., Prodr. Fl. Nov. Holland. 327 (1810)	4a:20
	<b>Pterostylis × ingens</b> (Rupp) D.L.Jones, Orchadian 5: 54 (1976)	4a:12
	<i>Pterostylis longifolia</i> R.Br. sensu Curtis (1979) = <b>Pterostylis melagramma</b> & <b>P. williamsonii</b>	4a:27
	<b>Pterostylis lustra</b> D.L.Jones, Austral. Orchid Res. 5: 87 (2006)	
	<b>Pterostylis melagramma</b> D.L.Jones, Austral. Orchid Res. 3: 145 (1998)	4a:27
	<b>Pterostylis mutica</b> R.Br., Prodr. Fl. Nov. Holland. 328 (1810)	4a:25
	<b>Pterostylis nana</b> R.Br., Prodr. Fl. Nov. Holland. 327 (1810)	4a:14
	<b>Pterostylis nutans</b> R.Br., Prodr. Fl. Nov. Holland. 327 (1810)	4a:13
	<i>Pterostylis obtusa</i> R.Br. sensu Curtis (1979) = <b>Pterostylis atrans</b>	4a:22
	<b>Pterostylis parviflora</b> R.Br., Prodr. Fl. Nov. Holland. 327 (1810)	4a:23
	<b>Pterostylis pedoglossa</b> Fitzg., Austral. Orchids 1(3): 5th plate (1877)	4a:13
	<b>Pterostylis pedunculata</b> R.Br., Prodr. Fl. Nov. Holland. 327 (1810)	4a:14
	<b>Pterostylis plumosa</b> Cady, Austral. Pl. 5: 138 (1969)	4a:24
	<i>Pterostylis praecox</i> Lindl. sensu Rodway (1903) = <b>Pterostylis alata</b>	4a:20
e	<b>Pterostylis pratensis</b> D.L.Jones, Austral. Orchid Res. 3: 149 (1998)	
	<i>Pterostylis pusilla</i> R.S.Rogers sensu Curtis (1953) = <b>Pterostylis squamata</b>	
e	<b>Pterostylis rubenachii</b> D.L.Jones, Austral. Orchid Res. 3: 150 (1998)	
	<i>Pterostylis rufa</i> R.Br. sensu Curtis (1979) = <b>Pterostylis squamata</b>	4a:27
	<b>Pterostylis sanguinea</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 1: 126 (1989)	4a:28
e	<b>Pterostylis scabrida</b> Lindl., Gen. Sp. Orchid. Pl. 389 (1840)	4a:19
	<b>Pterostylis squamata</b> R.Br., Prodr. Fl. Nov. Holland. 327 (1810)	4a:27
e	<b>Pterostylis stenochila</b> D.L.Jones, Austral. Orchid Res. 3: 153 (1998)	
	<b>Pterostylis tasmanica</b> D.L.Jones, Muelleria 8: 190 (1994)	
	<b>Pterostylis × toveyana</b> Ewart & Sharman, Proc. Roy. Soc. Victoria 28: 235 (1916)	4a:21
	<b>Pterostylis tunstallii</b> D.L.Jones & M.A.Clem., Austral. Orchid Res. 1: 128 (1989)	
	<b>Pterostylis uliginosa</b> D.L.Jones, Austral. Orchid Res. 3: 155 (1998)	
	<i>Pterostylis vereenae</i> R.S.Rogers sensu Curtis (1953) = <b>Pterostylis foliata</b>	
	<i>Pterostylis vittata</i> Lindl. sensu Curtis (1979) = <b>Pterostylis sanguinea</b>	4a:28
e	<b>Pterostylis wapstrarum</b> D.L.Jones, Austral. Orchid Res. 3: 156 (1998)	
e	<b>Pterostylis williamsonii</b> D.L.Jones, Austral. Orchid Res. 3: 157 (1998)	4a:27
e	<b>Pterostylis ziegeleri</b> D.L.Jones, Austral. Orchid Res. 3: 158 (1998)	4a:26
	<b>Pyrorchis nigricans</b> (R.Br.) D.L.Jones & M.A.Clem., Phytologia 77: 449 (1994)	4a:93
	<b>Sarcochilus australis</b> (Lindl.) Rchb.f., Ann. Bot. Syst. (Walpers) 6: 501 (1863)	4a:131
	<i>Sarcochilus parviflorus</i> Lindl. sensu Rodway (1903) = <b>Sarcochilus australis</b>	4a:131
	<i>Simpliglottis cornuta</i> (Hook.f.) Szlach. = <b>Chiloglottis cornuta</b>	4a:85
	<i>Simpliglottis grammata</i> (G.W.Carr) Jeanes = <b>Chiloglottis grammata</b>	
	<i>Simpliglottis gunnii</i> (Lindl.) Szlach. = <b>Chiloglottis gunnii</b>	4a:84
	<i>Simpliglottis triceratops</i> (D.L.Jones) Jeanes = <b>Chiloglottis triceratops</b>	
	<i>Simpliglottis valida</i> (D.L.Jones) Szlach. = <b>Chiloglottis valida</b>	

	<i>Specularantha aphylla</i> (Lindl.) D.L.Jones & M.A.Clem. = <b>Pterostylis aphylla</b>	4a:24
	<i>Specularantha atriola</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis atriola</b>	
	<i>Specularantha parviflora</i> (R.Br.) D.L.Jones & M.A.Clem. = <b>Pterostylis parviflora</b>	4a:23
	<i>Specularantha uliginosa</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Pterostylis uliginosa</b>	
	<i>Spiculaea huntiana</i> (F.Muell.) Schltr. = <b>Thynninorchis huntiana</b>	4a:89
	<b>Spiranthes alticola</b> D.L.Jones, <i>Orchadian</i> 15: 555 (2008)	
	<b>Spiranthes australis</b> (R.Br.) Lindl., <i>Bot. Reg.</i> 10: subt.823 (1824)	4a:128
	<i>Spiranthes sinensis</i> (Pers.) Ames subsp. <i>australis</i> (R.Br.) Kitam. = <b>Spiranthes australis</b>	4a:128
	<i>Stegostyla alpina</i> (R.S.Rogers) D.L.Jones & M.A.Clem. = <b>Caladenia alpina</b>	4a:111
	<i>Stegostyla angustata</i> (Lindl.) D.L.Jones & M.A.Clem. = <b>Caladenia angustata</b>	4a:108
	<i>Stegostyla atrata</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia atrata</b>	
	<i>Stegostyla congesta</i> (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia congesta</b>	4a:112
	<i>Stegostyla cracens</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia cracens</b>	
	<i>Stegostyla gracilis</i> (R.Br.) D.L.Jones & M.A.Clem. = <b>Caladenia gracilis</b>	4a:110
	<i>Stegostyla transitoria</i> (D.L.Jones) D.L.Jones & M.A.Clem. = <b>Caladenia transitoria</b>	
	<i>Sullivania minor</i> (R.Br.) D.L.Jones & M.A.Clem. = <b>Paracaleana minor</b>	4a:88
	<i>Taurantha concinna</i> (R.Br.) D.L.Jones & M.A.Clem. = <b>Pterostylis concinna</b>	4a:11
	× <i>Taurodium toveyanum</i> (Ewart & Sharman) D.L.Jones & M.A.Clem. sensu Buchanan (2005) = <b>Pterostylis toveyana</b>	
e	<b>Thelymitra aggericola</b> D.L.Jones, <i>The Orchadian</i> 12: 517 (1999)	
	<i>Thelymitra angustifolia</i> R.Br. sensu Hooker (1860) – a name of uncertain application	
	<b>Thelymitra antennifera</b> (Lindl.) Hook.f., <i>Bot. Antarct. Voy. Ill.</i> (Fl. Tasman.) 2: 4, t.101A (1858)	4a:51
	<b>Thelymitra arenaria</b> Lindl., <i>Gen. Sp. Orchid. Pl.</i> 519 (1840)	
	<b>Thelymitra aristata</b> Lindl., <i>Gen. Sp. Orchid. Pl.</i> 521 (1840)	4a:42
	<b>Thelymitra atronitida</b> Jeanes, <i>Muelleria</i> 14: 91 (2000)	
	<i>Thelymitra azurea</i> R.S.Rogers sensu Buchanan (1995) = <b>Thelymitra jonesii</b>	
	<b>Thelymitra benthamiana</b> Rchb.f., <i>Beitr. Syst. Pflanzenk.</i> 55 (1871)	
	<b>Thelymitra bracteata</b> J.Z.Weber ex Jeanes, <i>Muelleria</i> 19: 43 (2004)	
	<b>Thelymitra brevifolia</b> Jeanes, <i>Muelleria</i> 19: 30 (2004)	
	<i>Thelymitra canaliculata</i> R.Br. sensu Curtis (1979) = <b>Thelymitra jonesii</b>	4a:46
	<b>Thelymitra carnea</b> R.Br., <i>Prodr. Fl. Nov. Holland.</i> 314 (1810)	4a:50
	<i>Thelymitra chasmogama</i> R.S.Rogers sensu Curtis (1979) a plant of uncertain hybrid origin	4a:47
	<b>Thelymitra circumsepta</b> Fitzg., <i>Austral. Orchids</i> 1: t.1 (1878)	4a:48
	<b>Thelymitra cyanea</b> (Lindl.) Benth., <i>Fl. Austral.</i> 6: 323 (1873)	4a:49
	<i>Thelymitra decora</i> Cheeseman sensu Buchanan (1995) = <b>Thelymitra simulata</b>	
	<b>Thelymitra erosa</b> D.L.Jones & M.A.Clem., <i>Austral. Orchid Res.</i> 3: 184 (1998)	
	<b>Thelymitra exigua</b> Jeanes, <i>Muelleria</i> 19: 28 (2004)	
	<b>Thelymitra flexuosa</b> Endl., <i>Nov. Stirp. Dec.</i> 3: 23 (1839)	4a:51
	<i>Thelymitra grandiflora</i> Fitzg. sensu Curtis (1953) = <b>Thelymitra aristata</b>	
	<b>Thelymitra holmesii</b> Nicholls, <i>Vict. Naturalist</i> 49: 263 (1933)	4a:43
e	<b>Thelymitra imbricata</b> D.L.Jones & M.A.Clem., <i>Austral. Orchid Res.</i> 3: 186 (1998)	4a:43
	<b>Thelymitra improcera</b> D.L.Jones & M.A.Clem., <i>Austral. Orchid Res.</i> 3: 187 (1998)	4a:46
	<b>Thelymitra inflata</b> Jeanes, <i>Muelleria</i> 19: 71 (2004)	
	<b>Thelymitra × irregularis</b> Nicholls, <i>Vict. Naturalist</i> 63: 126 (1946)	4a:48
	<b>Thelymitra ixiooides</b> Sw., <i>Ksvenska Vet. Akad. Handl.</i> 21: 253 (1800)	4a:44



- e **Thelymitra jonesii** Jeanes, Muellera 15: 81 (2001) 4a:46  
**Thelymitra juncifolia** Lindl., Gen. Sp. Orchid. Pl. 522 (1840)  
*Thelymitra longifolia* J.R.Forst. & G.Forst. sensu Bentham (1873) = **Thelymitra nuda**  
**Thelymitra longiloba** D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 191 (1998) 4a:46  
**Thelymitra lucida** Jeanes, Muellera 19: 70 (2004)  
*Thelymitra luteocilium* Fitzg. sensu Curtis (1979) = **Thelymitra rubra** 4a:47  
*Thelymitra* × *macmillanii* F.Muell. sensu Curtis (1979) = a hybrid involving **Thelymitra nuda** 4a:49  
**Thelymitra malvina** M.A.Clem., D.L.Jones & Molloy, Austral. Orchid Res. 1: 141 (1989)  
*Thelymitra media* R.Br. sensu Curtis (1979) = **Thelymitra improcera** & **T. longiloba** 4a:46  
*Thelymitra megalyptra* Fitzg. sensu Curtis (1979) = **Thelymitra imbricata** 4a:43  
**Thelymitra** × **merraniae** Nicholls, Vict. Naturalist 46: 139 (1929)  
**Thelymitra mucida** Fitzg., Gard. Chron. 17: 495 (1882) 4a:46  
**Thelymitra nuda** R.Br., Prodr. Fl. Nov. Holland. 314 (1810) 4a:44  
*Thelymitra nuda* R.Br. var. *grandiflora* Lindl. = **Thelymitra imbricata**  
**Thelymitra pauciflora** R.Br., Prodr. Fl. Nov. Holland. 314 (1810) 4a:42  
*Thelymitra pauciflora* R.Br. var. *holmesii* (Nicholls) Nicholls = **Thelymitra holmesii** 4a:43  
**Thelymitra peniculata** Jeanes, Muellera 19: 50 (2004)  
e **Thelymitra polychroma** D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 193 (1998)  
*Thelymitra pulchella* Hook.f. recorded in error  
*Thelymitra resecta* Rupp = **Thelymitra circumsepta** 4a:48  
**Thelymitra rubra** Fitzg., Gard. Chron. 17: 495 (1882) 4a:50  
e **Thelymitra silena** D.L.Jones, The Orchadian 12: 518 (1999)  
**Thelymitra simulata** D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 195 (1998)  
*Thelymitra smithiana* (Gunn ex Lindl.) Hook.f. = **Thelymitra flexuosa**  
e **Thelymitra spadicea** D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 196 (1998)  
e **Thelymitra sparsa** D.L.Jones & M.A.Clem., Austral. Orchid Res. 3: 197 (1998)  
**Thelymitra** × **truncata** R.S.Rogers, Trans. & Proc. Roy. Soc. South Australia 41: 343, t.17 (1917) 4a:45  
*Thelymitra venosa* R.Br. sensu Curtis (1979) = **Thelymitra cyanea** 4a:48  
*Thelymitra versicolor* Lindl. = **Thelymitra nuda**  
e **Thelymitra viridis** Jeanes, Muellera 19: 36 (2004)  
X **Thynninorchis huntiana** (F.Muell.) D.L.Jones & M.A.Clem., Orchadian 13: 457 (2002) 4a:89  
e **Thynninorchis nothofagicola** (D.L.Jones) D.L.Jones & M.A.Clem., Orchadian 13: 457 (2002)  
t **Townsonia viridis** (Hook.f.) Schltr., Repert. Spec. Nov. Regni Veg. 9: 250 (1911) 4a:118  
*Urochilus sanguineus* (D.L.Jones & M.A.Clem.) D.L.Jones & M.A.Clem. = **Pterostylis sanguinea** 4a:28

**POACEAE (GRAMINEAE)**

- Achnatherum caudatum* (Trin.) S.W.L.Jacobs & J.Everett = **Amelichloa caudata** 4b:192  
*Agropyron elongatum* (Host) P.Beauv. = **Thinopyrum elongatum** 4b:295  
*Agropyron junceiforme* (Á.Löve & D.Löve) Á.Löve & D.Löve = **Thinopyrum junceiforme** 4b:293  
*Agropyron junceum* (L.) P.Beauv. = **Thinopyrum junceiforme** 4b:293  
*Agropyron pectinatum* (Labill.) P.Beauv. = **Australopyrum pectinatum** 4b:301  
*Agropyron repens* (L.) P.Beauv. = **Elytrigia repens** 4b:295  
*Agropyron scabrum* (R.Br.) P.Beauv. = **Anthosachne scabra** 4b:296  
*Agropyron velutinum* Nees = **Australopyrum velutinum** 4b:301  
*Agrostis aemula* R.Br. = **Lachnagrostis aemula**

	<i>Agrostis aemula</i> R.Br. var. <i>aemula</i> = <b>Lachnagrostis aemula</b>	4b:258
	<i>Agrostis aemula</i> R.Br. var. <i>setifolia</i> (Hook.f.) Vickery = <b>Lachnagrostis punicea</b> subsp. <b>punicea</b>	4b:259
	<i>Agrostis aequata</i> Nees = <b>Lachnagrostis rudis</b>	4b:262
	<b>Agrostis australiensis</b> Mez, Feddes Repert. Spec. Nov. Regni Veg. 17: 302 (1921)	4b:257
	<i>Agrostis australiensis</i> Mez sensu Curtis & Morris (1994) = <b>Lachnagrostis rudis</b>	4b:257
	<i>Agrostis</i> sp. aff. <i>australiensis</i> Mez sensu Curtis & Morris (1994) = <b>Agrostis diemenica</b>	4b:257
	<i>Agrostis avenacea</i> J.F.Gmel. = <b>Lachnagrostis filiformis</b>	4b:259
	<b>Agrostis bettyae</b> S.W.L.Jacobs, Telopea 9: 679 (2001)	
	<i>Agrostis billardierei</i> R.Br. var. <i>billardierei</i> = <b>Lachnagrostis billardierei</b> subsp. <b>billardierei</b>	4b:260
	<i>Agrostis billardierei</i> R.Br. var. <i>collicola</i> D.I.Morris = <b>Lachnagrostis collicola</b>	4b:261
	<i>Agrostis billardierei</i> R.Br. var. <i>filifolia</i> Vickery = <b>Lachnagrostis punicea</b> subsp. <b>filifolia</b>	4b:261
	<i>Agrostis billardierei</i> R.Br. var. <i>robusta</i> Vickery = <b>Lachnagrostis robusta</b>	4b:261
	<i>Agrostis billardierei</i> R.Br. var. <i>tenuiseta</i> D.I.Morris = <b>Lachnagrostis billardierei</b> subsp. <b>tenuiseta</b>	4b:261
i	<b>Agrostis capillaris</b> L. var. <i>aristata</i> (Parn.) Druce, Fl. Oxfordshire, ed. 2: 474 (1927)	4b:264
i	<b>Agrostis capillaris</b> L. var. <i>capillaris</i> , Sp. Pl. 1: 62 (1753)	4b:264
	<i>Agrostis contracta</i> F.Muell. ex Hook.f. = <b>Deyeuxia contracta</b>	
	<i>Agrostis crinita</i> (L.f.) R.Br. = <b>Dichelachne crinita</b>	
e	<b>Agrostis diemenica</b> D.I.Morris, Telopea 10: 765 (2004)	4b:257
i	<b>Agrostis gigantea</b> Roth, Tent. Fl. Germ. 1: 31 (1788)	4b:264
	<i>Agrostis</i> sp. aff. <i>hiemalis</i> (Walter) Britton, Sterns & Poggenb. sensu Curtis & Morris (1994) = <b>Agrostis propinqua</b>	4b:258
	<b>Agrostis joyceae</b> S.W.L.Jacobs, Telopea 9: 680 (2001)	
	<i>Agrostis lacunarum</i> D.I.Morris = <b>Lachnagrostis lacunarum</b>	4b:260
	<i>Agrostis lobata</i> R.Br. = <b>Deyeuxia quadriseta</b>	
	<i>Agrostis montana</i> R.Br. = <b>Deyeuxia monticola</b>	
	<b>Agrostis muelleriana</b> Vickery, Contr. New South Wales Natl. Herb. 1: 103 (1941)	4b:256
	<i>Agrostis ovata</i> G.Forst. sensu Labillardiere (1805) = <b>Echinopogon ovatus</b>	
	<b>Agrostis parviflora</b> R.Br., Prodr. Fl. Nov. Holland. 170 (1810)	4b:256
	<i>Agrostis</i> sp. aff. <i>parviflora</i> R.Br. sensu Curtis & Morris (1994) = <b>Agrostis thompsoniae</b>	4b:257
	<b>Agrostis propinqua</b> S.W.L.Jacobs, Telopea 9: 681 (2001)	4b:258
	<i>Agrostis quadrifida</i> Labill. = <b>Pentapogon quadrifidus</b>	
	<i>Agrostis quadriseta</i> (Labill.) R.Br. = <b>Deyeuxia quadriseta</b>	
	<i>Agrostis rudis</i> Roem. & Schult. sensu Curtis & Morris (1994) = <b>Lachnagrostis morrisii</b>	4b:261
	<i>Agrostis scabra</i> R.Br. sensu Rodway (1903) = <b>Agrostis parviflora</b> & <b>A. propinqua</b>	
	<i>Agrostis</i> sp. aff. <i>scabra</i> R.Br. sensu Curtis & Morris (1994) = <b>Agrostis propinqua</b>	4b:258
	<i>Agrostis sciurea</i> R.Br. = <b>Dichelachne micrantha</b>	
	<i>Agrostis semiverticillata</i> (Forssk.) C.Chr. = <b>Agrostis viridis</b>	4b:283
i	<b>Agrostis stolonifera</b> L., Sp. Pl. 1: 62 (1753)	4b:262
	<i>Agrostis tenuis</i> Sibth. = <b>Agrostis capillaris</b>	4b:264
	<b>Agrostis thompsoniae</b> S.W.L.Jacobs, Telopea 9: 682 (2001)	4b:257
	<b>Agrostis venusta</b> Trin., Mém. Acad. Imp. Sci. St.-Petersbourg, Ser. 6 Sci. Math., Seconde Pt. Sci. Nat. 4: 340 (1841)	4b:256
	<i>Agrostis virginica</i> L. sensu Labillardiere (1805) = <b>Sporobolus virginicus</b>	
i	<b>Agrostis viridis</b> Gouan, Hortus Monsp. 546 (1762)	4b:283

	Agrostis vulgaris With. sensu Spicer (1878) = <b>Agrostis capillaris</b>	4b:264
i	<b>Aira caryophyllea</b> L. subsp. <b>caryophyllea</b> , Sp. Pl. 1: 66 (1753)	4b:246
n i *	Aira cupaniana Guss., Fl. Sicul. Syn. 1: 148 (1843)	
	Aira elegans Willd. ex Gaudin = <b>Aira elegantissima</b>	4b:247
i	<b>Aira elegantissima</b> Schur, Verh. Mitth. Siebenbürg. Vereins Naturwiss. Hermannstadt 4: 85 (1853)	4b:247
i	<b>Aira praecox</b> L., Sp. Pl. 1: 65 (1753)	4b:246
	Alopecurus agrestis L. = <b>Alopecurus myosuroides</b>	
i	<b>Alopecurus geniculatus</b> L., Sp. Pl. 1: 60 (1753)	4b:284
i	<b>Alopecurus myosuroides</b> Huds., Fl. Angl. (Hudson): 23 (1762)	4b:285
i	<b>Alopecurus pratensis</b> L. subsp. <b>pratensis</b> , Sp. Pl. 1: 60 (1753)	4b:284
i	<b>Amelichloa caudata</b> (Trin.) Arriaga & Barkworth, Sida 22: 148 (2006)	4b:192
i	<b>Ammophila arenaria</b> (L.) Link subsp. <b>arenaria</b> , Hort. Berol. [Link] 1: 105 (1827)	4b:274
	<b>Amphibromus archeri</b> (Hook.f.) P.Morris, Vict. Naturalist 51: 146 (1934)	4b:232
?i #	<b>Amphibromus fluitans</b> Kirk, Trans. & Proc. New Zealand Inst. 16: 374 (1884)	
	<b>Amphibromus macrorhinus</b> S.W.L.Jacobs & Lapinpuro, Telopea 2: 723 (1986)	4b:233
	<b>Amphibromus neesii</b> Steud., Syn. Pl. Glumac. 1: 328 (1854)	4b:234
	<b>Amphibromus nervosus</b> (Hook.f.) Baill., Hist. Pl. 12: 203 (1894)	
	<b>Amphibromus recurvatus</b> Swallen, Amer. J. Bot. 18: 415 (1931)	4b:232
	<b>Amphibromus sinuatus</b> S.W.L.Jacobs & Lapinpuro, Telopea 2: 727 (1986)	4b:233
	Anthistiria australis R.Br. = <b>Themeda triandra</b>	
	Anthistiria ciliata L.f. sensu Rodway (1903) = <b>Themeda triandra</b>	4b:358
i	<b>Anthosachne kingiana</b> (Endl.) Govaerts subsp. <b>multiflora</b> (Banks & Sol. ex Hook.f.) Govaerts, J. Adelaide Bot. Gard. 27: 24 (2014)	
	<b>Anthosachne scabra</b> (R.Br.) Nevski, Trudy Sredne-Aziatsk. Gosud. Univ., Ser. 8b, Bot. 17: 65 (1934)	4b:296
	Anthoxanthum crinitum L.f. sensu Brown (1810) = <b>Dichelachne crinita</b>	
i	<b>Anthoxanthum odoratum</b> L., Sp. Pl. 1: 28 (1753)	4b:250
i	<b>Aristida benthamii</b> Henrard var. <b>benthamii</b> , Meded. Rijks-Herb. 58A: 246, t.117 (1932)	4b:323
	Aristida jerichoensis (Domin) Henrard sensu Buchanan et al. (1989) = <b>Aristida benthamii</b> var. <b>benthamii</b>	4b:323
	Arrhenatherum avenaceum (Scop.) P.Beauv. sensu Rodway (1903) = <b>Arrhenatherum elatius</b>	4b:234
i	<b>Arrhenatherum elatius</b> (L.) P.Beauv. ex J.Presl & C.Presl var. <b>bulbosum</b> (Willd.) Spenn., Fl. Friburg. 1: 113 (1825)	4b:234
	Arundo penicillatum Labill. = <b>Rytidosperma penicillatum</b>	
	Arundo phragmites L. sensu Brown (1810) = <b>Phragmites australis</b>	
	Arundo poiformis Labill. sensu Labillardiere (1805) = <b>Poa poiformis</b>	
	Arundo semiannularis Labill. = <b>Rytidosperma semiannulare</b>	
	<b>Australopyrum pectinatum</b> (Labill.) Á.Löve, Feddes Repert. 95: 443 (1984)	4b:301
	<b>Australopyrum velutinum</b> (Nees) B.K.Simon, Austrobaileya 2: 241 (1986)	4b:301
	Austrodanthonia alpicola (Vickery) H.P.Linder sensu Fl. Australia 44B: 51 = <b>Rytidosperma oreophilum</b> (Tasmanian plants)	
	Austrodanthonia caespitosa (Gaudich.) H.P.Linder = <b>Rytidosperma caespitosum</b>	4b:312
	Austrodanthonia carphoides (F.Muell. ex Benth.) H.P.Linder = <b>Rytidosperma carphoides</b>	4b:306
	Austrodanthonia diemenica (D.I.Morris) H.P.Linder = <b>Rytidosperma diemenicum</b>	4b:311

	Austrodanthonia geniculata (J.M.Black) H.P.Linder = <b>Rytidosperma geniculatum</b>	4b:308
	Austrodanthonia induta (Vickery) H.P.Linder = <b>Rytidosperma indutum</b>	4b:312
	Austrodanthonia laevis (Vickery) H.P.Linder = <b>Rytidosperma laeve</b>	4b:312
	Austrodanthonia penicillata (Labill.) H.P.Linder = <b>Rytidosperma penicillatum</b>	4b:313
	Austrodanthonia pilosa (R.Br.) H.P.Linder = <b>Rytidosperma pilosum</b>	4b:313
	Austrodanthonia popinensis (D.I.Morris) H.P.Linder = <b>Rytidosperma fulvum</b>	4b:316
	Austrodanthonia procera (Vickery) S.W.L.Jacobs = <b>Rytidosperma indutum</b>	4b:312
	Austrodanthonia racemosa (R.Br.) H.P.Linder = <b>Rytidosperma racemosum</b>	4b:314
	Austrodanthonia racemosa (R.Br.) H.P.Linder var. racemosa = <b>Rytidosperma racemosum</b> var. <b>racemosum</b>	4b:314
	Austrodanthonia remota (D.I.Morris) H.P.Linder = <b>Rytidosperma remotum</b>	4b:316
	Austrodanthonia setacea (R.Br.) H.P.Linder = <b>Rytidosperma setaceum</b>	4b:311
	Austrodanthonia tenuior (Steud.) H.P.Linder = <b>Rytidosperma tenuius</b>	4b:310
i t	<b>Austroderia richardii</b> (Endl.) N.P.Barker & H.P.Linder, Ann. Missouri Bot. Gard. 97: 344 (2010)	4b:319
	Austrofestuca hookeriana (F.Muell. ex Hook.f.) S.W.L.Jacobs = <b>Hookerochloa hookeriana</b>	4b:223
	<b>Austrofestuca littoralis</b> (Labill.) E.B.Alexeev, Byull. Moskovsk. Obshch. Isp. Prir., Otd. Biol. 81(5): 55 (1976)	4b:223
e	<b>Austrostipa aphylla</b> (Rodway) S.W.L.Jacobs & J.Everett, Telopea 6: 584 (1996)	4b:192
	<b>Austrostipa bigeniculata</b> (Hughes) S.W.L.Jacobs & J.Everett, Telopea 6: 584 (1996)	4b:190
	<b>Austrostipa blackii</b> (C.E.Hubb.) S.W.L.Jacobs & J.Everett, Telopea 6: 584 (1996)	4b:191
	<b>Austrostipa flavescens</b> (Labill.) S.W.L.Jacobs & J.Everett, Telopea 6: 585 (1996)	4b:190
	<b>Austrostipa mollis</b> (R.Br.) S.W.L.Jacobs & J.Everett, Telopea 6: 587 (1996)	4b:189
	<b>Austrostipa nodosa</b> (S.T.Blake) S.W.L.Jacobs & J.Everett, Telopea 6: 587 (1996)	4b:188
	<b>Austrostipa pubinodis</b> (Trin. & Rupr.) S.W.L.Jacobs & J.Everett, Telopea 6: 588 (1996)	4b:187
	<b>Austrostipa rudis</b> (Spreng.) S.W.L.Jacobs & J.Everett subsp. <b>australis</b> (J.Everett & S.W.L.Jacobs) S.W.L.Jacobs & J.Everett, Telopea 6: 588 (1996)	4b:186
	<b>Austrostipa scabra</b> (Lindl.) S.W.L.Jacobs & J.Everett subsp. <b>falcata</b> (Hughes) S.W.L.Jacobs & J.Everett, Telopea 6: 588 (1996)	4b:188
	<b>Austrostipa scabra</b> (Lindl.) S.W.L.Jacobs & J.Everett subsp. <b>scabra</b> , Telopea 6: 588 (1996)	4b:188
	<b>Austrostipa semibarbata</b> (R.Br.) S.W.L.Jacobs & J.Everett, Telopea 6: 588 (1996)	4b:189
	<b>Austrostipa stipoides</b> (Hook.f.) S.W.L.Jacobs & J.Everett, Telopea 6: 589 (1996)	4b:186
	<b>Austrostipa stuposa</b> (Hughes) S.W.L.Jacobs & J.Everett, Telopea 6: 589 (1996)	4b:191
n i *	Avellinia michelii (Savi) Parl., Pl. Nov. 61 (1842)	
	Avena alba Vahl sensu Townrow (1969) = <b>Avena barbata</b>	4b:236
i	<b>Avena barbata</b> Pott ex Link, J. Bot. (Shrader) 2: 314,315 (1799)	4b:236
i	<b>Avena fatua</b> L., Sp. Pl. 1: 80 (1753)	4b:237
	Avena filiformis G.Forst. sensu Labillardiere (1805) = <b>Lachnagrostis filiformis</b>	
i	<b>Avena ludoviciana</b> Durieu, Actes Soc. Linn. Bordeaux 20: 41 (1855)	4b:238
	Avena quadriseta Labill. = <b>Deyeuxia quadriseta</b>	
i	<b>Avena sativa</b> L., Sp. Pl. 1: 79 (1753)	4b:237
	Avena sterilis L. subsp. ludoviciana (Durieu) Gillet & Magne = <b>Avena ludoviciana</b>	4b:238
i	<b>Avena strigosa</b> Schreb., Spic. Fl. Lips. 52 (1771)	4b:237
i	<b>Bothriochloa macra</b> (Steud.) S.T.Blake, Proc. Roy. Soc. Queensland 80: 64 (1969)	4b:355
i	<b>Brachypodium distachyon</b> (L.) P.Beauv., Ess. Agrostogr. 101, 155, 156 (1812)	4b:292

i	<b>Briza maxima</b> L., Sp. Pl. 1: 70 (1753)	4b:208
i	<b>Briza minor</b> L., Sp. Pl. 1: 70 (1753)	4b:208
i	<b>Bromus alopecuroides</b> Poir., Voy. Barbarie 2: 100 (1789)	4b:288
	<b>Bromus arenarius</b> Labill., Nov. Holl. Pl. 1: 23, t.28 (1805)	4b:288
i	<b>Bromus brevis</b> Nees ex Steud., Syn. Pl. Glumac. 1: 326 (1854)	4b:290
i	<b>Bromus catharticus</b> Vahl, Symb. Bot. (Vahl) 2: 22 (1791)	4b:290
i	<b>Bromus cecadilla</b> Steud., Syn. Pl. Glumac. 1: 321 (1854)	4b:290
i	<b>Bromus diandrus</b> Roth, Bot. Abh. Beobacht. 44 (1787)	4b:288
	<i>Bromus fonkii</i> Phil. = <b>Bromus lithobius</b>	4b:290
i	<b>Bromus hordeaceus</b> L., Sp. Pl. 1: 77 (1753)	4b:287
	<i>Bromus japonicus</i> Murray var. <i>vestitus</i> sensu Buchanan (2005) = <b>Bromus arenarius</b>	4b:288
i	<b>Bromus lithobius</b> Trin., Linnaea 10: 303 (1836)	4b:290
	<i>Bromus macrostachys</i> Desf. sensu Townrow (1969) = <b>Bromus alopecuroides</b>	4b:288
i	<b>Bromus madritensis</b> L. var. <b>ciliatus</b> Guss., Fl. Sicul. Syn. 1: 78 (1843)	4b:289
	<i>Bromus mollis</i> L. = <b>Bromus hordeaceus</b>	4b:287
i	<b>Bromus sterilis</b> L., Sp. Pl. 1: 77 (1753)	4b:289
	<i>Bromus thominii</i> Hardouin sensu Townrow (1969) = <b>Bromus hordeaceus</b>	4b:287
	<i>Bromus unioloides</i> Kunth = <b>Bromus catharticus</b>	4b:290
	<i>Bromus willdenowii</i> Kunth = <b>Bromus catharticus</b>	4b:290
i # t x	<b>Calamagrostis epigejos</b> (L.) Roth, Tent. Fl. Germ. 1: 34 (1788)	4b:273
i	<b>Catapodium marinum</b> (L.) C.E.Hubb., Kew Bull. 9: 375 (1954)	4b:225
i	<b>Catapodium rigidum</b> (L.) C.E.Hubb., Fl. Bedfordshire: 437 (1953)	4b:224
i	<b>Cenchrus clandestinus</b> (Hochst. ex Chiov.) Morrone, Ann. Bot. (Oxford) 106(1): 127 (2010)	4b:348
i	<b>Cenchrus longisetus</b> M.C.Johnst., Sida 1(3): 182 (1963)	4b:349
i	<b>Cenchrus macrourus</b> (Trin.) Morrone, Ann. Bot. (Oxford) 106(1): 128 (2010)	4b:351
	<i>Cenchrus purpurascens</i> Thunb. recorded in error	4b:351
	<i>Ceratochloa unioloides</i> DC. = <b>Bromus catharticus</b>	4b:290
i	<b>Cortaderia jubata</b> (Lemoine) Stapf, Bot. Mag. 124, t.7607 (1898)	4b:320
	<i>Cortaderia richardii</i> (Endl.) Zotov = <b>Austroderia richardii</b>	4b:319
i	<b>Cortaderia selloana</b> (Schult. & Schult.f.) Asch. & Graebn., Syn. Mitteleur. Fl. [Ascherson & Graebner] 2: 325 (1900)	4b:320
	<i>Critesion marinum</i> (Huds.) Á.Löve = <b>Hordeum marinum</b>	4b:300
	<i>Critesion murinum</i> (L.) Á.Löve = <b>Hordeum murinum</b>	4b:298
i	<b>Cynodon dactylon</b> (L.) Pers. var. <b>dactylon</b> , Syn. Pl. (Persoon) 1: 85 (1805)	4b:330
i	<b>Cynosurus cristatus</b> L., Sp. Pl. 1: 72 (1753)	4b:207
i	<b>Cynosurus echinatus</b> L., Sp. Pl. 1: 72 (1753)	4b:207
i	<b>Dactylis glomerata</b> L., Sp. Pl. 1: 71 (1753)	4b:221
	<i>Danthonia archeri</i> Hook.f. = <b>Amphibromus archeri</b>	
	<i>Danthonia caespitosa</i> Gaudich. = <b>Rytidosperma caespitosum</b>	4b:312
	<i>Danthonia carphoides</i> F.Muell. ex Benth. var. <i>angustior</i> Vickery = <b>Rytidosperma carphoides</b>	4b:306
i	<b>Danthonia decumbens</b> (L.) DC., Lam., Fl. Franc. (DC. & Lamarck), ed. 3. 3: 33 (1805)	4b:317
	<i>Danthonia diemenica</i> D.I.Morris = <b>Rytidosperma diemenicum</b>	4b:311
	<i>Danthonia dimidiata</i> Vickery = <b>Rytidosperma dimidiatum</b>	4b:317
	<i>Danthonia eriantha</i> Lindl. sensu Townrow (1969) identity uncertain	
	<i>Danthonia fortuneae-hibernae</i> Renvoize = <b>Rytidosperma fortuneae-hibernae</b>	4b:310

	Danthonia geniculata J.M.Black = <b>Rytidosperma geniculatum</b>	4b:308
	Danthonia gracilis Hook.f. = <b>Rytidosperma gracile</b>	4b:309
	Danthonia laevis Vickery = <b>Rytidosperma laeve</b>	4b:312
	Danthonia longifolia R.Br. sensu Townrow (1969) = <b>Rytidosperma tenuius</b>	4b:310
	Danthonia nervosa Hook.f. = <b>Amphibromus nervosus</b>	
	Danthonia nitens D.I.Morris = <b>Rytidosperma nitens</b>	4b:315
	Danthonia nivicola Vickery = <b>Rytidosperma nivicola</b>	4b:315
	Danthonia nudiflora P.Morris = <b>Rytidosperma nudiflorum</b>	4b:314
	Danthonia pallida R.Br. sensu Bentham (1878) = <b>Rytidosperma pallidum</b>	
	Danthonia pauciflora R.Br. = <b>Rytidosperma pauciflorum</b>	4b:309
	Danthonia penicillata (Labill.) R.Br. ex P.Beauv. = <b>Rytidosperma penicillatum</b>	4b:313
	Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. pallida (R.Br.) Rodway sensu Rodway (1903) = <b>Rytidosperma pallidum</b>	
	Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. pilosa (R.Br.) Rodway = <b>Rytidosperma pilosum</b>	
	Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. racemosa (R.Br.) Rodway = <b>Rytidosperma racemosum</b>	
	Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. semiannularis (Labill.) Rodway = <b>Rytidosperma semiannulare</b>	
	Danthonia penicillata (Labill.) R.Br. ex P.Beauv. var. setacea (R.Br.) Rodway = <b>Rytidosperma setaceum</b>	
	Danthonia pilosa R.Br. = <b>Rytidosperma pilosum</b>	4b:313
	Danthonia popinensis D.I.Morris = <b>Rytidosperma fulvum</b>	4b:316
	Danthonia procera Vickery = <b>Rytidosperma indutum</b>	4b:312
	Danthonia pulvinorum D.I.Morris = <b>Rytidosperma nivicola</b>	4b:315
	Danthonia purpurascens Vickery sensu Townrow (1969) = <b>Rytidosperma tenuius</b>	4b:310
	Danthonia racemosa R.Br. = <b>Rytidosperma racemosum</b>	4b:314
	Danthonia racemosa R.Br. var. penicillata (Labill.) Benth. nom. illeg. = <b>Rytidosperma penicillatum</b>	
	Danthonia remota D.I.Morris = <b>Rytidosperma remotum</b>	4b:316
	Danthonia semiannularis (Labill.) R.Br. var. gracilis (Hook.f.) Hook.f. = <b>Rytidosperma gracile</b>	4b:308
	Danthonia semiannularis (Labill.) R.Br. var. semiannularis = <b>Rytidosperma semiannulare</b>	4b:309
	Danthonia setacea R.Br. = <b>Rytidosperma setaceum</b>	4b:311
	Danthonia subulata Hook.f. = <b>Rytidosperma setaceum</b>	
	Danthonia tenuior (Steud.) Conert = <b>Rytidosperma tenuius</b>	4b:310
	<b>Deschampsia cespitosa</b> (L.) P.Beauv., Ess. Agrostogr. 91: 160, t.18, fig.3 (1812)	4b:242
t	<b>Deschampsia gracillima</b> Kirk, J. Bot. 29: 237 (1891)	4b:243
	Deyeuxia accedens Vickery = <b>Deyeuxia frigida</b>	4b:271
	Deyeuxia aequata (Nees) Benth. = <b>Lachnagrostis rudis</b>	4b:261
e	<b>Deyeuxia apseyensis</b> D.I.Morris, Muelleria 7: 160 (1990)	4b:273
	Deyeuxia benthamiana Vickery = <b>Deyeuxia scaberula</b>	4b:270
	Deyeuxia billardierei (R.Br.) Kunth = <b>Lachnagrostis billardierei</b>	4b:260
	<b>Deyeuxia brachyathera</b> (Stapf) Vickery, Contr. New South Wales Natl. Herb. 1: 68 (1940)	4b:272
	<b>Deyeuxia carinata</b> Vickery, Contr. New South Wales Natl. Herb. 1: 58 (1940)	4b:272

	<b>Deyeuxia contracta</b> (F.Muell. ex Hook.f.) Vickery, Contr. New South Wales Natl. Herb. 1: 65 (1940)	4b:269
	<b>Deyeuxia decipiens</b> (R.Br.) Vickery, Contr. New South Wales Natl. Herb. 1: 70 (1940)	4b:269
	<b>Deyeuxia densa</b> Benth., Fl. Austral. 7: 582 (1878)	4b:270
	<i>Deyeuxia forsteri</i> Kunth = <b>Lachnagrostis filiformis</b>	4b:259
	<b>Deyeuxia frigida</b> F.Muell. ex Benth., Fl. Austral. 7: 583 (1878)	4b:271
	<b>Deyeuxia gunniana</b> (Nees) Benth., Fl. Austral. 7: 584 (1878)	4b:267
	<b>Deyeuxia innominata</b> D.I.Morris, Muelleria 7: 164 (1990)	4b:268
e x	<b>Deyeuxia lawrencei</b> Vickery, Contr. New South Wales Natl. Herb. 1: 48 (1940)	4b:266
	<b>Deyeuxia minor</b> F.Muell. ex Benth., Fl. Austral. 7: 582 (1878)	4b:270
	<i>Deyeuxia montana</i> (R.Br.) Benth. = <b>Deyeuxia monticola</b>	4b:268
	<b>Deyeuxia monticola</b> (Roem. & Schult.) Vickery, Contr. New South Wales Natl. Herb. 1: 56 (1940)	4b:268
	<i>Deyeuxia parviseta</i> Vickery sensu Townrow (1969) = <b>Deyeuxia innominata</b>	4b:268
	<b>Deyeuxia quadriseta</b> (Labill.) Benth., Fl. Austral. 7: 581 (1878)	4b:267
	<b>Deyeuxia rodwayi</b> Vickery, Contr. New South Wales Natl. Herb. 1: 60 (1940)	4b:271
	<b>Deyeuxia scaberula</b> Vickery, Contr. New South Wales Natl. Herb. 1: 64 (1940)	4b:270
	<i>Deyeuxia scabra</i> Kunth sensu Rodway (1903) = <b>Deyeuxia contracta &amp; D. scaberula</b>	
n	<i>Dichanthium sericeum</i> (R.Br.) A.Camus subsp. <i>sericeum</i> – previously listed as naturalised but insufficient evidence exists to support this	4b:355
	<b>Dichelachne crinita</b> (L.f.) Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 293 (1853)	4b:277
	<b>Dichelachne inaequiglumis</b> (Hack. ex Cheeseman) Edgar & Connor, New Zealand J. Bot. 20: 307 (1982)	4b:278
	<b>Dichelachne micrantha</b> (Cav.) Domin, Biblioth. Bot. 20(85): 353 (1915)	4b:278
	<b>Dichelachne montana</b> Endl., Prodr. Fl. Norfolk. 20 (1833)	
	<b>Dichelachne parva</b> B.K.Simon, Austrobaileya 1: 462 (1982)	
	<b>Dichelachne rara</b> (R.Br.) Vickery, Contr. New South Wales Natl. Herb. 1: 337 (1951)	4b:279
	<i>Dichelachne sciurea</i> (R.Br.) Hook.f. sensu Rodway (1903) = <b>Dichelachne spp.</b> excluding <b>Dichelachne crinita</b>	
	<b>Dichelachne sieberiana</b> Trin. & Rupr., Sp. Gram. Stipac. 2–3 (1842)	
	<i>Dichelachne stipoides</i> Hook.f. = <b>Austrostipa stipoides</b>	
n	<i>Digitaria ciliaris</i> (Retz.) Koeler – previously listed as naturalised but insufficient evidence exists to support this	4b:347
i	<b>Digitaria sanguinalis</b> (L.) Scop., Fl. Carniol., ed. 2. 2: 52 (1772)	4b:346
n	<i>Digitaria ternata</i> (Hochst. ex A.Rich.) Stapf – previously listed as naturalised but insufficient evidence exists to support this	4b:346
	<i>Diplax tasmanica</i> Hook.f. = <b>Microlaena tasmanica</b>	
	<i>Disarrenum antarcticum</i> Labill. = <b>Hierochloe redolens</b>	
	<b>Distichlis distichophylla</b> (Labill.) Fassett, Rhodora 27: 71 (1925)	4b:324
	<i>Distichlis maritima</i> Raf. sensu Rodway (1903) = <b>Distichlis distichophylla</b>	4b:324
	<b>Dryopoa dives</b> (F.Muell.) Vickery, Contr. New South Wales Natl. Herb. 3: 196 (1963)	4b:199
i	<b>Echinochloa crus-galli</b> (L.) P.Beauv., Ess. Agrostogr. 53 (1812)	4b:336
	<i>Echinochloa crus-galli</i> (L.) P.Beauv. var. <i>frumentacea</i> W.Wright = <b>Echinochloa frumentacea</b>	4b:338
i	<b>Echinochloa esculenta</b> (A.Braun) H.Scholz, Taxon 41: 523 (1992)	4b:336
i	<b>Echinochloa frumentacea</b> Link, Hort. Berol. [Link] 1: 204 (1827)	4b:338

n	Echinochloa oryzoides (Ard.) Fritsch – previously listed as naturalised but insufficient evidence exists to support this	
	Echinochloa utilis Ohwi & Yabuno = <b>Echinochloa esculenta</b>	4b:336
	<b>Echinopogon ovatus</b> (G.Forst.) P.Beauv., Ess. Agrostogr. 42, 161, t.9 fig.5 (1812)	4b:275
	Ehrharta acuminata (R.Br.) Spreng. = <b>Tetrarrhena acuminata</b>	4b:182
i	<b>Ehrharta calycina</b> Sm., Pl. Icon. Ined. 2: t.33 (1790)	4b:178
	Ehrharta distichophylla Labill. = <b>Tetrarrhena distichophylla</b>	4b:183
i	<b>Ehrharta erecta</b> Lam. var. <b>erecta</b> , Encycl. (Lamarck) 2: 347 (1786)	4b:179
	Ehrharta juncea (R.Br.) Spreng. = <b>Tetrarrhena juncea</b>	4b:183
i	<b>Ehrharta longiflora</b> Sm., Pl. Icon. Ined. 2: t.32 (1790)	4b:179
	Ehrharta oreophila (D.I.Morris) L.P.M.Willemse var. minor (D.I.Morris) L.P.M.Willemse = <b>Tetrarrhena oreophila</b> var. <b>minor</b>	4b:182
	Ehrharta oreophila (D.I.Morris) L.P.M.Willemse var. oreophila = <b>Tetrarrhena oreophila</b> var. <b>oreophila</b>	4b:182
	Ehrharta stipoides Labill. = <b>Microlaena stipoides</b>	4b:181
	Ehrharta tasmanica (Hook.f.) L.P.M.Willemse var. subalpina (F.Muell. ex Benth.) L.P.M.Willemse = <b>Microlaena tasmanica</b> var. <b>subalpina</b>	4b:182
	Ehrharta tasmanica (Hook.f.) L.P.M.Willemse var. tasmanica = <b>Microlaena tasmanica</b> var. <b>tasmanica</b>	4b:181
i	<b>Ehrharta villosa</b> Schult.f., Syst. Veg., ed. 15 bis [Roemer & Schultes] 7: 1374 (1830)	4b:184
i #	<b>Eleusine indica</b> (L.) Gaertn., Fruct. Sem. Pl. 1: 8 (1788)	
n i	<b>Eleusine tristachya</b> (Lam.) Lam., Tabl. Encycl. 1: 57 (1791)	
	Elymus elongatus (Host) Runemark = <b>Thinopyrum elongatum</b>	4b:295
	Elymus farctus (Viv.) Runemark ex Melderis subsp. boreali-atlanticus (Simonet & Guin.) Melderis = <b>Thinopyrum junceiforme</b>	
	Elymus multiflorus (Banks & Sol. ex Hook.f.) Á.Löve & Connor = <b>Anthosachne kingiana</b> subsp. <b>multiflora</b>	
	Elymus repens (L.) Gould = <b>Elytrigia repens</b>	4b:295
	Elymus scaber (R.Br.) Á.Löve = <b>Anthosachne scabra</b>	4b:296
i	<b>Elytrigia repens</b> (L.) Desv. ex Nevski, Trudy Bot. Inst. Akad. Nauk S.S.S.R., Ser. I, Fl. Sist. Vyssh. Rast. . 1: 14 (1933)	4b:295
i	<b>Enneapogon nigricans</b> (R.Br.) P.Beauv., Ess. Agrostogr. 82, 161, 171 (1812)	
?i	<b>Eragrostis brownii</b> (Kunth) Nees, Cat. Ind. Pl. 2: 105 (1834)	4b:326
i	<b>Eragrostis cilianensis</b> (All.) Vignolo ex Janch., Mitt. Naturwiss. Vereins Univ. Wien 5(9): 110 (1907)	4b:326
n i	<b>Eragrostis curvula</b> (Schrad.) Nees, Fl. Afr. Austral. Ill. 397 (1841)	4b:327
i	<b>Eragrostis minor</b> Host, Icon. Descr. Gram. Austriac. 4: 15 (1809)	
	Eragrostis molybdea Vickery = <b>Eragrostis brownii</b>	4b:326
i	<b>Eragrostis parviflora</b> (R.Br.) Trin., Mém. Acad. Imp. Sci. St.Pétersbourg Hist. Acad. 11: 411 (1830)	
i	<b>Eragrostis pilosa</b> (L.) P.Beauv., Ess. Agrostogr. 71, 162, 175 (1812)	
n i *	Eragrostis tenuifolia (A.Rich.) Hochst. ex Steud., Syn. Pl. Glumac. 1: 268 (1854)	4b:327
	Erythranthera australis (Petrie) Zotov = <b>Rytidosperma australe</b>	4b:318
	Festuca archeri E.B.Alexeev = <b>Festuca arundinacea</b>	4b:428
i	<b>Festuca arundinacea</b> Schreb., Spic. Fl. Lips. 57 (1771)	4b:197
	Festuca asperula Vickery sensu Townrow (1969) = <b>Festuca plebeia</b>	4b:196



	<i>Festuca bromoides</i> L. = <b>Vulpia bromoides</b>	4b:203
	<i>Festuca distichophylla</i> (Labill.) Hook.f. nom. illeg. = <b>Distichlis distichophylla</b>	
	<i>Festuca dives</i> F.Muell. = <b>Dryopoa dives</b>	4b:199
	<i>Festuca duriuscula</i> L. sensu Benth. (1878) = <b>Festuca plebeia</b> (misapplied in Tasmania)	
	<i>Festuca elatior</i> L. subsp. <i>arundinacea</i> (Schreb.) Hack. = <b>Festuca arundinacea</b>	4b:197
	<i>Festuca hookeriana</i> F.Muell. ex Hook.f. = <b>Hookerchloa hookeriana</b>	4b:223
	<i>Festuca littoralis</i> Labill. = <b>Austrofestuca littoralis</b>	4b:223
i	<b>Festuca nigrescens</b> Lam., Encycl. (Lamarck) 2: 460 (1788)	4b:197
	<i>Festuca ovina</i> L. sensu Rodway (1903) = <b>Festuca plebeia</b>	4b:196
	<i>Festuca pectinata</i> Labill. = <b>Australopyrum pectinatum</b>	
e	<b>Festuca plebeia</b> R.Br., Prodr. Fl. Nov. Holland. 178 (1810)	4b:196
i	<b>Festuca rubra</b> L., Sp. Pl. 1: 74 (1753)	4b:197
	<i>Festuca scabra</i> Labill. nom. illeg., non Vahl = <b>Anthosachne scabra</b>	
	<i>Gastridium lendigerum</i> (L.) Gaudin sensu Rodway (1903) = <b>Gastridium ventricosum</b>	4b:280
i t	<b>Gastridium ventricosum</b> (Gouan) Schinz & Thell., Vierteljahrsschr. Naturf. Ges. Zürich 58: 39 (1913)	4b:280
i	<b>Gaudinia fragilis</b> (L.) P.Beauv., Ess. Agrostogr. 95 (1812)	4b:238
	<b>Glyceria australis</b> C.E.Hubb., Bull. Misc. Inform. Kew. 1934: 450 (1934)	4b:228
i	<b>Glyceria declinata</b> Bréb., Fl. Normandie, ed. 3: 354 (1859)	4b:229
i t	<b>Glyceria fluitans</b> (L.) R.Br., Prodr. Fl. Nov. Holland. 179 (1810)	4b:229
	<i>Glyceria fluitans</i> (L.) R.Br. sensu Rodway (1903) = <b>Glyceria australis</b>	4b:228
i	<b>Glyceria maxima</b> (Hartm.) Holmb., Bot. Not. 72: 97 (1919)	4b:229
	<i>Glyceria notata</i> Chevall. sensu Buchanan (2005) = <b>Glyceria plicata</b>	
n i *	<i>Glyceria plicata</i> (Fr.) Fr., Novitiae Florae Suecicae Mantissa 3: 176 (1845)	4b:229
	<i>Glyceria stricta</i> Hook.f. = <b>Puccinellia stricta</b>	4b:205
i	<b>Hainardia cylindrica</b> (Willd.) Greuter, Boissiera 13: 177 (1967)	4b:227
	<i>Hemarthria compressa</i> (L.f.) R.Br. sensu Benth. (1878) = <b>Hemarthria uncinata</b> (misapplied in Tasmania)	
	<b>Hemarthria uncinata</b> R.Br. var. <i>uncinata</i> , Prodr. Fl. Nov. Holland. 207 (1810)	4b:359
	<i>Hierochloe antarctica</i> (Labill.) R.Br. = <b>Hierochloe redolens</b>	
	<i>Hierochloe borealis</i> Roem. & Schult. sensu Hooker (1860) = <b>Hierochloe fraseri</b> (misapplied in Tasmania)	
e	<b>Hierochloe fraseri</b> Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 93 (1845)	4b:248
	<b>Hierochloe rariflora</b> Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 93 (1845)	4b:249
	<b>Hierochloe redolens</b> (Vahl) R.Br. ex Roem. & Schult., Syst. Veg., ed. 15 bis [Roemer & Schultes] 2: 514 (1817)	4b:247
	<i>Hierochloe redolens</i> (Vahl) R.Br. ex Roem. & Schult. var. <i>fraseri</i> (Hook.f.) Benth. = <b>Hierochloe fraseri</b>	
i	<b>Holcus lanatus</b> L., Sp. Pl. 2: 1048 (1753)	4b:245
n i *	<i>Holcus mollis</i> L., Syst. Nat., ed. 10, 2: 1305 (1759)	4b:245
	<b>Hookerchloa hookeriana</b> (F.Muell. ex Hook.f.) E.B.Alexeev, Byull. Moskovsk. Obshch. Isp. Prir., Otd. Biol. 90: 106 (1985)	4b:223
i	<b>Hordeum distichon</b> L., Sp. Pl. 1: 85 (1753)	4b:297
i	<b>Hordeum glaucum</b> Steud., Syn. Pl. Glumac. 1: 352 (1854)	4b:298
n i	<b>Hordeum hystrix</b> Roth, Catal. Bot. 1: 23 (1797)	4b:300

i	<b>Hordeum leporinum</b> Link, Linnaea 9: 133 (1834)	4b:298
i	<b>Hordeum marinum</b> Huds., Fl. Angl. (Hudson) (ed. 2) 1: 57 (1778)	4b:300
	Hordeum marinum Huds. subsp. gussoneanum (Parl.) Thell. = <b>Hordeum hystrix</b>	4b:300
i	<b>Hordeum murinum</b> L., Sp. Pl. 1: 85 (1753)	4b:298
	Hordeum murinum L. subsp. glaucum (Steud.) Tzvelev = <b>Hordeum glaucum</b>	4b:298
	Hordeum murinum L. subsp. leporinum (Link) Arcang. = <b>Hordeum leporinum</b>	4b:298
	Hordeum nodosum L. sensu Bentham (1878) – a name of uncertain application	
i	<b>Hordeum vulgare</b> L., Sp. Pl. 1: 84 (1753)	4b:297
	Imperata arundinacea Cirillo sensu Rodway (1903) = <b>Imperata cylindrica</b> var. <b>major</b>	4b:353
	<b>Imperata cylindrica</b> (L.) P.Beauv. var. <b>major</b> (Nees) C.E.Hubb., Grasses of Mauritius & Rodriguez: 96 (1940)	4b:353
	<b>Isachne globosa</b> (Thunb. ex Murray) Kuntze, Revis. Gen. Pl. 2: 778 (1891)	
	Koeleria cristata Pers. = <b>Rostraria cristata</b>	4b:241
	Koeleria cristata (L.) Bertol. sensu Rodway (1903) = <b>Koeleria macrantha</b>	4b:240
i x	<b>Koeleria macrantha</b> (Ledeb.) Schult., Mant. 2 (Schultes): 345 (1824)	4b:240
	Koeleria phleoides (Vill.) Pers. sensu Rodway (1903) = <b>Rostraria cristata</b>	4b:241
	<b>Lachnagrostis aemula</b> (R.Br.) Trin., Fund. Agrost. (Trinius): 128 (1820)	4b:258
	Lachnagrostis aequata (Nees) S.W.L.Jacobs = <b>Lachnagrostis rudis</b>	4b:261
	<b>Lachnagrostis billardierei</b> (R.Br.) Trin. subsp. <b>billardierei</b> , Fund. Agrost. (Trinius): 128 (1820)	4b:260
e	<b>Lachnagrostis billardierei</b> (R.Br.) Trin. subsp. <b>tenuiseta</b> (D.I.Morris) S.W.L.Jacobs, Telopea 9: 445 (2001)	4b:261
e	<b>Lachnagrostis collicola</b> (D.I.Morris) S.W.L.Jacobs, Telopea 9: 445 (2001)	4b:261
	<b>Lachnagrostis filiformis</b> (G.Forst.) Trin., Fund. Agrost. (Trinius): 128 (1820)	4b:259
e	<b>Lachnagrostis lacunarum</b> (D.I.Morris) S.W.L.Jacobs, Telopea 9: 446 (2001)	4b:260
e	<b>Lachnagrostis morrisii</b> A.J.Br., Muelleria 24: 127 (2006)	4b:261
	<b>Lachnagrostis punicea</b> (A.J.Br. & N.G.Walsh) S.W.L.Jacobs subsp. <b>filifolia</b> (Vickery) S.W.L.Jacobs, Telopea 10: 840 (2004)	4b:261
	<b>Lachnagrostis punicea</b> (A.J.Br. & N.G.Walsh) S.W.L.Jacobs subsp. <b>punicea</b> , Telopea 9: 837 (2002)	4b:259
	<b>Lachnagrostis robusta</b> (Vickery) S.W.L.Jacobs, Telopea 9: 447 (2001)	4b:261
	<b>Lachnagrostis rudis</b> (Roem. & Schult.) Trin. subsp. <b>nana</b> A.J.Br., Muelleria 33: 93 (2015)	
	<b>Lachnagrostis rudis</b> (Roem. & Schult.) Trin. subsp. <b>rudis</b> , Fund. Agrost. 128 (1820)	
	Lachnagrostis scabra (P.Beauv.) Nees ex Steud. sensu de Salas & Baker 2014 = <b>Lachnagrostis rudis</b>	4b:261
i	<b>Lagurus ovatus</b> L., Sp. Pl. 1: 81 (1753)	4b:281
	Lepturus cylindricus (Willd.) Trin. = <b>Hainardia cylindrica</b>	4b:227
	Lepturus incurvatus Trin. = <b>Parapholis incurva</b>	4b:226
i t	<b>Leymus arenarius</b> (L.) Hochst., Flora 31: 118 (1848)	
i	<b>Lolium loliaceum</b> (Bory & Chaub. ex Fauché) Hand.-Mazz., Ann. K. K. Naturhist. Hofmus. 28: 32 (1914)	4b:202
i	<b>Lolium multiflorum</b> Lam., Fl. Franç. (Lamarck) 3: 621 (1779)	4b:201
i	<b>Lolium perenne</b> L., Sp. Pl. 1: 83 (1753)	4b:201
	Lolium perenne L. var. italicum (A.Braun) Rodway = <b>Lolium multiflorum</b>	
i	<b>Lolium rigidum</b> Gaudin, Agrost. Helv. 1: 334 (1811)	4b:201

i	<b>Lolium temulentum</b> L. f. <b>arvense</b> (With.) Junge, Jahrb. Hamburg. Wiss. Anst. Beih. 30: 314 (1913)	4b:200
	Lophochloa cristata (L.) Hyl. = <b>Rostraria cristata</b>	4b:241
	Lophopyrum elongatum (Host) Á.Löve = <b>Thinopyrum elongatum</b>	4b:295
	Lophopyrum ponticum (Podp.) Á.Löve sensu Buchanan (2005) = <b>Thinopyrum elongatum</b>	4b:295
	Microlaena gunnii Hook.f. = <b>Microlaena stipoides</b>	
	<b>Microlaena stipoides</b> (Labill.) R.Br. var. <b>stipoides</b> , Prodr. Fl. Nov. Holland. 210 (1810)	4b:181
e	<b>Microlaena tasmanica</b> (Hook.f.) Benth. var. <b>subalpina</b> F.Muell. ex Benth., Fl. Austral. 7: 553 (1878)	4b:182
e	<b>Microlaena tasmanica</b> (Hook.f.) Benth. var. <b>tasmanica</b> , Fl. Austral. 7: 552 (1878)	4b:181
n i *	Molineriella minuta (L.) Rouy, Fl. France [Rouy & Foucaud] 14: 102 (1913)	
	Monerma cylindrica (Willd.) Coss. & Durieu = <b>Hainardia cylindrica</b>	4b:227
i t	<b>Nardus stricta</b> L., Sp. Pl. 1: 53 (1753)	4b:184
i	<b>Nassella leucotricha</b> (Trin. & Rupr.) R.W.Pohl, Taxon 39: 610 (1990)	
i	<b>Nassella neesiana</b> (Trin. & Rupr.) Barkworth, Taxon 39: 611 (1990)	
i	<b>Nassella trichotoma</b> (Nees) Hack. ex Arechav., Anales Mus. Nac. Montevideo 1: 366 (1896)	4b:193
	Notodanthonia caespitosa (Gaudich.) Zotov = <b>Rytidosperma caespitosum</b>	4b:312
	Notodanthonia carphoides (F.Muell. ex Benth.) Zotov = <b>Rytidosperma carphoides</b>	4b:306
	Notodanthonia diemenica (D.I.Morris) H.P.Linder = <b>Rytidosperma diemenicum</b>	4b:311
	Notodanthonia geniculata (J.M.Black) Zotov = <b>Rytidosperma geniculatum</b>	4b:308
	Notodanthonia gracilis (Hook.f.) Zotov = <b>Rytidosperma gracile</b>	4b:309
	Notodanthonia laevis (Vickery) Zotov = <b>Rytidosperma laeve</b>	4b:312
	Notodanthonia penicillata (Labill.) Zotov = <b>Rytidosperma penicillatum</b>	4b:313
	Notodanthonia pilosa (R.Br.) Zotov = <b>Rytidosperma pilosum</b>	4b:313
	Notodanthonia popinensis (D.I.Morris) H.P.Linder = <b>Rytidosperma fulvum</b>	4b:316
	Notodanthonia racemosa (R.Br.) Zotov = <b>Rytidosperma racemosum</b>	4b:314
	Notodanthonia remota (D.I.Morris) H.P.Linder = <b>Rytidosperma remotum</b>	4b:316
	Notodanthonia semiannularis (Labill.) Zotov = <b>Rytidosperma semiannulare</b>	4b:308
	Notodanthonia setacea (R.Br.) Veldkamp = <b>Rytidosperma setaceum</b>	4b:311
	Notodanthonia tenuior (Steud.) Conert = <b>Rytidosperma tenuius</b>	4b:310
	Oryzopsis miliacea (L.) Benth. & Hook.f. ex Asch. & Schweinf. = <b>Piptatherum miliaceum</b>	4b:194
n	Panicum capillare L. – previously listed as naturalised but insufficient evidence exists to support this	4b:335
n	Panicum capillare L. var. occidentale Rydb. = Panicum capillare	4b:335
n i *	Panicum gilvum Launert, Mit. Bot. Staatssamml. München 8: 153 (1970)	4b:335
	Panicum gracile R.Br. sensu Rodway (1903) = Paspalidium gracile	
i	<b>Panicum hillmanii</b> Chase, J. Wash. Acad. Sci. 14: 345 (1934)	4b:334
i	<b>Panicum miliaceum</b> L., Sp. Pl. 1: 58 (1753)	4b:333
	Panicum sanguinale L. = <b>Digitaria sanguinalis</b>	4b:346
	Panicum schinzii Hack. sensu Curtis & Morris (1994) = <b>Panicum gilvum</b>	4b:335
i	<b>Parapholis incurva</b> (L.) C.E.Hubb., Blumea Suppl. 3: 14 (1946)	4b:226
i	<b>Parapholis strigosa</b> (Dumort.) C.E.Hubb., Blumea suppl. 3: 14 (1946)	4b:226
	Paspalidium gracile (R.Br.) Hughes sensu Townrow (1969) identity uncertain	
i	<b>Paspalum dilatatum</b> Poir., Encycl. (Lamarck) 5: 35 (1804)	4b:340
i	<b>Paspalum distichum</b> L., Syst. Nat., ed. 10. 2: 855 (1759)	4b:340
i	<b>Paspalum urvillei</b> Steud., Syn. Pl. Glumac. 1: 24 (1853)	4b:340

	Pennisetum alopecuroides (L.) Spreng. = <i>Cenchrus purpurascens</i>	
	Pennisetum clandestinum Hochst. ex Chiov. = <b>Cenchrus clandestinus</b>	4b:348
	Pennisetum macrourum Trin. = <b>Cenchrus macrourus</b>	4b:351
	Pennisetum villosum R.Br. ex Fresen. = <b>Cenchrus longisetus</b>	4b:349
	Pentapogon billardierei R.Br. sensu Rodway (1903) = <b>Pentapogon quadrifidus</b>	4b:275
	Pentapogon billardierei R.Br. var. parviflorus Benth. = <b>Pentapogon quadrifidus</b> var. <b>parviflorus</b>	
e	<b>Pentapogon quadrifidus</b> (Labill.) Baill. var. <b>parviflorus</b> (Benth.) D.I.Morris, Muelleria 7: 167 (1990)	4b:276
	<b>Pentapogon quadrifidus</b> (Labill.) Baill. var. <b>quadrifidus</b> , Hist. Pl. (Baillon) 12: 280 (1893)	4b:275
i	<b>Phalaris aquatica</b> L., Cent. Pl. 1. 4 (1755)	4b:252
i	<b>Phalaris arundinacea</b> L. var. <b>arundinacea</b> , Sp. Pl. 1: 55 (1753)	4b:251
i	<b>Phalaris canariensis</b> L., Sp. Pl. 1: 54 (1753)	4b:251
i	<b>Phalaris minor</b> Retz., Observ. Bot. (Retzius) 3: 8 (1783)	4b:252
i	<b>Phalaris paradoxa</b> L., Sp. Pl., ed. 2. 2: 1665 (1763)	4b:251
	<i>Phalaris tuberosa</i> L. = <b>Phalaris aquatica</b>	4b:252
i	<b>Phleum pratense</b> L. subsp. <b>pratense</b> , Sp. Pl. 1: 59 (1753)	4b:286
	<b>Phragmites australis</b> (Cav.) Trin. ex Steud., Nomencl. Bot. [Steudel], ed. 2. 2: 324 (1841)	4b:322
	<i>Phragmites communis</i> Trin. sensu Rodway (1903) = <b>Phragmites australis</b>	4b:322
i	<b>Piptatherum miliaceum</b> (L.) Coss., Notes sur quelques Plantes Critiques, Rares ou Nouvelles: 129 (1851)	4b:194
	<i>Poa affinis</i> R.Br. sensu Hooker (1860) – a name of uncertain application	
i	<b>Poa annua</b> L., Sp. Pl. 1: 68 (1753)	4b:211
	<i>Poa australis</i> R.Br. = <b>Poa poiformis</b>	
	<i>Poa billardierei</i> (Spreng.) St.-Yves sensu Rodway (1903) = <b>Poa poiformis</b> var. <b>poiformis</b>	4b:220
i	<b>Poa bulbosa</b> L., Sp. Pl. 1: 70 (1753)	4b:213
	<i>Poa caespitosa</i> G.Forst. ex Spreng. var. <i>alpina</i> F.Muell. ex Benth. sensu Rodway (1903) = <b>Poa gunnii</b>	4b:218
	<i>Poa caespitosa</i> G.Forst. ex Spreng. var. <i>australis</i> (R.Br.) Benth. sensu Rodway (1903) = <b>Poa rodwayi</b>	4b:215
	<i>Poa caespitosa</i> G.Forst. ex Spreng. var. <i>laevis</i> (R.Br.) Benth. = <b>Poa poiformis</b>	
	<i>Poa caespitosa</i> G.Forst. ex Spreng. var. <i>tenera</i> (F.Muell.) Benth. sensu Rodway (1903) = <b>Poa tenera</b>	4b:217
	<b>Poa clelandii</b> Vickery, Contr. New South Wales Natl. Herb. 4: 193 (1970)	4b:221
	<b>Poa clivicola</b> Vickery, Contr. New South Wales Natl. Herb. 4: 213 (1970)	
i	<b>Poa compressa</b> L., Sp. Pl. 1: 69 (1753)	4b:214
	<b>Poa costiniana</b> Vickery, Contr. New South Wales Natl. Herb. 4: 214 (1970)	4b:218
	<i>Poa distichophylla</i> (Labill.) R.Br. = <b>Distichlis distichophylla</b>	
	<b>Poa fawcettiae</b> Vickery, Contr. New South Wales Natl. Herb. 4: 232 (1970)	4b:217
	<b>Poa gunnii</b> Vickery, Contr. New South Wales Natl. Herb. 4: 217 (1970)	4b:218
	<b>Poa halmaturina</b> J.M.Black, Trans. Roy. Soc. South Australia 66: 248 (1943)	4b:220
	<b>Poa hiemata</b> Vickery, Contr. New South Wales Natl. Herb. 4: 230 (1970)	4b:217
	<b>Poa hookeri</b> Vickery, Contr. New South Wales Natl. Herb. 4: 222 (1970)	4b:216
i	<b>Poa infirma</b> Kunth, Bonpl. & Kunth, Nov. Gen. Sp. [H.B.K.] 1: 158 (1816)	4b:213
e	<b>Poa jugicola</b> D.I.Morris, Muelleria 7: 167 (1990)	4b:219

	<b>Poa labillardierei</b> Steud. var. <b>acris</b> Vickery, Contr. New South Wales Natl. Herb. 4: 205 (1970)	4b:220
	<b>Poa labillardierei</b> Steud. var. <b>labillardierei</b> , Syn. Pl. Glumac. 1: 262 (1854)	4b:219
	<i>Poa laevis</i> R.Br. = <b>Poa poiformis</b>	
e	<b>Poa mollis</b> Vickery, Contr. New South Wales Natl. Herb. 4: 241 (1970)	4b:215
	<b>Poa poiformis</b> (Labill.) Druce var. <b>poiformis</b> , Rep. Bot. Soc. Exch. Club Brit. Isles 1916, Suppl. 2: 640 (1917)	4b:220
	<b>Poa poiformis</b> (Labill.) Druce var. <b>ramifer</b> D.I.Morris, Muelleria 7: 169 (1990)	4b:220
i	<b>Poa pratensis</b> L., Sp. Pl. 1: 67 (1753)	4b:214
	<i>Poa pubinervis</i> (Vickery) S.W.L.Jacobs sensu Baker & Duretto (2011) = <b>Austrofestuca littoralis</b>	
	<i>Poa rigida</i> L. = <b>Catapodium rigidum</b>	4b:224
	<b>Poa rodwayi</b> Vickery, Contr. New South Wales Natl. Herb. 4: 235 (1970)	4b:215
	<i>Poa saxicola</i> R.Br. = <b>Saxipoa saxicola</b>	4b:214
	<b>Poa sieberiana</b> Spreng. var. <b>hirtella</b> Vickery, Contr. New South Wales Natl. Herb. 4: 228 (1970)	
	<b>Poa sieberiana</b> Spreng. var. <b>sieberiana</b> , Syst. Veg. (ed. 16) [Sprengel] 4(2, Cur. Post.): 35 (1827)	4b:216
	<b>Poa tenera</b> F.Muell. ex Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 2: 124, t.164a (1858)	4b:217
i	<b>Poa trivialis</b> L., Sp. Pl. 1: 67 (1753)	4b:213
	<i>Polypogon littoralis</i> Sm. = <b>Polypogon lutosus</b>	4b:283
i	<b>Polypogon lutosus</b> (Poir.) Hitchc., U.S.D.A. Bull. (1915–23) 772: 138 (1920)	4b:283
i	<b>Polypogon maritimus</b> Willd. var. <b>subspatheaceus</b> (Req.) Parl., Fl. Ital. (Parlatore) 1: 200 (1850)	4b:282
i	<b>Polypogon monspeliensis</b> (L.) Desf., Fl. Atlant. 1: 67 (1798)	4b:282
	<i>Polypogon viridis</i> (Gouan) Breistr. = <b>Agrostis viridis</b>	4b:283
	<i>Psamma arenaria</i> (L.) P.Beauv. = <b>Ammophila arenaria</b>	4b:274
i	<b>Puccinellia distans</b> (Jacq.) Parl., Fl. Ital. (Parlatore) 1: 366 (1850)	4b:205
e	<b>Puccinellia harcusiana</b> A.R.Williams, Fl. Australia 44A: 386 (2009)	
	<b>Puccinellia perlaxa</b> (Stapf ex N.G.Walsh) N.G.Walsh & A.R.Williams, Nuytsia 16: 464 (2007)	4b:205
	<b>Puccinellia stricta</b> (Hook.f.) C.H.Blom, Acta Horti Gothob. 5: 89 (1930)	4b:205
	<i>Puccinellia stricta</i> (Hook.f.) C.H.Blom var. <i>perlaxa</i> Stapf ex N.G.Walsh = <b>Puccinellia perlaxa</b>	4b:205
i	<b>Rostraria cristata</b> (L.) Tzvelev, Novosti Sist. Vyssh. Rast. 7: 47 (1971)	4b:241
	<b>Rytidosperma australe</b> (Petrie) Clayton & Renvoize ex Connor & Edgar, New Zealand J. Bot. 25: 166 (1987)	4b:318
	<b>Rytidosperma caespitosum</b> (Gaudich.) Connor & Edgar, New Zealand J. Bot. 17: 325 (1979)	4b:312
	<b>Rytidosperma carphoides</b> (F.Muell. ex Benth.) Connor & Edgar, New Zealand J. Bot. 17: 331 (1979)	4b:306
e	<b>Rytidosperma diemenicum</b> (D.I.Morris) A.M.Humphreys & H.P.Linder, Ann. Missouri Bot. Gard. 97: 358 (2010)	4b:311
	<b>Rytidosperma dimidiatum</b> (Vickery) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979)	4b:317
e	<b>Rytidosperma fortuneae-hibernae</b> (Renvoize) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979)	4b:310
?i	<b>Rytidosperma fulvum</b> (Vickery) A.M.Humphreys & H.P.Linder, Ann. Missouri Bot. Gard. 97: 358 (2010)	
	<b>Rytidosperma geniculatum</b> (J.M.Black) Connor & Edgar, New Zealand J. Bot. 17: 323 (1979)	4b:308

t	<b>Rytidosperma gracile</b> (Hook.f.) Connor & Edgar, New Zealand J. Bot. 17: 330 (1979)	4b:309
	<b>Rytidosperma indutum</b> (Vickery) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979)	4b:312
	<b>Rytidosperma laeve</b> (Vickery) Connor & Edgar, New Zealand J. Bot. 17: 325 (1979)	4b:312
e	<b>Rytidosperma nitens</b> (D.I.Morris) H.P.Linder, Telopea 6: 614 (1996)	4b:315
	<b>Rytidosperma nivicola</b> (Vickery) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979)	4b:315
	<b>Rytidosperma nudiflorum</b> (P.Morris) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979)	4b:314
	<b>Rytidosperma oreophilum</b> H.P.Linder & N.G.Walsh, Muelleria 8: 283 (1995)	
	<i>Rytidosperma pallidum</i> (R.Br.) A.M.Humphreys & H.P.Linder recorded in error	
e	<b>Rytidosperma pauciflorum</b> (R.Br.) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979)	4b:309
	<b>Rytidosperma penicillatum</b> (Labill.) Connor & Edgar, New Zealand J. Bot. 17: 327 (1979)	4b:313
	<b>Rytidosperma pilosum</b> (R.Br.) Connor & Edgar, New Zealand J. Bot. 17: 326 (1979)	4b:313
	<i>Rytidosperma popinensis</i> (D.I.Morris) A.M.Humphreys & H.P.Linder = <b>Rytidosperma fulvum</b>	4b:316
	<i>Rytidosperma procerum</i> (Vickery) Connor & Edgar = <b>Rytidosperma indutum</b>	4b:312
	<b>Rytidosperma racemosum</b> (R.Br.) Connor & Edgar var. <b>racemosum</b> , New Zealand J. Bot. 17: 327 (1979)	4b:314
e	<b>Rytidosperma remotum</b> (D.I.Morris) A.M.Humphreys & H.P.Linder, Ann. Missouri Bot. Gard. 97: 359 (2010)	4b:316
	<b>Rytidosperma semiannulare</b> (Labill.) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979)	4b:308
	<b>Rytidosperma setaceum</b> (R.Br.) Connor & Edgar, New Zealand J. Bot. 17: 332 (1979)	4b:311
	<b>Rytidosperma tenuius</b> (Steud.) A.Hansen & Sunding, Fl. Macaronesia, ed. 2, 1: 93 (1979)	4b:310
	<b>Saxipoa saxicola</b> (R.Br.) Soreng, L.J.Gillespie & S.W.L.Jacobs, Austral. Syst. Bot. 22: 407 (2010)	4b:214
	<i>Schedonorus hookerianus</i> (F.Muell. ex Hook.f.) P.Beauv. = <b>Hookerochloa hookeriana</b>	4b:223
	<i>Schedonorus littoralis</i> (Labill.) P.Beauv. = <b>Austrofestuca littoralis</b>	4b:223
	<i>Schedonorus phoenix</i> (Scop.) Holub = <b>Festuca arundinacea</b>	4b:197
i	<b>Sclerochloa dura</b> (L.) P.Beauv., Ess. Agrostogr. 98, 177, t.19 fig.4 (1812)	4b:225
i	<b>Secale cereale</b> L., Sp. Pl. 1: 84 (1753)	4b:302
	<i>Setaria geniculata</i> P.Beauv. var. <i>pauciseta</i> Desv. = <b>Setaria parviflora</b>	4b:344
	<i>Setaria gracilis</i> Kunth var. <i>pauciseta</i> (Desv.) B.K.Simon = <b>Setaria parviflora</b>	4b:344
i	<b>Setaria italica</b> (L.) P.Beauv., Ess. Agrostogr. 51, 170, 178 (1812)	4b:343
i	<b>Setaria parviflora</b> (Poir.) Kerguélen, Lejeunia 120: 161 (1987)	4b:344
n	<i>Setaria pumila</i> (Poir.) Roem. & Schult. subsp. <i>pumila</i> – previously listed as naturalised but insufficient evidence exists to support this	
i	<b>Setaria verticillata</b> (L.) P.Beauv., Ess. Agrostogr. 51, 171, 178 (1812)	4b:342
i	<b>Setaria viridis</b> (L.) P.Beauv., Ess. Agrostogr. 51, 171, 178 (1812)	4b:343
	<i>Sieglingia decumbens</i> (L.) Bernh. = <b>Danthonia decumbens</b>	4b:317
n	<i>Sorghum bicolor</i> (L.) Moench – previously listed as naturalised but insufficient evidence exists to support this	
n i *	<i>Sorghum halepense</i> (L.) Pers., Syn. Pl. (Persoon) 1: 101 (1805)	
i	<b>Spartina anglica</b> C.E.Hubb., Bot. J. Linn. Soc. 76: 364 (1978)	4b:330
	<i>Spartina townsendii</i> H.Groves & J.Groves sensu Townrow (1969) = <b>Spartina anglica</b>	4b:330
	<i>Spinifex hirsutus</i> Labill. sensu Rodway (1903) = <b>Spinifex sericeus</b>	4b:351
	<b>Spinifex sericeus</b> R.Br., Prodr. Fl. Nov. Holland. 198 (1810)	4b:351
i	<b>Sporobolus africanus</b> (Poir.) Robyns & Tournay, Bull. Jard. Bot. État Bruxelles 25: 242 (1955)	4b:328
	<i>Sporobolus capensis</i> Kunth = <b>Sporobolus africanus</b>	4b:328
	<i>Sporobolus indicus</i> (L.) R.Br. = <b>Sporobolus africanus</b>	4b:328

	<b>Sporobolus virginicus</b> (L.) Kunth, Révis. Gramin. 1: 67 (1829)	4b:328
	Stenotaphrum americanum Schrank = <b>Stenotaphrum secundatum</b>	4b:344
i	<b>Stenotaphrum secundatum</b> (Walter) Kuntze, Revis. Gen. Pl. 2: 794 (1891)	4b:344
	Stipa aphanoneura Hughes = <b>Austrostipa flavescens</b>	4b:190
	Stipa aphylla (Rodway) J.Townrow = <b>Austrostipa aphylla</b>	4b:192
	Stipa bigeniculata Hughes = <b>Austrostipa bigeniculata</b>	4b:190
	Stipa blackii C.E.Hubb. = <b>Austrostipa blackii</b>	4b:191
	Stipa caudata Trin. = <b>Amelichloa caudata</b>	4b:192
	Stipa compacta Hughes = <b>Austrostipa flavescens</b>	4b:190
	Stipa elatior (Benth.) Hughes = <b>Austrostipa flavescens</b>	4b:190
	Stipa elegantissima Labill. sensu Labillardiere (1805) and Bentham (1878) recorded in error	
	Stipa eremophila Reader sensu Townrow (1969) identity uncertain	
	Stipa falcata Hughes sensu Townrow (1978) = <b>Austrostipa scabra</b> p.p.maj.	4b:188
	Stipa flavescens Labill. = <b>Austrostipa flavescens</b>	4b:190
	Stipa mollis R.Br. = <b>Austrostipa mollis</b>	4b:189
	Stipa neesiana Trin. & Rupr. = <b>Nassella neesiana</b>	
	Stipa nervosa Vickery var. <i>neuralis</i> sensu Townrow (1978) = <b>Austrostipa rudis</b> subsp.	
	<b>australis</b>	4b:186
	Stipa nodosa S.T.Blake = <b>Austrostipa nodosa</b>	4b:188
	Stipa pubescens R.Br. sensu Rodway (1903) = <b>Austrostipa pubinodis</b>	4b:192
	Stipa pubescens R.Br. var. <i>aphylla</i> Rodway = <b>Austrostipa aphylla</b>	4b:187
	Stipa pubinodis Trin. & Rupr. = <b>Austrostipa pubinodis</b>	4b:187
	Stipa rudis Spreng. = <b>Austrostipa rudis</b>	4b:186
	Stipa scabra Lindl. = <b>Austrostipa scabra</b>	4b:188
	Stipa semibarbata R.Br. = <b>Austrostipa semibarbata</b>	4b:189
	Stipa setacea R.Br. sensu Rodway (1903) = <b>Austrostipa stuposa</b>	4b:191
	Stipa stipoides (Hook.f.) Veldkamp = <b>Austrostipa stipoides</b>	4b:186
	Stipa stuposa Hughes = <b>Austrostipa stuposa</b>	4b:191
	Stipa teretifolia Steud. = <b>Austrostipa stipoides</b>	4b:186
	Stipa variabilis Hughes sensu Townrow (1969) = <b>Austrostipa nodosa</b>	4b:188
	<b>Tetrarrhena acuminata</b> R.Br., Prodr. Fl. Nov. Holland. 210 (1810)	4b:182
	<b>Tetrarrhena distichophylla</b> (Labill.) R.Br., Prodr. Fl. Nov. Holland. 210 (1810)	4b:183
	<b>Tetrarrhena juncea</b> R.Br., Prodr. Fl. Nov. Holland. 210 (1810)	4b:183
e	<b>Tetrarrhena oreophila</b> D.I.Morris var. <i>minor</i> D.I.Morris, Records of the Queen Victoria Museum, n.ser., 55: 4 (1977)	4b:182
e	<b>Tetrarrhena oreophila</b> D.I.Morris var. <i>oreophila</i> , Records of the Queen Victoria Museum, n.ser., 55: 4 (1977)	4b:182
	Tetrarrhena tenacissima Nees sensu Hooker (1860) = <b>Tetrarrhena juncea</b>	
	Themeda australis (R.Br.) Stapf = <b>Themeda triandra</b>	4b:358
	<b>Themeda triandra</b> Forssk., Fl. Aegypt.-Arab. 178 (1775)	4b:358
i	<b>Thinopyrum elongatum</b> (Host) D.R.Dewey, Gene Manipulation Pl. Improv. 274 (1984)	4b:295
i	<b>Thinopyrum junceiforme</b> (Á.Löve & D.Löve) Á.Löve, Taxon 29: 351 (1980)	4b:293
	Thonandia gracilis (Hook.f.) H.P.Linder = <b>Rytidosperma gracile</b>	4b:309
	Thonandia semiannularis (Labill.) H.P.Linder = <b>Rytidosperma semiannulare</b>	4b:308

- Trisetum spicatum** (L.) K.Richt. subsp. **australiense** Hultén ex Veldkamp, Gard. Bull. Singapore 36: 135 (1983) 4b:239
- Trisetum subspicatum (L.) P.Beauv. sensu Rodway (1903) = **Trisetum spicatum** subsp. **australiense** 4b:239
- i **Triticum aestivum** L., Sp. Pl. 1: 85 (1753) 4b:303
- Triticum pectinatum (Labill.) R.Br. = **Australopyrum pectinatum**
- Triticum scabrum R.Br. = **Anthosachne scabra**
- Triticum velutinum (Nees) Hook.f. nom. illeg., nom. superfl. = **Australopyrum velutinum**
- Uniola distichophylla Labill. = **Distichlis distichophylla**
- i **Vulpia bromoides** (L.) Gray, Nat. Arr. Brit. Pl. 2: 124 (1821) 4b:203
- i **Vulpia fasciculata** (Forssk.) Fritsch, Exkursionsfl. Oesterreich (Ed. 2): 74 (1909) 4b:204
- Vulpia megalura (Nutt.) Rydb. = **Vulpia myuros** f. **megalura** 4b:204
- i **Vulpia muralis** (Kunth) Nees, Linnaea 19: 694 (1847)
- i **Vulpia myuros** (L.) C.C.Gmel. f. **megalura** (Nutt.) Stace & R.Cotton, Watsonia 11: 72 (1976) 4b:204
- i **Vulpia myuros** (L.) C.C.Gmel. f. **myuros**, Fl. Bad. 1: 8 (1805)
- Zoysia macrantha** Desv. subsp. **walshii** Night., Fl. Australia 43: 376 (2002) 4b:332
- Zoysia pungens Willd. sensu Rodway (1903) = **Zoysia macrantha** subsp. **walshii** 4b:332
- ×Agropogon littoralis (Sm.) C.E.Hubb. = **Polypogon lutosus** 4b:283

#### POSIDONIACEAE

- Caulinia antarctica R.Br. = **Amphibolis antarctica** (Cymodoceaceae)
- Caulinia oceanica R.Br. = **Potamogeton australiensis** (Potamogetonaceae)
- Posidonia australis** Hook.f., Bot. Antarct. Voy. Ill. (Fl. Tasman.) 2: 43 (1858) 4b:25

#### POTAMOGETONACEAE

- Potamogeton australiensis** A.Benn., J. Bot. 48: 149 (1910) 4b:19
- Potamogeton cheesemanii** A.Benn., J. Bot. 66: 66 (1883) 4b:19
- Potamogeton crispus** L., Sp. Pl. 1: 126 (1753) 4b:18
- Potamogeton drummondii** Benth., Fl. Austral. 7: 171 (1878)
- Potamogeton gramineus L. sensu Hooker (1860) = **Potamogeton ochreatus** identity somewhat uncertain
- Potamogeton heterophyllus Schreb. sensu Hooker (1860) = **Potamogeton perfoliatus** identity somewhat uncertain
- Potamogeton natans L. sensu Rodway (1903) misapplied to **P. cheesemanii**, **P. australiensis** and **P. drummondii** 4b:19
- Potamogeton obtusifolius Mert. & W.D.J.Koch sensu Rodway (1903) = **Potamogeton crispus** 4b:18
- Potamogeton ochreatus** Raoul, Ann. Sci. Nat., Bot., sér. 3, 2: 117 (1844) 4b:18
- Potamogeton pectinatus L. = **Stuckenia pectinata** 4b:18
- Potamogeton perfoliatus** L., Sp. Pl. 1: 126 (1753) 4b:18
- Potamogeton praelongus Wulfen sensu Rodway (1903) = **Potamogeton ochreatus** 4b:18
- Potamogeton tricarinatus F.Muell. & A.Benn. ex A.Benn. sensu Baker & de Salas (2013) misapplied to **P. australiensis**, **P. cheesemanii** and **P. drummondii** 4b:19
- Stuckenia pectinata** (L.) Börner, Fl. Deut. Volk. 713 (1912) 4b:18

#### RESTIONACEAE

- Acion hookeri (D.I.Morris) B.G.Briggs & L.A.S.Johnson = **Chordifex hookeri** 4b:46
- Acion monocephalum (R.Br.) B.G.Briggs & L.A.S.Johnson = **Chordifex monocephalus** 4b:47



	<b>Apodasmia brownii</b> (Hook.f.) B.G.Briggs & L.A.S.Johnson, <i>Telopea</i> 7: 371 (1998)	4b:39
	<b>Baloskion australe</b> (R.Br.) B.G.Briggs & L.A.S.Johnson, <i>Telopea</i> 8: 23 (1998)	4b:46
	<b>Baloskion tetraphyllum</b> (Labill.) B.G.Briggs & L.A.S.Johnson subsp. <b>tetraphyllum</b> , <i>Telopea</i> 8: 23 (1998)	4b:45
	<i>Calorophus ater</i> L.A.S.Johnson & B.G.Briggs = <b>Calorophus erostris</b>	4b:42
	<b>Calorophus elongatus</b> Labill., <i>Nov. Holl. Pl.</i> 2: 78, t.228 (1806)	4b:41
e	<b>Calorophus erostris</b> (C.B.Clarke) L.A.S.Johnson & B.G.Briggs, <i>The Student's Flora of Tasmania</i> 4b: 425 (1994)	4b:42
	<i>Calorophus lateriflorus</i> (R.Br.) F.Muell. = <b>Empodisma minus</b>	4b:43
e	<b>Chordifex hookeri</b> (D.I.Morris) B.G.Briggs, <i>Telopea</i> 10: 685 (2004)	4b:46
e	<b>Chordifex monocephalus</b> (R.Br.) B.G.Briggs, <i>Telopea</i> 10: 685 (2004)	4b:47
	<b>Empodisma minus</b> (Hook.f.) L.A.S.Johnson & D.F.Cutler, <i>Kew Bull.</i> 28: 383 (1974)	4b:43
	<b>Eurychorda complanata</b> (R.Br.) B.G.Briggs & L.A.S.Johnson, <i>Telopea</i> 7: 359 (1998)	4b:46
	<i>Hypolaena exsulca</i> R.Br. recorded in error	
	<b>Hypolaena fastigiata</b> R.Br., <i>Prodr. Fl. Nov. Holland.</i> 251 (1810)	4b:40
	<i>Hypolaena lateriflora</i> (R.Br.) Benth. = <b>Empodisma minus</b>	4b:43
	<i>Hypolaena longissima</i> Benth. sensu Rodway (1903) = <b>Calorophus elongatus</b>	4b:41
	<i>Leptocarpus brownii</i> Hook.f. = <b>Apodasmia brownii</b>	4b:39
	<b>Leptocarpus tenax</b> (Labill.) R.Br., <i>Prodr. Fl. Nov. Holland.</i> 250 (1810)	4b:39
	<b>Lepyrodia muelleri</b> Benth., <i>Fl. Austral.</i> 7: 215 (1878)	4b:43
	<i>Lepyrodia tasmanica</i> Hook.f. = <b>Sporadanthus tasmanicus</b>	4b:44
	<i>Restio australis</i> R.Br. = <b>Baloskion australe</b>	4b:46
	<i>Restio complanatus</i> R.Br. = <b>Eurychorda complanata</b>	4b:46
	<i>Restio glaber</i> (Rodway) L.A.S.Johnson & B.G.Briggs = <b>Chordifex monocephalus</b>	4b:47
	<i>Restio gracilis</i> R.Br. sensu Rodway (1903) = <b>Baloskion australe</b>	4b:46
	<i>Restio hookeri</i> D.I.Morris = <b>Chordifex hookeri</b>	4b:46
	<i>Restio lateriflorus</i> R.Br. nom. illeg. = <b>Calorophus elongatus</b>	
	<i>Restio monocephalus</i> R.Br. = <b>Chordifex monocephalus</b>	4b:47
	<i>Restio oligocephalus</i> F.Muell. = <b>Chordifex monocephalus</b>	4b:46
	<i>Restio oligocephalus</i> F.Muell. var. <i>glabrum</i> Rodway = <b>Chordifex monocephalus</b>	
	<i>Restio oligocephalus</i> F.Muell. var. <i>intermedius</i> Rodway = <b>Chordifex hookeri</b>	
	<i>Restio tetraphyllum</i> Labill. = <b>Baloskion tetraphyllum</b>	4b:45
	<i>Schoenodum tenax</i> Labill. = <b>Leptocarpus tenax</b>	
	<b>Sporadanthus tasmanicus</b> (Hook.f.) B.G.Briggs & L.A.S.Johnson, <i>Telopea</i> 8: 31 (1998)	4b:44
e	<b>Winifredia sola</b> L.A.S.Johnson & B.G.Briggs, <i>Telopea</i> 2: 738 (1986)	4b:47
<b>RUPPIACEAE</b>		
	<i>Ruppia maritima</i> L. sensu Rodway (1903) = <b>Ruppia polycarpa</b> (misapplied in Tasmania)	4b:21
	<b>Ruppia megacarpa</b> R.Mason, <i>New Zealand J. Bot.</i> 5: 525 (1967)	4b:21
	<b>Ruppia polycarpa</b> R.Mason, <i>New Zealand J. Bot.</i> 5: 524 (1967)	4b:21
	<b>Ruppia tuberosa</b> J.S.Davis & Toml., <i>J. Arnold Arbor.</i> 55: 60 (1974)	4b:21
<b>TYPHACEAE</b>		
	<i>Typha angustifolia</i> L. sensu Rodway (1903) = <b>Typha domingensis</b> & <b>T. orientalis</b>	
	<b>Typha domingensis</b> Pers., <i>Syn. Pl. (Persoon)</i> 2: 532 (1807)	4b:364
i	<b>Typha latifolia</b> L., <i>Sp. Pl.</i> 2: 971 (1753)	4b:363
	<b>Typha orientalis</b> C.Presl, <i>Epimel. Bot.</i> 239 (1851)	4b:363

### XANTHORRHOACEAE

- Lomandra glauca (R.Br.) Ewart sensu Willis (1973) = **Lomandra nana** 4b:422  
**Lomandra longifolia** Labill., Nov. Holl. Pl. 1: 92, t.119 (1805) 4b:422  
**Lomandra nana** (A.T.Lee) A.T.Lee, Fl. Australia 46: 223 (1986) 4b:422  
e **Xanthorrhoea arenaria** D.J.Bedford, Fl. Australia 46: 225 (1986) 4b:421  
**Xanthorrhoea australis** R.Br., Prodr. Fl. Nov. Holland. 288 (1810) 4b:419  
e **Xanthorrhoea bracteata** R.Br., Prodr. Fl. Nov. Holland. 288 (1810) 4b:421  
Xanthorrhoea hastilis R.Br. sensu Hooker (1860) = **Xanthorrhoea australis** (misapplied in Tasmania)  
Xanthorrhoea minor R.Br. sensu Rodway (1903) = **Xanthorrhoea arenaria** & **X. bracteata** 4b:421  
Xerotes glauca R.Br. sensu Rodway (1903) = **Lomandra nana** 4b:422  
Xerotes longifolia (Labill.) R.Br. = **Lomandra longifolia** 4b:422

### XYRIDACEAE

- Xyris gracilis R.Br. sensu Bentham (1878), misapplied to **X. marginata**, **X. muelleri**, and **X. tasmanica**  
Xyris gracilis R.Br. subsp. tasmanica D.I.Morris = **Xyris tasmanica** 4b:35  
Xyris gracilis R.Br. var. bracteata Benth. = **Xyris muelleri**  
e **Xyris marginata** Rendle, J. Bot. 37: 503 (1899) 4b:35  
e **Xyris muelleri** Malme, Svensk. Bot. Tidskr. 21: 381 (1928) 4b:35  
**Xyris operculata** Labill., Nov. Holl. Pl. 1: 14, t.10 (1805) 4b:34  
e **Xyris tasmanica** (D.I.Morris) Doust & B.J.Conn, Austral. Syst. Bot. 10(2): 212 (1997) 4b:35

### ZANNICHELLIACEAE

- x **Lepilaena australis** J.Drumm. ex Harv., Hooker's J. Bot. Kew Gard. Misc. 7: 58 (1855) 4b:22  
**Lepilaena bilocularis** Kirk, Trans. & Proc. New Zealand Inst. 28: 500 (1896) 4b:24  
**Lepilaena cylindrocarpa** (Körn. ex Müll.Stuttg) Benth., Fl. Austral. 7: 180 (1878) 4b:22  
**Lepilaena marina** E.L.Robertson, Marine Benthic Fl. South. Australia 1: 80 (1984) 4b:23  
**Lepilaena patentifolia** E.L.Robertson, Fl. S. Australia 4: 1736 (1986) 4b:23  
**Lepilaena preissii** (Lehm.) F.Muell., Fragm. (Mueller) 8: 217 (1874) 4b:24  
Zannichellia palustris L. sensu Hooker (1860) probably misapplied to **Lepilaena** spp.

### ZOSTERACEAE

- Heterozostera nigricaulis** J.Kuo, Aquatic Bot. 81(2): 110 (2005)  
**Heterozostera tasmanica** (G.Martens ex Asch.) Hartog, Verh. Kon. Ned. Akad. Wetensch., Afd. Natuurk., Sect. 2. 59: 116 (1970) 4b:28  
Nanozostera muelleri (Irmisch ex Asch.) Toml. & Posl. = **Zostera muelleri** subsp. **muelleri** 4b:29  
Zostera marina R.Br. = **Heterozostera nigricaulis**  
**Zostera muelleri** Irmisch ex Asch. subsp. **muelleri**, Linnaea 35: 168 (1868) 4b:29  
Zostera nana Mert. ex Roth sensu Rodway (1903) = **Zostera muelleri** subsp. **muelleri** 4b:29  
Zostera tasmanica G.Martens ex Asch. = **Heterozostera tasmanica** 4b:28

## GYMNOSPERMAE

### CUPRESSACEAE

- e **Athrotaxis cupressoides** D.Don, Ann. Nat. Hist. 1: 234 (1838) 1:6

- e **Athrotaxis × laxifolia** Hook., Icon. Pl. 6: t.573 (1843) = **A. cupressoides** × **A. selaginoides** 1:6
- e **Athrotaxis selaginoides** D.Don, Ann. Nat. Hist. 1: 234 (1838) 1:7
- Athrotaxis tetragona Hook. = **Microcachrys tetragona** (Podocarpaceae)
- Callitris cupressiformis F.Muell. var. tasmanica (Benth.) Maiden = **Callitris rhomboidea**
- e **Callitris oblonga** Rich. & A.Rich. subsp. **oblonga**, Comm. Bot. Conif. Cycad. 49, t.18 fig.2 (1826) 1:5
- Callitris rhomboidea** R.Br. ex Rich. & A.Rich., Comm. Bot. Conif. Cycad. 47, t.18 fig.1 (1826) 1:5
- Callitris rhomboidea R.Br. ex Rich. & A.Rich. var. tasmanica (Benth.) Ewart, B.Rees & B.Wood = **Callitris rhomboidea**
- Callitris tasmanica (Benth.) R.T.Baker & H.G.Sm. = **Callitris rhomboidea**
- e **Diselma archeri** Hook.f., Bot. Antarct. Voy. Ill. (Fl. Tasman.) 1: 353 t.98 (1857) 1:5
- Fitzroya archeri (Hook.f.) Benth. & Hook.f. = **Diselma archeri** 1:5
- Frenela australis R.Br. ex Benth. nom. illeg., sensu Bentham (1878) = **Callitris oblonga**
- Frenela rhomboidea (R.Br. ex Rich. & A.Rich.) Endl. var. tasmanica Benth. = **Callitris rhomboidea**
- i **Hesperocyparis macrocarpa** (Hartw. ex Gordon) Bartel, Phytologia 91: 182 (2009)

## PINACEAE

- i **Pinus pinaster** Aiton, Hortus Kew. (W. Aiton) 3: 367 (1789)
- i **Pinus radiata** D.Don, Trans. Linn. Soc. London 17: 442 (1837) 1:8
- i # **Pinus wallichiana** A.B.Jacks., Bull. Misc. Inform. Kew 1938(2): 85 (1938)

## PODOCARPACEAE

- Dacrydium franklinii Hook.f. = **Lagarostrobos franklinii** 1:3
- e **Lagarostrobos franklinii** (Hook.f.) Quinn, Austral. J. Bot. 30: 316 (1982) 1:3
- e **Microcachrys tetragona** (Hook.) Hook.f., London J. Bot. 4: 150 (1845) 1:3
- Microstrobos niphophilus J.Garden & L.A.S.Johnson = **Pherosphaera hookeriana** 1:2
- e **Pherosphaera hookeriana** W.Archer bis, Hooker's J. Bot. Kew Gard. Misc. 2: 52 (1850) 1:2
- e **Phyllocladus aspleniifolius** (Labill.) A.Rich. ex Hook.f., London J. Bot. 4: 151 (1845) 1:2
- Phyllocladus rhomboidalis A.Rich. = **Phyllocladus aspleniifolius** 1:2
- Podocarpus alpinus Hook.f. = **Podocarpus lawrencei** 1:4
- Podocarpus aspleniifolia Labill. = **Phyllocladus aspleniifolius**
- Podocarpus lawrencei** Hook.f., London J. Bot. 4: 151 (1845) 1:4

## PTERIDOPHYTA

### ADIANTACEAE

- Adiantum aethiopicum** L., Syst. Nat., ed. 10, 2: 1329 (1759)
- Adiantum assimile Sw. sensu Brown (1810) = **Adiantum aethiopicum**
- Adiantum trigonum Labill. = **Adiantum aethiopicum**
- Anogramma leptophylla** (L.) Link, Fil. Spec. 137 (1841)
- Cheilanthes austrotenuifolia** H.M.Quirk & T.C.Chambers, Austral. J. Bot. 31: 510 (1983)
- Cheilanthes distans** (R.Br.) Mett., Abh. Senckenberg. Naturf. Ges. 3: 69 (1859)
- Cheilanthes sieberi** Kunze subsp. **sieberi**, Pl. Preiss. [J.G.C.Lehman] 2: 112 (1846)
- Cheilanthes tenuifolia (Burm.f.) Sw. sensu Rodway (1903) = **Cheilanthes austrotenuifolia**

**Pellaea calidirupium** Brownsey & Lovis, New Zealand J. Bot. 28: 197 (1990)

**Pellaea falcata** (R.Br.) Fée, Mem. Foug., 5. Gen. Filic. 129 (1850)

*Platylooma falcata* (R.Br.) J.Sm. = **Pellaea falcata**

#### ASPLENIACEAE

**Asplenium appendiculatum** (Labill.) C.Presl subsp. **appendiculatum**, Tent. Pterid. 106 (1836)

*Asplenium brownii* J.Sm. sensu Hooker (1860) = **Diplazium australe**

**Asplenium bulbiferum** G.Forst. subsp. **gracillimum** (Colenso) Brownsey, New Zealand J. Bot. 15: 60 (1977)

*Asplenium bulbiferum* G.Forst. var. *hookeriana* (Colenso) Rodway = **Asplenium hookerianum**

**Asplenium flabellifolium** Cav., Descr. Pl. (Cavanilles): 257 (1801)

**Asplenium flaccidum** G.Forst. subsp. **flaccidum**, Fl. Ins. Austr. 80 (1786)

**Asplenium hookerianum** Colenso, Tasmanian J. Nat. Sci. 2: 169 (1844)

*Asplenium laxum* R.Br. nom. illeg. = **A. appendiculatum** × **A. bulbiferum** subsp. **gracillimum**

*Asplenium lucidum* G.Forst. sensu Hooker (1860) = **Asplenium obtusatum**

*Asplenium obliquum* G.Forst. sensu Labillardiere (1805) = **Asplenium obtusatum** subsp.

**northlandicum** (misapplied in Tasmania)

**Asplenium obtusatum** G.Forst. subsp. **northlandicum** Brownsey, New Zealand J. Bot. 15: 49, t.6 (1977)

*Asplenium terrestre* Brownsey = **Asplenium appendiculatum**

**Asplenium trichomanes** L. subsp. **quadrialeans** D.E.Mey., Ber. Deutsch. Bot. Ges. 74: 456 (1961)

**Asplenium trichomanes** L. subsp. **trichomanes**, Sp. Pl. 2: 1080 (1753)

*Asplenium umbrosum* J.Sm. sensu Bentham (1878) = **Diplazium australe** (Athyriaceae) misapplied in Tasmania

*Caenopteris appendiculata* Labill. = **Asplenium appendiculatum**

**Pleurosorus rutifolius** (R.Br.) Fée, Mem. Foug., 5. Gen. Filic. 179, t.16c (1852)

#### ATHYRIACEAE

*Allantodia australis* R.Br. = **Diplazium australe**

*Athyrium australe* (R.Br.) C.Presl = **Diplazium australe**

*Cystopteris fragilis* (L.) Bernh. var. *tasmanica* (Hook.) Hook.f. = **Cystopteris tasmanica**

**Cystopteris tasmanica** Hook., Sp. Fil. 1: 199 (1846)

**Diplazium australe** (R.Br.) N.A.Wakef., Vict. Naturalist 58: 142 t.3 (1942)

#### AZOLLACEAE

*Azolla filiculoides* Lam. = **Azolla rubra**

**Azolla rubra** R.Br., Prodr. Fl. Nov. Holland. 167 (1810)

#### BLECHNACEAE

*Blechnum aggregatum* (Colenso) Tindale sensu Willis (1973) = **Blechnum chambersii**

**Blechnum cartilagineum** Sw., Syn. Fil. (Swartz): 114 (1806)

**Blechnum chambersii** Tindale, Flora of the Sydney Region, ed. 2: 86 (1972)

**Blechnum fluviatile** (R.Br.) E.J.Lowe ex Salomon, Nomencl. Gefässkrypt. 115 (1883)

**Blechnum minus** (R.Br.) Ettingsh., Denkschr. Kaiserl. Akad. Wiss. Math.-Naturwiss. Kl. 23: 63, t.8 (1864)

**Blechnum neohollandicum** Christenh., Phytotaxa 19: 20 (2011)

**Blechnum nudum** (Labill.) Mett. ex Luerss., lora oder Allgemeinen Botanischer Zeitung 59: 292 (1876)

**Blechnum parrissiae** Christenh., Phytotaxa 19: 20 (2011)

**Blechnum patersonii** (R.Br.) Mett. subsp. **patersonii**, Fil. Hort. Bot. Lips. 64, t.4 figs. 4-10 (1856)

**Blechnum pennamarina** (Poir.) Kuhn subsp. **alpina** (R.Br.) T.C.Chambers & P.A.Farrant, Fern Gaz. 15: 96 (1996)

*Blechnum procerum* (G.Forst.) Sw. sensu Labillardiere (1805) = **Blechnum wattsii**

*Blechnum rupestre* (Kaulf. ex Link) Christenh. (cited in error) = **Blechnum spinulosum**

**Blechnum spinulosum** Poir., Encycl. (Lamarck) Suppl. 1: 644 (1810)

**Blechnum vulcanicum** (Blume) Kuhn, Ann. Mus. Bot. Lugduno-Batavi4: 284 (1869)

**Blechnum wattsii** Tindale, Contr. New South Wales Natl. Herb. 3: 247 (1963)

*Doodia aspera* R.Br. = **Blechnum neohollandicum**

*Doodia australis* (Parris) Parris = **Blechnum parrissiae**

*Doodia caudata* (Cav.) R.Br. = **Blechnum spinulosum**

*Doodia media* R.Br. = **Blechnum parrissiae** (Tasmanian plants)

*Lomaria alpina* Spreng. = **Blechnum pennamarina** subsp. **alpina**

*Lomaria capensis* (L.) Willd. sensu Bentham (1878) = **Blechnum wattsii** (misapplied in Tasmania)

*Lomaria discolor* (G.Forst.) Willd. sensu Rodway (1903) = **Blechnum nudum**

*Lomaria fluviatilis* Spreng. = **Blechnum fluviatile**

*Lomaria lanceolata* Spreng. = **Blechnum chambersii**

*Lomaria patersonii* Spreng. = **Blechnum patersonii**

*Lomaria procera* Spreng. sensu Rodway (1903) = **Blechnum wattsii**

*Lomaria procera* Spreng. var. *paludosa* Rodway = **Blechnum minus**

*Lomaria vulcanica* Blume = **Blechnum vulcanicum**

*Onoclea nuda* Labill. = **Blechnum nudum**

*Stegania alpina* R.Br. = **Blechnum pennamarina** subsp. **alpina**

*Stegania falcata* R.Br. = **Blechnum nudum**

*Stegania fluviatilis* R.Br. = **Blechnum fluviatile**

*Stegania lanceolata* R.Br. = **Blechnum chambersii**

*Stegania minor* R.Br. = **Blechnum minus**

*Stegania nuda* R.Br. = **Blechnum nudum**

*Stegania patersonii* R.Br. = **Blechnum patersonii**

*Stegania procera* R.Br. = **Blechnum wattsii**

## CULCITACEAE

**Calochlaena dubia** (R.Br.) M.D.Turner & R.A.White, Amer. Fern J. 78: 92 (1988)

*Culcita dubia* (R.Br.) Maxon = **Calochlaena dubia**

*Davallia dubia* R.Br. = **Calochlaena dubia**

## CYATHEACEAE

*Alsophila australis* R.Br. = **Cyathea australis**

**Cyathea australis** (R.Br.) Domin subsp. **australis**, Pteridophyta: 262 (1929)

**Cyathea cunninghamii** Hook.f., Icon. Pl. 10: t.985 (1854)

**Cyathea × marcescens** N.A.Wakef., Vict. Naturalist 59: 33, figs.1-5 (1942)

*Cyathea medullaris* (G.Forst.) Sw. sensu Bentham (1878) = **Cyathea cunninghamii** (misapplied in Tasmania)

## DENNSTAEDTIACEAE

**Histiopteris incisa** (Thunb.) J.Sm., Hist. Fil. 295 (1875)

**Hypolepis amaurochis** (Kunze) Hook., Sp. Fil. 2: 62 (1852)

*Hypolepis australis* N.A.Wakef. = **Hypolepis amaurochis**

- t **Hypolepis distans** Hook., Sp. Fil. 2: 70, t.95c (1852)  
**Hypolepis glandulifera** Brownsey & Chinnock, J. Adelaide Bot. Gard. 10: 16, fig.8 (1987)  
**Hypolepis muelleri** N.A.Wakef., Vict. Naturalist 60: 42 (1943)  
Hypolepis punctata (Thunb.) Mett. ex Kuhn sensu Willis (1973) = **Hypolepis glandulifera**  
**Hypolepis rugosula** (Labill.) J.Sm., Bot. Mag., 72 (Companion): 8 (1846)  
Hypolepis tenuifolia (G.Forst.) Bernh. ex C.Presl sensu Rodway (1903) = **Hypolepis amaurorachis & H. rugosula**  
Pteridium esculentum (G.Forst.) Cockayne = **Pteridium esculentum** subsp. **esculentum**  
**Pteridium esculentum** (G.Forst.) Cockayne subsp. **esculentum**, Report of the Botanical Survey of Tongariro National Park: 34 (1908)

#### DICKSONIACEAE

- Dicksonia antarctica** Labill., Nov. Holl. Pl. 2: 100, t.249 (1807)  
Dicksonia dubia (R.Br.) Gaudich. sensu Hooker (1860) = **Calochlaena dubia**

#### DRYOPTERIDACEAE

- Aspidium aculeatum (L.) Sw. sensu Bentham (1878) = **Polystichum proliferum** (misapplied in Tasmania)  
Aspidium capense Willd. nom. illeg. = **Rumohra adiantiformis**  
Aspidium coriaceum (Sw.) Sw. sensu Brown (1810) = **Rumohra adiantiformis**  
Aspidium decompositum (R.Br.) Spreng. sensu Bentham (1878) = **Lastreopsis acuminata** (misapplied in Tasmania)  
Aspidium proliferum R.Br. = **Polystichum proliferum**  
**Lastreopsis acuminata** (Houlston) C.V.Morton, Contr. U.S. Natl. Herb. 30: 245 (1973)  
**Lastreopsis hispida** (Sw.) Tindale, Vict. Naturalist 73: 183 (1957)  
Lastreopsis shepherdii (Kuntze ex Mett.) Tindale = **Lastreopsis acuminata**  
Nephrodium decompositum R.Br. sensu Rodway (1903) = **Lastreopsis acuminata**  
Nephrodium hispidum (Sw.) Hook. = **Lastreopsis hispida**  
Polystichum coriaceum Schott = **Rumohra adiantiformis**  
**Polystichum proliferum** (R.Br.) C.Presl, Tent. Pterid. 83 (1836)  
Polystichum vestitum (G.Forst.) C.Presl sensu Rodway (1903) = **Polystichum proliferum**  
**Rumohra adiantiformis** (G.Forst.) Ching, Sinensia 5: 70 (1934)

#### EQUISETACEAE

- i **Equisetum hyemale** L., Sp. Pl. 2: 1062 (1753)

#### GLEICHENIACEAE

- e **Gleichenia abscida** Rodway, The Tasmanian Flora: 289 (1903)  
e **Gleichenia alpina** R.Br., Prodr. Fl. Nov. Holland. 161 (1810)  
Gleichenia circinata Sw. sensu Rodway (1903) = **Gleichenia microphylla**  
**Gleichenia dicarpa** R.Br., Prodr. Fl. Nov. Holland. 161 (1810)  
Gleichenia flabellata R.Br. sensu Rodway (1903) = **Sticherus tener & S. urceolatus**  
**Gleichenia microphylla** R.Br., Prodr. Fl. Nov. Holland. 161 (1810)  
Gleichenia tenera R.Br. = **Sticherus tener**  
**Sticherus lobatus** N.A.Wakef., Vict. Naturalist 60: 110 (1943)  
**Sticherus tener** (R.Br.) Ching, Sunyatsenia 5: 285 (1940)  
Sticherus tener form A sensu Garrett (1996) = **Sticherus urceolatus**  
Sticherus tener form B sensu Garrett (1996) = **Sticherus tener**  
**Sticherus urceolatus** M.Garrett & Kantvilas, Muelleria 11: 103 (1998)

## GRAMMITIDACEAE

- Ctenopteris heterophylla (Labill.) Tindale = **Notogrammitis heterophylla**  
Grammitis sp. A sensu Garrett 1996 = **Notogrammitis garrettii**  
Grammitis sp. B sensu Garrett 1996 = **Notogrammitis gunnii**  
Grammitis armstrongii Tindale = **Notogrammitis crassior**  
Grammitis australis R.Br. = **Notogrammitis billardierei**  
Grammitis billardierei Willd. = **Notogrammitis billardierei**  
Grammitis garrettii Parris = **Notogrammitis garrettii**  
Grammitis gunnii Parris = **Notogrammitis gunnii**  
Grammitis heterophylla Labill. = **Notogrammitis heterophylla**  
Grammitis leptophylla (L.) Sw. sensu Bentham (1878) = **Anogramma leptophylla** (Adiantaceae)  
Grammitis magellanica Desv. subsp. nothofageti Parris = **Notogrammitis angustifolia** subsp.  
**nothofageti**  
Grammitis meridionalis Parris = **Notogrammitis angustifolia** subsp. **nothofageti** p.p.maj.  
Grammitis poeppigiana (Mett.) Pic.Serm. = **Notogrammitis crassior**  
Grammitis pseudociliata Parris = **Notogrammitis pseudociliata**  
Grammitis rutifolia R.Br. = **Pleurosorus rutifolius** (Aspleniaceae)  
**Notogrammitis angustifolia** (Jacq.) Parris subsp. **nothofageti** (Parris) Parris, New Zealand J. Bot. 50:  
466 (2012)  
**Notogrammitis billardierei** (Willd.) Parris, New Zealand J. Bot. 50: 466 (2012)  
**Notogrammitis crassior** (Kirk) Parris, New Zealand J. Bot. 50: 467 (2012)  
e **Notogrammitis garrettii** (Parris) Parris, New Zealand J. Bot. 50: 468 (2012)  
t **Notogrammitis gunnii** (Parris) Parris, New Zealand J. Bot. 50: 468 (2012)  
**Notogrammitis heterophylla** (Labill.) Parris, New Zealand J. Bot. 50: 469 (2012)  
t **Notogrammitis pseudociliata** (Parris) Parris, New Zealand J. Bot. 50: 469 (2012)

## HEMIONITIDIACEAE

- Gymnogramma leptophylla (L.) Desv. = **Anogramma leptophylla** (Adiantaceae)  
Gymnogramma rutaefolia Hook. = **Pleurosorus rutifolius** (Aspleniaceae)

## HYMENOPHYLLACEAE

- Apteropteris applanata A.M.Gray & R.G.Williams = **Hymenophyllum applanatum**  
Crepidomanes venosum (R.Br.) Bostock = **Polyphlebium venosum**  
e **Hymenophyllum applanatum** (A.M.Gray & R.G.Williams) Ebihara & K.Iwats., Bull. Natl. Mus. Nat. Sci.,  
Tokyo, B. 36: 43 (2010)  
**Hymenophyllum australe** Willd., Sp. Pl., ed. 4 [Willdenow], 5: 527 (1810)  
Hymenophyllum crispatum Hook. & Grev. sensu Hooker (1860) = **Hymenophyllum australe**  
**Hymenophyllum cupressiforme** Labill., Nov. Holl. Pl. 2: 102, t.250 fig.2 (1807)  
**Hymenophyllum flabellatum** Labill., Nov. Holl. Pl. 2: 101, t.250 fig.1 (1807)  
Hymenophyllum javanicum Spreng. sensu Rodway (1903) = **Hymenophyllum australe**  
Hymenophyllum malingii (Hook.) Mett. sensu Rodway (1903) = **Hymenophyllum applanatum**  
**Hymenophyllum marginatum** Hook. & Grev., Icon. Filic. 1: t.34 (1827)  
Hymenophyllum nitens R.Br. = **Hymenophyllum flabellatum**  
**Hymenophyllum peltatum** (Poir.) Desv., Mém. Soc. Linn. Paris 6: 333 (1827)  
**Hymenophyllum rarum** R.Br., Prodr. Fl. Nov. Holland. 159 (1810)  
Hymenophyllum tunbridgense (L.) Sm. sensu Rodway (1903) = **Hymenophyllum cupressiforme**  
Hymenophyllum unilaterale Willd. sensu Hooker (1860) = **Hymenophyllum peltatum**

- Hymenophyllum wilsoni Hook. sensu Rodway (1903) = **Hymenophyllum peltatum**  
 Mecodium australe (Willd.) Copel. = **Hymenophyllum australe**  
 Mecodium flabellatum (Labill.) Copel. = **Hymenophyllum flabellatum**  
 Mecodium rarum (R.Br.) Copel. = **Hymenophyllum rarum**  
**Polyphlebium venosum** (R.Br.) Copel., Philipp. J. Sci. 67: 55 (1938)  
 Sphaerocionium applanatum (A.M.Gray & R.G.Williams) K.Iwats. = **Hymenophyllum applanatum**  
 Trichomanes venosum R.Br. = **Polyphlebium venosum**

#### ISOETACEAE

- Isoetes sp. A sensu Garrett 1996 = **Isoetes Maxwell River (S.J.Jarman HO314082) Tas Herbarium**  
 e **Isoetes sp. Maxwell River (S.J.Jarman HO314082) Tas Herbarium**  
**Isoetes drummondii** A.Braun subsp. **drummondii**, Monatsber. Königl. Preuss. Akad. Wiss. Berlin 1868: 542 (1869)  
 e **Isoetes elatior** F.Muell. ex A.Braun, Linnaea 25: 722 (1853)  
 e **Isoetes gunnii** A.Braun, Monatsber. K. Akad. Wiss. Berlin: 535 (1868)  
 e **Isoetes humilior** F.Muell. ex A.Braun, Linnaea 25: 722 (1853)  
 Isoetes lacustris L. sensu Rodway (1903) = **Isoetes elatior, I. humilior & I. gunnii**  
**Isoetes muelleri** A.Braun, Monatsber. K. Akad. Wiss. Berlin: 541 (1868)

#### LINDSAEACEAE

- Lindsaea cuneata (G.Forst.) C.Chr. = **Lindsaea trichomanoides**  
**Lindsaea linearis** Sw., J. Bot. (Schr.) 1800(2): 78 (1801)  
**Lindsaea trichomanoides** Dryand., Trans. Linn. Soc. London 3: 43, t.11 (1797)

#### LYCOPODIACEAE

- Huperzia australiana** (Herter) Holub, Folia Geobot. Phytotax. 20: 70 (1985)  
 Huperzia varia (R.Br.) Trevis. = **Phlegmariurus varius**  
 t **Lycopodiella diffusa** (R.Br.) B.Ollg., Opera Bot. 92: 176 (1987)  
**Lycopodiella lateralis** (R.Br.) B.Ollg., Opera Bot. 92: 176 (1987)  
**Lycopodiella serpentina** (Kunze) B.Ollg., Opera Bot. 92: 176 (1987)  
 Lycopodium australianum Herter = **Huperzia australiana**  
 Lycopodium carolinianum L. sensu Bentham (1878) = **Lycopodiella serpentina** (misapplied in Tasmania)  
 Lycopodium clavatum L. var. fastigiatum (R.Br.) Benth. nom. illeg. = **Lycopodium fastigiatum**  
 Lycopodium decurrens R.Br. = **Lycopodium scariosum**  
 Lycopodium densum Labill. = **Lycopodium deuterodensum**  
**Lycopodium deuterodensum** Herter, Estud. Bot. (Montevideo) 20: 15 (1949)  
 Lycopodium diffusum R.Br. = **Lycopodiella diffusa**  
**Lycopodium fastigiatum** R.Br., Prodr. Fl. Nov. Holland. 165 (1810)  
 Lycopodium laterale R.Br. = **Lycopodiella lateralis**  
 Lycopodium myrtifolium G.Forst. nom. illeg., non L. = **Phlegmariurus varius**  
**Lycopodium scariosum** G.Forst., Fl. Ins. Austr. 87 (1786)  
 Lycopodium selago L. sensu Rodway (1903) = **Huperzia australiana**  
 Lycopodium serpentinum Kunze = **Lycopodiella serpentina**  
 Lycopodium uliginosum Labill. = **Selaginella uliginosa**  
 Lycopodium varium R.Br. = **Phlegmariurus varius**  
**Phlegmariurus varius** (R.Br.) A.R.Field & Bostock, Phytokeys 20: 49 (2013)  
**Phylloglossum drummondii** Kunze, Bot. Zeitung (Berlin) 1: 721 (1843)



### MARSILEACEAE

- i **Marsilea mutica** Mett., Ann. Sci. Nat., Bot., sér. 4, 15: 88 (1861)  
Pilularia globulifera L. sensu Rodway (1903) = **Pilularia novae-hollandiae**  
**Pilularia novae-hollandiae** A.Braun, Monatsber. Königl. Preuss. Akad. Wiss. Berlin 1863: 435 (1863)

### OPHIOGLOSSACEAE

- x **Botrychium australe** R.Br., Prodr. Fl. Nov. Holland. 164 (1810)  
**Botrychium lunaria** (L.) Sw., J. Bot. (Schrader) 1800(2): 110 (1801)  
Botrychium ternatum (Thunb.) Sw. sensu Rodway (1903) = **Botrychium australe**  
Botrychium virginianum (L.) Sw. sensu Hooker (1860) = **Botrychium australe**  
**Ophioglossum lusitanicum** L. subsp. **coriaceum** (A.Cunn.) R.T.Clausen, Mem. Torrey Bot. Club 19: 161 (1938)  
Ophioglossum vulgatum L. sensu Bentham (1878) = **Ophioglossum lusitanicum** (misapplied in Tasmania)

### OSMUNDACEAE

- Todea africana Willd. ex Bernh. nom. illeg. = **Todea barbara**  
**Todea barbara** (L.) T.Moore, Index Fil. (T. Moore): 119 (1857)

### POLYPODIACEAE

- Microsorium diversifolium (Willd.) Copel. = **Microsorium pustulatum** subsp. **pustulatum**  
**Microsorium pustulatum** (G.Forst.) Copel. subsp. **pustulatum**, Gen. Fil. [Copeland]: 196 (1947)  
Phymatodes billardierei (R.Br.) C.Presl = **Microsorium pustulatum** subsp. **pustulatum**  
Phymatosorus pustulatus (G.Forst.) Large, Braggins & P.S.Green = **Microsorium pustulatum** subsp. **pustulatum**  
Polypodium australe (R.Br.) Mett. = **Notogrammitis billardierei** (Grammitidaceae)  
Polypodium billardierei R.Br. = **Microsorium pustulatum**  
Polypodium grammitidis R.Br. = **Notogrammitis heterophylla** (Grammitidaceae)  
Polypodium punctatum Thunb. sensu Rodway (1903) = **Hypolepis rugosula** & **H. amaurorachis** (Dennstaedtiaceae)  
Polypodium pustulatum G.Forst. = **Microsorium pustulatum**  
Polypodium rugosulum Labill. = **Hypolepis rugosula** (Dennstaedtiaceae)  
Polypodium scandens Labill. nom. illeg. = **Microsorium pustulatum** subsp. **pustulatum**

### PSILOTACEAE

- Psilotum truncatum R.Br. = **Tmesipteris obliqua** (misapplied in Tasmania)  
Tmesipteris billardierei Endl. = **Tmesipteris obliqua**  
**Tmesipteris elongata** P.A.Dang., Botaniste 2: 213 (1890-91)  
Tmesipteris forsteri Endl. nom. illeg. sensu Hooker (1860) = **Tmesipteris obliqua**  
**Tmesipteris obliqua** Chinnock, Muelleria 8: 60 (1993)  
**Tmesipteris parva** N.A.Wakef., Vict. Naturalist 60: 143 (1944)  
Tmesipteris tannensis (Spreng.) Bernh. sensu Rodway (1903) = **Tmesipteris obliqua** p.p.maj.  
Tmesipteris tugana H.N.Barber = **Tmesipteris elongata**

### PTERIDACEAE

- Litobrochia comans (G.Forst.) C.Presl = **Pteris comans**  
Litobrochia incisa (Thunb.) C.Presl = **Histiopteris incisa** (Dennstaedtiaceae)  
Pteris aquilina L. sensu Rodway (1903) = **Pteridium esculentum** subsp. **esculentum** (Dennstaedtiaceae)  
**Pteris comans** G.Forst., Fl. Ins. Austr. 79 (1786)  
Pteris endlicheriana J.Agardh sensu Hooker (1860) = **Pteris comans**

*Pteris esculenta* G.Forst. sensu Labillardiere (1805) = ***Pteridium esculentum*** subsp. ***esculentum***  
(Dennstaedtiaceae)

*Pteris falcata* R.Br. = ***Pellaea falcata*** (Adiantaceae)

*Pteris incisa* Thunb. = ***Histiopteris incisa*** (Deddstaedtiaceae)

***Pteris tremula*** R.Br., Prodr. Fl. Nov. Holland. 154 (1810)

*Pteris vespertilionis* Labill. = ***Histiopteris incisa*** (Deddstaedtiaceae)

#### **SCHIZAEACEAE**

***Schizaea asperula*** N.A.Wakef., Vict. Naturalist 59: 89 (1942)

***Schizaea bifida*** Willd., Abh. Kurfurstl.-Mainz. Akad. Nützl. Wiss. Erfurt 5: 30, t.3 fig.3 (1802)

***Schizaea fistulosa*** Labill., Nov. Holl. Pl. 2: 103, t.250 fig.3 (1807)

#### **SELAGINELLACEAE**

***Selaginella gracillima*** (Kunze) Spring ex Salomon, Nomencl. Gefässkrypt. 353 (1883)

*Selaginella preissiana* Spring sensu Rodway (1903) = ***Selaginella gracillima***

***Selaginella uliginosa*** (Labill.) Spring, Mém. Acad. Roy. Sci. Belgique 24: 60 (1849)

#### **THELYPTERIDACEAE**

*Cyclosorus penniger* (G.Forst.) Ching = ***Pneumatopteris pennigera***

***Pneumatopteris pennigera*** (G.Forst.) Holttum, Blumea 21: 305 (1973)

# Macquarie Island Vascular Plant Census

## DICOTYLEDONEAE

### APIACEAE (UMBELLIFERAE)

- e **Azorella macquariensis** Orchard, Muellera 7: 16 (1989)  
Azorella selago Hook.f. (in Macquarie Island) = **Azorella macquariensis**
- t **Hydrocotyle novae-zeelandiae** DC., Prodr. [A. P. de Candolle] 4: 67 (1830)
- t **Stilbocarpa polaris** (Hombr. & Jacquinot ex Hook.f.) A.Gray, U.S. Expl. Exped., Phan. 15: 714 (1854)

### ASTERACEAE (COMPOSITAE)

- Cotula plumosa (Hook.f.) Hook.f. = **Leptinella plumosa**
- t **Leptinella plumosa** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 26 (1844)
- t **Pleurophyllum hookeri** Buchanan, Trans. & Proc. New Zealand Inst. 16: 395 (1884)

### BRASSICACEAE (CRUCIFERAE)

- Cardamine corymbosa** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 6 (1844)  
Cardamine depressa Hook.f. var. stellata (Hook.f.) Hook.f. = **Cardamine corymbosa**  
Cardamine glacialis DC. var. subcarnosa (Hook.f.) O.E.Schulz sensu Cheeseman (1919) =  
**Cardamine corymbosa**

### CALLITRICHACEAE

- Callitriche antarctica** Engelm. ex Hegelm., Verh. Bot. Vereins Prov. Brandenburg 9: 20 (1867)

### CARYOPHYLLACEAE

- i **Cerastium fontanum** Baumg. subsp. **fontanum**, Enum. Stirp. Transsilv. 1: 425 (1816)  
Cerastium triviale Link sensu Cheeseman (1919) = **Cerastium fontanum** subsp. **fontanum**  
(misapplied in Tasmania)
- Colobanthus affinis** (Hook.) Hook.f., Bot. Antarct. Voy. III. (Fl. Tasman.) 1: 45 (1855) 1:74
- t **Colobanthus apetalus** (Labill.) Druce var. **alpinus** (Kirk) L.B.Moore, Fl. New Zealand 1: 214 (1961)  
Colobanthus billardierei Fenzl sensu Cheeseman (1919) = **Colobanthus apetalus** var. **alpinus**  
(misapplied in Tasmania)
- t **Colobanthus muscoides** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 14 (1844)  
Stellaria decipiens Hook.f. sensu Cheeseman (1919) = **Stellaria parviflora** (misapplied in Tasmania)
- i **Stellaria media** (L.) Vill., Hist. Pl. Dauphiné (Villars) 3: 615 (1789) 1:71
- t **Stellaria parviflora** Banks & Sol. ex Hook.f., Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 25 (1852)

### CRASSULACEAE

- t **Crassula moschata** G.Forst., Commentat. Soc. Regiae Sci. Gott. 9: 26 (1789)  
Tillaea moschata (G.Forst.) DC. = **Crassula moschata**

### HALORAGACEAE

- t **Myriophyllum triphyllum** Orchard, Brunonia 2: 259 (1980)

**ONAGRACEAE**

- t **Epilobium brunnescens** (Cockayne) P.H.Raven & Engelhorn subsp. **brunnescens**, New Zealand J. Bot. 9: 350 (1971)  
 Epilobium linnaeoides Hook.f. = **Epilobium pedunculare**  
 Epilobium nummulariifolium A.Cunn. var. nerteroides Hook.f. sensu Cheeseman (1919) =  
**Epilobium brunnescens** subsp. **brunnescens**
- t **Epilobium pedunculare** A.Cunn., Ann. Nat. Hist. 3: 31 (1839)

**POLYGONACEAE**

FTO 95

- i # x **Rumex crispus** L., Sp. Pl. 1: 335 (1753)

**PORTULACACEAE**

FTO 100

- Montia fontana** L. subsp. **fontana**, Sp. Pl. 1: 87 (1753)

**RANUNCULACEAE**

FTO 47

- Ranunculus biternatus Sm. sensu Cheeseman (1919) = **Ranunculus crassipes**  
**Ranunculus crassipes** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 2: 224 (1845)

**ROSACEAE**

- Acaena adscendens Vahl Sensu Hooker (1844) = **Acaena magellanica**  
**Acaena magellanica** (Lam.) Vahl, Enum. Pl. [Vahl] 1: 297 (1804)
- t **Acaena minor** (Hook.f.) Allan, Fl. New Zealand 1: 363 (1961)  
 Acaena sanguisorbae Vahl var. minor Hook.f. = **Acaena minor**

**RUBIACEAE**

- t **Coprosma perpusilla** Colenso subsp. **subantarctica** Orchard, Brunonia 9: 133 (1987)  
 Coprosma repens Hook.f. (nom. illeg.) = **Coprosma perpusilla** subsp. **subantarctica**
- t **Galium antarcticum** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 303 bis (1846)

**MONOCOTYLEDONEAE**

**CYPERACEAE**

- t **Carex trifida** Cav., Icon. [Cavanilles] 5: 41, t.465 (1799)
- Isolepis aucklandica** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 88 (1844), pl.50 (1845) 4b:102  
 Scirpus aucklandicus (Hook.f.) Boeckeler = **Isolepis aucklandica** 4b:102
- t **Uncinia divaricata** Boott, Bot. Antarct. Voy. II. (Fl. Nov.-Zel.) 1: 286 (1853)
- t **Uncinia hookeri** Boott, Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 91 (1844); pl.51 (1845)  
 Uncinia riparia R.Br. var. hookeri (Boott) Kük. = **Uncinia hookeri**

**JUNCACEAE**

- t **Juncus scheuchzerioides** Gaudich., Ann. Sci. Nat. (Paris) 5: 100 (1825)  
 Luzula campestris DC. var. crinita (Hook.f.) Buchenau = **Luzula crinita**
- t **Luzula crinita** Hook.f., Bot. Antarct. Voy. I. (Fl. Antarct.) 1: 84, pl.48 (1844)

**ORCHIDACEAE**

- n e **Corybas dienemus** D.L.Jones, Fl. Australia 50: 572 (1993) 4a:121  
 Corybas macranthus (Hook.f.) Rchb.f. = **Corybas dienemus**
- n e **Corybas sulcatus** (M.A.Clem. & D.L.Jones) G.N.Backh., Vict. Naturalist 127: 56 (2010)
- n **Nematoceras dienemum** (D.L.Jones) D.L.Jones, M.A.Clem. & Molloy = **Corybas dienemus**

n *Nematoceras sulcatum* M.A.Clem. & D.L.Jones = **Corybas sulcatus**

### POACEAE (GRAMINEAE)

i # **Agrostis capillaris** L., Sp. Pl. 1: 62 (1753)

t **Agrostis magellanica** Lam., Tabl. Encycl. 1: 160 (1791)

i # **Agrostis stolonifera** L., Sp. Pl. 1: 62 (1753)

i # x **Anthoxanthum odoratum** L., L., Sp. Pl. 1: 28 (1753)

**Deschampsia cespitosa** (L.) P.Beauv., Ess. Agrostogr. 91: 160, t.18, fig.3 (1812) 4b:242

t **Deschampsia chapmanii** Petrie, Trans. & Proc. New Zealand Inst. 23: 401 (1891)

*Deschampsia penicillata* Kirk = **Deschampsia cespitosa**

t **Festuca contracta** Kirk, Trans. & Proc. New Zealand Inst. 27: 353 (1895)

*Festuca erecta* d'Urv. nom. illeg. = **Festuca contracta**

i **Poa annua** L., Sp. Pl. 1: 68 (1753) 4b:211

**Poa cookii** (Hook.f.) Hook.f., Philos. Trans. 168: 22 (1879)

t **Poa foliosa** (Hook.f.) Hook.f., Handb. N. Zeal. Fl. 338 (1864)

*Poa hamiltonii* Kirk = **Poa cookii**

t **Poa litorosa** Cheeseman, Man. New Zealand Fl. 1156 (1906)

e **Puccinellia macquariensis** (Cheeseman) Allan & Jansen, Trans. & Proc. Roy. Soc. New Zealand 69: 268 (1939)

*Triodia macquariensis* Cheeseman = **Puccinellia macquariensis**

## PTERIDOPHYTA

### BLECHNACEAE

**Blechnum pennamarina** (Poir.) Kuhn subsp. **alpina** (R.Br.) T.C.Chambers & P.A.Farrant, Fern Gaz. 15: 96 (1996)

*Lomaria penna-marina* Trevis. = **Blechnum pennamarina** subsp. **alpina**

### DRYOPTERIDACEAE

*Aspidium vestitum* (G.Forst.) Sw. = **Polystichum vestitum**

t **Polystichum vestitum** (G.Forst.) C.Presl, Tent. Pterid. 83 (1836)

### GRAMMITIDACEAE

*Grammitis poeppigiana* (Mett.) Pic.Serm. = **Notogrammitis crassior**

**Notogrammitis crassior** (Kirk) Parris, New Zealand J. Bot. 50: 467 (2012)

*Polypodium australe* (R.Br.) Mett. nom. illeg. sensu Cheeseman (1919) = **Notogrammitis crassior**

### HYMENOPHYLLACEAE

t **Hymenophyllum falklandicum** Baker, Syn. Fil. (Hooker & Baker), ed. 2: 68 (1874)

### LYCOPODIACEAE

**Huperzia australiana** (Herter) Holub, Folia Geobot. Phytotax. 20: 70 (1985)

*Lycopodium varium* R.Br. sensu Cheeseman (1919) = **Huperzia australiana**

---

**APPENDIX I: CHANGES IN THE CENSUS BETWEEN THE 2016 AND 2017 EDITIONS**

**Ia: Taxa new to the Tasmanian Census**

**DICOTYLEDONEAE**

**ASTERACEAE**

**Chrysocephalum semipapposum**

subsp. **asperum** (Steetz) Paul  
G.Wilson

**Chrysocephalum semipapposum**

subsp. **lineare** Paul G.Wilson

**Microseris walteri** Gand.  
e **Xerochrysum alpinum** Paul  
G.Wilson

**EPACRIDACEAE**

e **Leptecophylla pogonocalyx** subsp.  
**decipiens** Jarman

---

**APPENDIX I: CHANGES IN THE CENSUS BETWEEN THE 2016 AND 2017 EDITIONS**

**Ib: Taxa that have changed names between 2016 and 2017**

**DICOTYLEDONEAE**

**ASTERACEAE**

Helichrysum milliganii = **Xerochrysum milliganii** (*Taxon* 64: 106 [2015])

**EPACRIDACEAE**

Leptecophylla juniperina subsp. oxycedrus =  
**Leptecophylla oxycedrus** (*Swainsona* 31: 1 [2017])

Leptecophylla juniperina subsp. parvifolia =  
**Leptecophylla parvifolia** (*Swainsona* 31: 7 [2017])

**FABACEAE**

Medicago falcata = **Medicago sativa**  
nothosubsp. × **varia** (previously  
misapplied)

**MIMOSACEAE**

Acacia implexa = **Acacia uncifolia** (previously  
misapplied)

**THYMELEACEAE**

Pimelea sp. Freycinet (A.M.Buchanan 15902) Tas  
Herbarium = **Pimelea leiophylla**

**MONOCOTYLEDONEAE**

**ORCHIDACEAE**

Nematoceras dienemum = **Corybas dienemus**

Nematoceras sulcatum = **Corybas sulcatus**

## APPENDIX I: CHANGES IN THE CENSUS BETWEEN THE 2016 AND 2017 EDITIONS

### Ic: Taxa that have changed status between 2016 and 2017

#### DICOTYLEDONEAE

##### ASTERACEAE

- ix *Cynara cardunculus* subsp. *flavescens*
- ix *Onopordum acaulon*
- ix *Pilosella officinarum* subsp. *officinarum*
- i *Senecio angulatus*

##### BORAGINACEAE

- i *Lithospermum officinale*

##### BRASSICACEAE

*Cardamine tryssa* is no longer considered extinct in Tasmania (recently collected in the Central Plateau)

- i *Lunaria annua*
- i *Nasturtium microphyllum*

##### CRASSULACEAE

- i *Crassula natans* var. *minus*

##### CUSCUTACEAE

- ix *Cuscuta suaveolens*

##### FABACEAE

- i *Securigera varia*

##### GERANIACEAE

- i *Geranium yeoi*

##### MALVACEAE

- i *Malva pseudolavatera*

##### MIMOSACEAE

- i *Acacia baileyana*

##### MYRTACEAE

- e *Melaleuca virens* is endemic to Tasmania (previous versions were not listed as endemic in error)

##### PROTEACEAE

- ix *Hakea laurina*

##### RANUNCULACEAE

- i *Aquilegia vulgaris*
- i *Ranunculus acris* subsp. *acris*
- i *Ranunculus trilobus*

##### ROSACEAE

- i *Rubus philadelphicus*
- i *Rubus rubritinctus*

##### SOLANACEAE

- i *Physalis peruviana*
- i *Solanum triflorum*

#### MONOCOTYLEDONEAE

##### ALOACEAE

- i *Kniphofia uvaria*

##### CYPERACEAE

- i *Isolepis hystrix*

##### POACEAE

- i *Eleusine tristachya*
- i *Eragrostis curvula*
- i *Hordeum hystrix*



## APPENDIX I: CHANGES IN THE CENSUS BETWEEN THE 2016 AND 2017 EDITIONS

Id: Taxa now not considered to be part of the Tasmanian Flora

### DICOTYLEDONEAE

#### AMARANTHACEAE

- Amaranthus graecizans subsp. silvestris †
- Amaranthus spinosus †

#### APIACEAE

- Aegopodium podagraria †

#### ASTERACEAE

- i\* Centaurea calcitrapa
- i\* Centaurea cyanus
- Centaurea solstitialis †
- i\* Lactuca serriola f. integrifolia
- i\* Matricaria chamomilla
- Taraxacum kok-saghyz †

#### BASELLACEAE

- i\* Anredera cordifolia

#### BETULACEAE

- Alnus cordata †
- i\* Alnus glutinosa

#### BORAGINACEAE

- i\* Symphytum × uplandicum

#### BRASSICACEAE

- i\* Brassica oleracea
- Carrichtera annua †
- Eruca sativa †
- Lepidium heterophyllum †
- i\* Raphanus maritimus
- i\* Thlaspi arvense

#### CALLITRICHACEAE

- i\* Callitriche brutia

#### CAMPANULACEAE

- i\* Campanula rapunculoides

#### CAPRIFOLIACEAE

- i\* Viburnum tinus

#### CARYOPHYLLACEAE

- Silene conica †

- i\* Stellaria graminea
- Vaccaria hispanica †

#### CHENOPODIACEAE

- i\* Bassia scoparia
- Chenopodium foliosum †

#### CLUSIACEAE

- Hypericum humifusum †

#### EPACRIDACEAE

- Leptecophylla juniperina after the revision by Jarman & Kantvilas (2017) *L. juniperina* is no longer found in Tasmania

#### ERICACEAE

- i\* Arbutus unedo

#### EUPHORBIACEAE

- i\* Euphorbia stricta

#### FABACEAE

- Hedysarum coronarium †
- Onobrychis viciifolia †
- Trifolium uniflorum †

#### FUMARIACEAE

- i\* Fumaria officinalis
- Pseudofumaria alba subsp. alba †

#### GERANIACEAE

- i\* Erodium malacoides

#### LENTIBULARIACEAE

- i\* Utricularia gibba

#### MALVACEAE

- Hibiscus trionum †

#### ONAGRACEAE

- i\* Epilobium nummulariifolium

#### POLEMONIACEAE

- Collomia grandiflora †

**PORTULACACEAE**

*Claytonia perfoliata* subsp. *perfoliata* †

**PRIMULACEAE**

i\* *Lysimachia minima*

**RANUNCULACEAE**

i\* *Adonis microcarpa*  
*Ranunculus arvensis* †  
*Ranunculus flammula* subsp. *flammula* †  
*Ranunculus sceleratus* subsp. *sceleratus* †

**RUBIACEAE**

*Galium tricornutum* †

**SALICACEAE**

i\* *Salix alba*  
i\* *Salix matsudana*  
i\* *Salix* × *calodendron*  
i\* *Salix* × *rubens*  
i\* *Salix* × *sepulcralis*

**SCROPHULARIACEAE**

i\* *Kickxia spuria*  
*Veronica peregrina* †

**SOLANACEAE**

*Hyoscyamus albus* †  
*Nicotiana sylvestris* †  
i\* *Solanum nodiflorum*

**URTICACEAE**

i\* *Paretaria judaica*

**MONOCOTYLEDONEAE**

**CYPERACEAE**

i\* *Carex bauxbaumii*  
i\* *Carex pilulifera*  
i\* *Carex scoparia*  
*Carex testacea* †

**HYDROCHARITACEAE**

*Lagarosiphon major* †

**POACEAE**

i\* *Aira cupaniana*  
i\* *Avellinia michelii*  
*Dichanthium sericeum* subsp. *sericeum* †  
*Digitaria ciliaris* †  
*Digitaria ternata* †  
*Echinochloa oryzoides* †  
i\* *Eragrostis tenuifolia*  
i\* *Glyceria plicata*  
i\* *Holcus mollis*  
i\* *Molineriella minuta*  
*Panicum capillare* †  
i\* *Panicum gilvum*  
*Setaria pumila* subsp. *pumila* †  
*Sorghum bicolor* †  
i\* *Sorghum halepense*

\* Taxa marked as doubtfully naturalised are no longer considered to be part of the Tasmanian flora

† Taxa for which there is insufficient evidence that they have ever naturalised (Baker *et al.* in preparation)

## REFERENCES

- ABRS (Australian Biological Resources Study) (Australian Government: Department of the Environment, Water, Heritage and the Arts). <http://www.environment.gov.au/biodiversity/abrs/index.html>
- APG II (2003) An update of the Angiosperm Phylogeny Group classification for the orders and families of flowering plants: APG II. *Botanical Journal of the Linnean Society* 141: 399–436.
- Baker ML, de Salas MF (2012) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania & Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- Baker ML, de Salas MF (2013) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania & Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- Baker ML, Duretto MF (2011) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania & Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- Bentham G (1863) *Flora Australiensis*, Vol. I. (Lovell Reeve and Co.: London)
- Bentham G (1864) *Flora Australiensis*, Vol. II (Lovell Reeve and Co.: London)
- Bentham G (1867) *Flora Australiensis*, Vol. III (Lovell Reeve and Co.: London)
- Bentham G (1868) *Flora Australiensis*, Vol. IV (Lovell Reeve and Co.: London)
- Bentham G (1870) *Flora Australiensis*, Vol. V (Lovell Reeve and Co.: London)
- Bentham G (1873) *Flora Australiensis*, Vol. VI (Lovell Reeve and Co.: London)
- Bentham G (1878) *Flora Australiensis*, Vol. VII (Lovell Reeve and Co.: London)
- Brown R (1810) *Prodromus Florae Novae Hollandiae et Insulae van-Diemen* (Richard Taylor & Son: London)
- Buchanan AM (1995) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania*. Tasmanian Herbarium Occasional Publication No. 5 (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart)
- Buchanan AM (1999) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania*. 3rd edn. Tasmanian Herbarium Occasional Publication No. 6 (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart)
- Buchanan AM (2004) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- Buchanan AM (2005) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania*. 4th edn. Tasmanian Herbarium Occasional Publication No. 7 (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart)
- Buchanan AM (2007) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- Buchanan AM (2009) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- Buchanan AM, McGeary-Brown A, Orchard AE (1989) *A Census of the Vascular Plants of Tasmania*. Tasmanian Herbarium Occasional Publication No. 2 (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart)

- Cheeseman TF (1919) The vascular flora of Macquarie Island. *Australasian Antarctic Expedition 1911–1914. Scientific Reports. Series C. Zoology and Botany.* Volume VII, Part 3. 63 pp. (Government Printer: Sydney)
- Cronquist AJ (1981) *An Integrated System of Classification of Flowering Plants.* (Columbia University Press: New York)
- Curtis (1956) *The Student's Flora of Tasmania Part I – Gymnospermae; Angiospermae: Ranunculaceae to Myrtaceae.* 1<sup>st</sup> edn. (Government Printer: Hobart)
- Curtis WM (1963) *The Student's Flora of Tasmania Part 2 – Angiospermae: Lythraceae to Epacridaceae.* (Government Printer: Hobart)
- Curtis WM (1967) *The Student's Flora of Tasmania Part 3 – Angiospermae: Plumbaginaceae to Salicaceae.* (Government Printer: Hobart)
- Curtis WM (1979) *The Student's Flora of Tasmania Part 4A – Angiospermae: Orchidaceae.* (Government Printer: Hobart)
- Curtis WM, Morris DI (1975) *The Student's Flora of Tasmania Part I – Gymnospermae; Angiospermae: Ranunculaceae to Myrtaceae.* 2<sup>nd</sup> edn. (Government Printer: Hobart)
- Curtis WM, Morris DI (1994) *The Student's Flora of Tasmania Part 4B – Angiospermae: Alismataceae to Burmanniaceae.* (St. Davids Park Publishing: Hobart)
- de Salas MF, Baker ML (2014) *A Census of the Vascular Plants of Tasmania & Index to The Student's Flora of Tasmania & Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- de Salas MF, Baker ML (2015) *A Census of the Vascular Plants of Tasmania including Macquarie Island* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- de Salas MF, Baker ML (2016) *A Census of the Vascular Plants of Tasmania including Macquarie Island* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart) [www.tmag.tas.gov.au](http://www.tmag.tas.gov.au)
- Duretto MF (2009+) *Flora of Tasmania Online* (Tas. Herbarium, Tas. Museum & Art Gallery: Hobart). [www.tmag.tas.gov.au/floratasmania](http://www.tmag.tas.gov.au/floratasmania)
- Garrett M (1996) *The Ferns of Tasmania; Their Ecology and Distribution* (Tasmanian Forest Research Council: Hobart)
- Harris S, Buchanan A, Connolly A (2001) *One Hundred Islands: The Flora of the Outer Fureaux.* (Tasmanian DPIWE: Hobart). p. 197
- Henderson RJF in George AS (Ed) (1987) *Flora of Australia* 45: 209–214, 477–479 (ABRS: Canberra)
- Hewson HJ in George AS (Ed) (1982) *Flora of Australia* 8: 313, map 366 (ABRS: Canberra)
- Hooker JD (1844) *The Botany of the Antarctic Voyage of H. M. Discovery Ships Erebus and Terror in the Years 1839 – 1843 Under the Command of Captain Sir James Clark Ross. Part I: Flora Antarctica* (Reeve Brothers: London)
- Hooker JD (1847) Contributions towards a Flora of Van Diemen's Land. *London Journal of Botany* 6: 461(bis)–479(bis)
- Hooker JD (1860) *The Botany of the Antarctic Voyage of H. M. Discovery Ships Erebus and Terror in the Years 1839 – 1843 Under the Command of Captain Sir James Clark Ross. Part III: Flora Tasmaniae* (Reeve Brothers: London)
- Jarman SJ and Kantvilas GK (2017) *Leptecophylla* in Tasmania: a reassessment of four species. *Swainsona* 31: 1–16

- Labillardiere JJH de (1804–1807) *Novae Hollandiae Plantarum Specimen* (Typographia Dominae Huzard: Paris)
- Maiden JH, Betche E (1916) *A Census of New South Wales Plants* (W. A. Gullick, Government Printer: Sydney)
- McCarthy PM (1998) Ferns, Gymnosperms and Allied Groups. *Flora of Australia* 48. (ABRS: Canberra)
- Morris DI (1977) Miscellaneous notes on endemic Tasmanian plants in the genera *Olearia*, *Ixodia*, *Danthonia*, *Tetrarrhena*. *Records of the Queen Victoria Museum* 55: 1–5
- Raphael TD (1955) Tasmanian garden escapes. *Papers and Proceedings of the Royal Society of Tasmania* 89: 147–150
- Rodway L (1903) *The Tasmanian Flora*. (Government Printer: Hobart)
- Rye BL (1990) Thymelaeaceae. *Flora of Australia* 18: 181, Map 247 (ABRS: Canberra)
- Smith JJ (1912) Epacridaceae. *Nova Guinea* 8: 797
- Townrow JES (1969) A species list of and keys to the grasses of Tasmania. *Papers and Proceedings of the Royal Society of Tasmania* 103: 69–96
- Townrow J (1978) *Papers and Proceedings of the Royal Society of Tasmania* 112
- Willis JH (1970) *A Handbook to Plants in Victoria*. (Melbourne Uni. Press: Melbourne)
- Willis JH (1973) *A Handbook to Plants of Victoria. Vol. 1 – Ferns, Conifers and Monocotyledons*. Ed. 2 (Melbourne Uni. Press: Melbourne)

**APPENDIX 2. FAMILY PLACEMENT OF GENERA IN THIS CENSUS**

(see Introduction for details regarding this appendix. Names not in current use in Tasmania are indented and in italics)

Abrotanella	Asteraceae	D	Alopecurus	Poaceae	M
Acacia	Mimosaceae	D	<i>Alsophila</i>	Cyatheaceae	P
Acaena	Rosaceae	D	Alstroemeria	Liliaceae	M
Acanthus	Acanthaceae	D	Alternanthera	Amaranthaceae	D
Acer	Aceraceae	D	<i>Alyssum</i>	Brassicaceae	D
Acetosa	Polygonaceae	D	Alyxia	Apocynaceae	D
Acetosella	Polygonaceae	D	Amaranthus	Amaranthaceae	D
Achillea	Asteraceae	D	Amelichloa	Poaceae	M
<i>Achnatherum</i>	Poaceae	M	Ammi	Apiaceae	D
Acianthus	Orchidaceae	M	Ammobium	Asteraceae	D
<i>Acion</i>	Restionaceae	M	Ammophila	Poaceae	M
<i>Aciphylla</i>	Apiaceae	D	Amperea	Euphorbiaceae	D
Acradenia	Rutaceae	D	Amphibolis	Cymodoceaceae	M
Acrothamnus	Epacridaceae	D	Amphibromus	Poaceae	M
Acrotriche	Epacridaceae	D	Amsinckia	Boraginaceae	D
Actinotus	Apiaceae	D	<i>Anagallis</i>	Primulaceae	D
Actites	Asteraceae	D	Anchusa	Boraginaceae	D
Adiantum	Adiantaceae	P	Androstoma	Epacridaceae	D
Adonis	Ranunculaceae	D	Anemone	Ranunculaceae	D
<i>Adriana</i>	Euphorbiaceae	D	Angianthus	Asteraceae	D
<i>Aegopodium</i>	Apiaceae	D	<i>Anguillaria</i>	Liliaceae	M
Aeonium	Crassulaceae	D	Anigozanthos	Haemodoraceae	M
Agapanthus	Liliaceae	M	Anisodonteia	Malvaceae	D
Agastachys	Proteaceae	D	Anisotome	Apiaceae	D
Agave	Agavaceae	M	Anodopetalum	Cunoniaceae	D
<i>Agropyron</i>	Poaceae	M	Anogramma	Adiantaceae	P
Agrostemma	Caryophyllaceae	D	Anopterus	Escalloniaceae	D
Agrostis	Poaceae	M	Anredera	Basellaceae	D
Aira	Poaceae	M	Anthemis	Asteraceae	D
Ajuga	Lamiaceae	D	<i>Anthericum</i>	Liliaceae	M
<i>Albizia</i>	Mimosaceae	D	<i>Anthistiria</i>	Poaceae	M
<i>Alchemilla</i>	Rosaceae	D	<i>Anthocercis</i>	Solanaceae	D
<i>Alepyrum</i>	Centrolepidaceae	M	Anthosachne	Poaceae	M
<i>Aletris</i>	Liliaceae	M	Anthoxanthum	Poaceae	M
Alisma	Alismataceae	M	Anthriscus	Apiaceae	D
<i>Allantodia</i>	Athyriaceae	P	<i>Anthyllis</i>	Fabaceae	D
Allittia	Asteraceae	D	Antirrhinum	Scrophulariaceae	D
Allium	Liliaceae	M	<i>Anzybas</i>	Orchidaceae	M
Allocasuarina	Casuarinaceae	D	Aotus	Fabaceae	D
Almaleea	Fabaceae	D	Apalochlamys	Asteraceae	D
Alnus	Betulaceae	D	Aphanes	Rosaceae	D
Aloe	Aloeaceae	M	Aphelia	Centrolepidaceae	M

Apium	Apiaceae	D	Austrostipa	Poaceae	M
Apodasmia	Restionaceae	M	Avellinia	Poaceae	M
Aponogeton	Aponogetonaceae	M	Avena	Poaceae	M
<i>Aptenia</i>	Aizoaceae	D	Azolla	Azollaceae	P
<i>Apteropteris</i>	Hymenophyllaceae	P	Azorella	Apiaceae	D
Aquilegia	Ranunculaceae	D	Baeckea	Myrtaceae	D
Arabidopsis	Brassicaceae	D	Ballantinia	Brassicaceae	D
<i>Arachnorchis</i>	Orchidaceae	M	Baloskion	Restionaceae	M
Arbutus	Ericaceae	D	Banksia	Proteaceae	D
Archeria	Epacridaceae	D	Barbarea	Brassicaceae	D
Arctium	Asteraceae	D	Bartsia	Scrophulariaceae	D
Arctotheca	Asteraceae	D	Bassia	Chenopodiaceae	D
Arctotis	Asteraceae	D	<i>Batrachium</i>	Ranunculaceae	D
Arenaria	Caryophyllaceae	D	Bauera	Cunoniaceae	D
Argemone	Papaveraceae	D	Baumea	Cyperaceae	M
Argentina	Rosaceae	D	Bedfordia	Asteraceae	D
Argentipallium	Asteraceae	D	<i>Bellardia</i>	Scrophulariaceae	D
Argyrotegium	Asteraceae	D	Bellenden	Proteaceae	D
Aristida	Poaceae	M	Bellis	Asteraceae	D
Aristotelia	Elaeocarpaceae	D	Berberis	Berberidaceae	D
Arrhenatherum	Poaceae	M	Berkheya	Asteraceae	D
<i>Arthrochilus</i>	Orchidaceae	M	Bertya	Euphorbiaceae	D
<i>Arthrocnemum</i>	Chenopodiaceae	D	Beta	Chenopodiaceae	D
Arthropodium	Liliaceae	M	Betula	Betulaceae	D
<i>Arundo</i>	Poaceae	M	Beyeria	Euphorbiaceae	D
<i>Ascyrum</i>	Clusiaceae	D	Billardiera	Pittosporaceae	D
Asparagus	Liliaceae	M	Blandfordia	Liliaceae	M
Asperula	Rubiaceae	D	Blechnum	Blechnaceae	P
Asphodelus	Liliaceae	M	<i>Boisduvalia</i>	Onagraceae	D
<i>Aspidium</i>	Dryopteridaceae	P	Bolboschoenus	Cyperaceae	M
Asplenium	Aspleniaceae	P	Borago	Boraginaceae	D
Astelia	Liliaceae	M	Boronia	Rutaceae	D
<i>Aster</i>	Asteraceae	D	Bossiaea	Fabaceae	D
Asterotrichion	Malvaceae	D	Bothriochloa	Poaceae	M
Astroloma	Epacridaceae	D	Botrychium	Ophioglossaceae	P
Atherosperma	Monimiaceae	D	<i>Brachyglottis</i>	Asteraceae	D
Athrotaxis	Cupressaceae	G	Brachyloma	Epacridaceae	D
<i>Athyrium</i>	Athyriaceae	P	Brachypodium	Poaceae	M
Atriplex	Chenopodiaceae	D	Brachyscome	Asteraceae	D
Australina	Urticaceae	D	<i>Bracteantha</i>	Asteraceae	D
Australopyrum	Poaceae	M	Brassica	Brassicaceae	D
Austrocynoglossum	Boraginaceae	D	Briza	Poaceae	M
<i>Austrodanthonia</i>	Poaceae	M	Bromus	Poaceae	M
Austroderia	Poaceae	M	<i>Brossaea</i>	Ericaceae	D
Austrofestuca	Poaceae	M	Brunonia	Brunoniaceae	D

Buddleja	Buddlejaceae	D	<i>Carpodontos</i>	Eucryphiaceae	D
Buglossoides	Boraginaceae	D	<i>Carrichtera</i>	Brassicaceae	D
Bulbine	Liliaceae	M	Carthamus	Asteraceae	D
<i>Bunochilus</i>	Orchidaceae	M	Cassinia	Asteraceae	D
Burchardia	Liliaceae	M	Cassytha	Lauraceae	D
Burnettia	Orchidaceae	M	<i>Casuarina</i>	Casuarinaceae	D
Bursaria	Pittosporaceae	D	Catapodium	Poaceae	M
<i>Cacalia</i>	Asteraceae	D	<i>Caucalis</i>	Apiaceae	D
<i>Caenopteris</i>	Aspleniaceae	P	<i>Caulinia</i>	Posidoniaceae	M
Caesia	Liliaceae	M	Caustis	Cyperaceae	M
Cakile	Brassicaceae	D	Cedronella	Lamiaceae	D
Caladenia	Orchidaceae	M	Celmisia	Asteraceae	D
Calamagrostis	Poaceae	M	<i>Celsia</i>	Scrophulariaceae	D
Calandrinia	Portulacaceae	D	Cenarrhenes	Proteaceae	D
<i>Caldasia</i>	Apiaceae	D	Cenchrus	Poaceae	M
<i>Calea</i>	Asteraceae	D	Centaurea	Asteraceae	D
Caleana	Orchidaceae	M	Centaurium	Gentianaceae	D
Calendula	Asteraceae	D	Centella	Apiaceae	D
Callistachys	Fabaceae	D	Centipeda	Asteraceae	D
<i>Callistemon</i>	Myrtaceae	D	Centranthus	Valerianaceae	D
Callitriche	Callitrichaceae	D	Centrolepis	Centrolepidaceae	M
Callitris	Cupressaceae	G	Centropappus	Asteraceae	D
Calluna	Ericaceae	D	Cerastium	Caryophyllaceae	D
Calocephalus	Asteraceae	D	<i>Ceratochloa</i>	Poaceae	M
Calochilus	Orchidaceae	M	<i>Chaetospora</i>	Cyperaceae	M
Calochlaena	Culcitaceae	P	Chamaecytisus	Fabaceae	D
<i>Calonema</i>	Orchidaceae	M	Chamaemelum	Asteraceae	D
<i>Calonemorchis</i>	Orchidaceae	M	Chamaescilla	Liliaceae	M
Calorophus	Restionaceae	M	<i>Chamaesyce</i>	Euphorbiaceae	D
<i>Caltha</i>	Ranunculaceae	D	Chasmanthe	Iridaceae	M
<i>Calycotrix</i>	Myrtaceae	D	<i>Cheesemania</i>	Brassicaceae	D
Calystegia	Convolvulaceae	D	Cheilanthes	Adiantaceae	P
Calytrix	Myrtaceae	D	Chenopodium	Chenopodiaceae	D
Camelina	Brassicaceae	D	Chiloglottis	Orchidaceae	M
Campanula	Campanulaceae	D	Chionogentias	Gentianaceae	D
Campynema	Liliaceae	M	<i>Chionohebe</i>	Scrophulariaceae	D
<i>Candollea</i>	Stylidiaceae	D	<i>Chlorophytum</i>	Liliaceae	M
<i>Canthium</i>	Rubiaceae	D	<i>Chondrilla</i>	Asteraceae	D
Capsella	Brassicaceae	D	Chordifex	Restionaceae	M
Cardamine	Brassicaceae	D	Chorizandra	Cyperaceae	M
<i>Cardaria</i>	Brassicaceae	D	Chrysanthemoides	Asteraceae	D
Carduus	Asteraceae	D	<i>Chrysanthemum</i>	Asteraceae	D
Carex	Cyperaceae	M	Chrysocephalum	Asteraceae	D
Carpha	Cyperaceae	M	<i>Chrysocoma</i>	Asteraceae	D
Carpobrotus	Aizoaceae	D	Cicendia	Gentianaceae	D



Cichorium	Asteraceae	D	Cullen	Fabaceae	D
Cirsium	Asteraceae	D	<i>Cuphonotus</i>	Brassicaceae	D
Cistus	Cistaceae	D	Cuscuta	Cuscutaceae	D
Cladium	Cyperaceae	M	<i>Cyanicula</i>	Orchidaceae	M
<i>Claytonia</i>	Portulacaceae	D	Cyathea	Cyatheaceae	P
Clematis	Ranunculaceae	D	Cyathodes	Epacridaceae	D
<i>Cnicus</i>	Asteraceae	D	<i>Cyclosorus</i>	Thelypteridaceae	P
Coleonema	Rutaceae	D	Cyclosporum	Apiaceae	D
<i>Collomia</i>	Polemoniaceae	D	Cynogeton	Juncaginaceae	M
Colobanthus	Caryophyllaceae	D	Cymbalaria	Scrophulariaceae	D
Comesperma	Polygalaceae	D	Cymbonotus	Asteraceae	D
Conium	Apiaceae	D	<i>Cymodocea</i>	Cymodoceaceae	M
Conospermum	Proteaceae	D	Cynara	Asteraceae	D
Convolvulus	Convolvulaceae	D	Cynodon	Poaceae	M
Conyza	Asteraceae	D	Cynoglossum	Boraginaceae	D
Cooperhooikia	Goodeniaceae	D	Cynosurus	Poaceae	M
Coprosma	Rubiaceae	D	Cyperus	Cyperaceae	M
Cordyline	Agavaceae	M	Cyphanthera	Solanaceae	D
Coronidium	Asteraceae	D	Cyrtostylis	Orchidaceae	M
<i>Coronilla</i>	Fabaceae	D	<i>Cystanthe</i>	Epacridaceae	D
<i>Coronopus</i>	Brassicaceae	D	Cystopteris	Athyriaceae	P
Correa	Rutaceae	D	Cytisus	Fabaceae	D
Cortaderia	Poaceae	M	<i>Dacrydium</i>	Podocarpaceae	G
Corunastylis	Orchidaceae	M	Dactylis	Poaceae	M
Corybas	Orchidaceae	M	Damasonium	Alismataceae	M
<i>Corysanthes</i>	Orchidaceae	M	Dampiera	Goodeniaceae	D
<i>Cota</i>	Asteraceae	D	Danthonia	Poaceae	M
Cotoneaster	Rosaceae	D	Daphne	Thymelaeaceae	D
Cotula	Asteraceae	D	Datura	Solanaceae	D
Cotyledon	Crassulaceae	D	Daucus	Apiaceae	D
<i>Crangonorchis</i>	Orchidaceae	M	<i>Davallia</i>	Culcitaceae	P
<i>Crantzia</i>	Apiaceae	D	Daviesia	Fabaceae	D
Craspedia	Asteraceae	D	<i>Decaspora</i>	Epacridaceae	D
Crassula	Crassulaceae	D	Delairea	Asteraceae	D
Crataegus	Rosaceae	D	<i>Dendrobium</i>	Orchidaceae	M
<i>Crepidomanes</i>	Hymenophyllaceae	P	<i>Derwentia</i>	Scrophulariaceae	D
Crepis	Asteraceae	D	Deschampsia	Poaceae	M
<i>Critesion</i>	Poaceae	M	Desmodium	Fabaceae	D
Crocsmia	Iridaceae	M	<i>Devauxia</i>	Centrolepidaceae	M
Cryptandra	Rhamnaceae	D	Deyeuxia	Poaceae	M
<i>Cryptostemma</i>	Asteraceae	D	Dianella	Liliaceae	M
Cryptostylis	Orchidaceae	M	Dianthus	Caryophyllaceae	D
<i>Ctenopteris</i>	Grammitidaceae	P	<i>Dichanthium</i>	Poaceae	M
Cucumis	Cucurbitaceae	D	Dichelachne	Poaceae	M
<i>Culcita</i>	Culcitaceae	P	Dichondra	Convolvulaceae	D

<i>Dichopogon</i>	Liliaceae	M	Echium	Boraginaceae	D
Dichosciadium	Apiaceae	D	Egeria	Hydrocharitaceae	M
Dicksonia	Dicksoniaceae	P	Ehrharta	Poaceae	M
<i>Didiscus</i>	Apiaceae	D	Einadia	Chenopodiaceae	D
<i>Didymotheca</i>	Gyrostemonaceae	D	Elaeocarpus	Elaeocarpaceae	D
Digitalis	Scrophulariaceae	D	Elatine	Elatinaceae	D
Digitaria	Poaceae	M	Eleocharis	Cyperaceae	M
<i>Dillenia</i>	Dilleniaceae	D	Eleusine	Poaceae	M
Dillwynia	Fabaceae	D	Elodea	Hydrocharitaceae	M
Dimorphotheca	Asteraceae	D	<i>Elymus</i>	Poaceae	M
Diplarrena	Iridaceae	M	<i>Elynanthus</i>	Cyperaceae	M
Diplaspis	Apiaceae	D	Elytrigia	Poaceae	M
<i>Diplax</i>	Poaceae	M	<i>Embothrium</i>	Proteaceae	D
Diplazium	Athyriaceae	P	Emex	Polygonaceae	D
<i>Diplodium</i>	Orchidaceae	M	Empodisma	Restionaceae	M
Diplotaxis	Brassicaceae	D	Enneapogon	Poaceae	M
Dipodium	Orchidaceae	M	Epacris	Epacridaceae	D
Dipogon	Fabaceae	D	Epilobium	Onagraceae	D
Dipsacus	Dipsacaceae	D	<i>Epipactis</i>	Orchidaceae	M
Disa	Orchidaceae	M	Equisetum	Equisetaceae	P
<i>Disarrenum</i>	Poaceae	M	Eragrostis	Poaceae	M
Discaria	Rhamnaceae	D	<i>Erechtites</i>	Asteraceae	D
Diselma	Cupressaceae	G	Erica	Ericaceae	D
<i>Disperis</i>	Orchidaceae	M	Erigeron	Asteraceae	D
Disphyma	Aizoaceae	D	Eriochilus	Orchidaceae	M
Distichlis	Poaceae	M	<i>Eriostemon</i>	Rutaceae	D
Dittrichia	Asteraceae	D	Erodium	Geraniaceae	D
Diuris	Orchidaceae	M	Erophila	Brassicaceae	D
Dockrillia	Orchidaceae	M	<i>Eruca</i>	Brassicaceae	D
Dodonaea	Sapindaceae	D	Eryngium	Apiaceae	D
<i>Dolichos</i>	Fabaceae	D	<i>Erythraea</i>	Gentianaceae	D
Donatia	Donatiaceae	D	Erythranthe	Scrophulariaceae	D
<i>Doodia</i>	Blechnaceae	P	<i>Erythranthera</i>	Poaceae	M
Draba	Brassicaceae	D	Eschscholzia	Papaveraceae	D
Dracophyllum	Epacridaceae	D	Eucalyptus	Myrtaceae	D
<i>Drapetes</i>	Thymelaeaceae	D	Euchiton	Asteraceae	D
<i>Drimys</i>	Winteraceae	D	Eucryphia	Eucryphiaceae	D
Drosanthemum	Aizoaceae	D	Euonymus	Celastraceae	D
Drosera	Droseraceae	D	<i>Eupatorium</i>	Asteraceae	D
Drymophila	Liliaceae	M	Euphorbia	Euphorbiaceae	D
Dryopoa	Poaceae	M	Euphrasia	Scrophulariaceae	D
Dysphania	Chenopodiaceae	D	<i>Eurybia</i>	Asteraceae	D
Eballium	Cucurbitaceae	D	Eurychorda	Restionaceae	M
Echinochloa	Poaceae	M	Euryomyrtus	Myrtaceae	D
Echinopogon	Poaceae	M	Euryops	Asteraceae	D

Eutaxia	Fabaceae	D	Glossodia	Orchidaceae	M
Ewartia	Asteraceae	D	Glossostigma	Scrophulariaceae	D
<i>Exacum</i>	Gentianaceae	D	Glyceria	Poaceae	M
<i>Exarrhena</i>	Boraginaceae	D	Glycine	Fabaceae	D
Exocarpos	Santalaceae	D	Gnaphalium	Asteraceae	D
<i>Fabricia</i>	Myrtaceae	D	Gomphocarpus	Asclepiadaceae	D
<i>Fagus</i>	Fagaceae	D	Gompholobium	Fabaceae	D
Fallopia	Polygonaceae	D	Gonocarpus	Haloragaceae	D
<i>Felicia</i>	Asteraceae	D	Goodenia	Goodeniaceae	D
Festuca	Poaceae	M	Goodia	Fabaceae	D
Ficinia	Cyperaceae	M	<i>Grammitis</i>	Grammitidaceae	P
<i>Filago</i>	Asteraceae	D	Gratiola	Scrophulariaceae	D
<i>Fitzroya</i>	Cupressaceae	G	Grevillea	Proteaceae	D
Foeniculum	Apiaceae	D	Griselinia	Cornaceae	D
Forstera	Stylidiaceae	D	Gunnera	Gunneraceae	D
Frankenia	Frankeniaceae	D	<i>Gunnia</i>	Orchidaceae	M
Fraxinus	Oleaceae	D	<i>Gymnogramma</i>	Hemionitidiaceae	P
Freesia	Iridaceae	M	Gymnoschoenus	Cyperaceae	M
<i>Frenela</i>	Cupressaceae	G	Gynatrix	Malvaceae	D
Fuchsia	Onagraceae	D	Gypsophila	Caryophyllaceae	D
Fumaria	Fumariaceae	D	Gyrostemon	Gyrostemonaceae	D
<i>Fuscospora</i>	Fagaceae	D	Haemodorum	Haemodoraceae	M
Gahnia	Cyperaceae	M	Hainardia	Poaceae	M
Gaimardia	Centrolepidaceae	M	Hakea	Proteaceae	D
Galenia	Aizoaceae	D	Halophila	Hydrocharitaceae	M
Galinsoga	Asteraceae	D	Haloragis	Haloragaceae	D
Galium	Rubiaceae	D	Hardenbergia	Fabaceae	D
Gamochaeta	Asteraceae	D	<i>Hebe</i>	Scrophulariaceae	D
Gastridium	Poaceae	M	Hedera	Araliaceae	D
Gastrodia	Orchidaceae	M	Hedycarya	Monimiaceae	D
Gaudinia	Poaceae	M	<i>Hedypnois</i>	Asteraceae	D
Gaultheria	Ericaceae	D	<i>Hedysarum</i>	Fabaceae	D
Gazania	Asteraceae	D	Helichrysum	Asteraceae	D
Genista	Fabaceae	D	<i>Helipterum</i>	Asteraceae	D
<i>Genoplesium</i>	Orchidaceae	M	<i>Helminthia</i>	Asteraceae	D
<i>Genosiris</i>	Iridaceae	M	Helminthotheca	Asteraceae	D
<i>Gentiana</i>	Gentianaceae	D	<i>Helxine</i>	Urticaceae	D
Gentianella	Gentianaceae	D	Hemarthria	Poaceae	M
Geococcus	Brassicaceae	D	Hemichroa	Amaranthaceae	D
Geranium	Geraniaceae	D	Herpolirion	Liliaceae	M
Geum	Rosaceae	D	Hesperantha	Iridaceae	M
<i>Githago</i>	Caryophyllaceae	D	Hesperocyparis	Cupressaceae	G
Gladiolus	Iridaceae	M	Heterozostera	Zosteraceae	M
Glaucium	Papaveraceae	D	<i>Hewardia</i>	Iridaceae	M
Gleichenia	Gleicheniaceae	P	Hibbertia	Dilleniaceae	D

<i>Hibiscus</i>	Malvaceae	D	<i>Jonesiopsis</i>	Orchidaceae	M
<i>Hieracium</i>	Asteraceae	D	Juncus	Juncaceae	M
Hierochloa	Poaceae	M	Kelleria	Thymelaeaceae	D
Hirschfeldia	Brassicaceae	D	Kennedia	Fabaceae	D
Histiopteris	Dennstaedtiaceae	P	Kickxia	Scrophulariaceae	D
Holcus	Poaceae	M	Kniphofia	Aloeaceae	M
<i>Homeria</i>	Iridaceae	M	Koeleria	Poaceae	M
Hookerochloa	Poaceae	M	<i>Kohlrauschia</i>	Caryophyllaceae	D
Hordeum	Poaceae	M	Kunzea	Myrtaceae	D
Hornungia	Brassicaceae	D	Laburnum	Fabaceae	D
Hovea	Fabaceae	D	Lachnagrostis	Poaceae	M
Humulus	Cannabaceae	D	Lactuca	Asteraceae	D
Huperzia	Lycopodiaceae	P	<i>Lagarosiphon</i>	Hydrocharitaceae	M
<i>Hutchinsia</i>	Brassicaceae	D	Lagarostrobos	Podocarpaceae	G
Hyalosperma	Asteraceae	D	Lagenophora	Asteraceae	D
<i>Hydatella</i>	Hydatellaceae	M	Lagurus	Poaceae	M
Hydrocotyle	Apiaceae	D	Lamium	Lamiaceae	D
Hydrorchis	Orchidaceae	M	<i>Lampocarya</i>	Cyperaceae	M
<i>Hylogyne</i>	Proteaceae	D	Lampranthus	Aizoaceae	D
<i>Hymenanthera</i>	Violaceae	D	Landoltia	Lemnaceae	M
<i>Hymenochilus</i>	Orchidaceae	M	Lapsana	Asteraceae	D
<i>Hymenolobus</i>	Brassicaceae	D	Lasiopetalum	Sterculiaceae	D
Hymenophyllum	Hymenophyllaceae	P	Lasiospermum	Asteraceae	D
<i>Hyoscyamus</i>	Solanaceae	D	Lastreopsis	Dryopteridaceae	P
Hypericum	Clusiaceae	D	Lathyrus	Fabaceae	D
Hypochaeris	Asteraceae	D	Lavandula	Lamiaceae	D
Hypolaena	Restionaceae	M	<i>Lavatera</i>	Malvaceae	D
Hypolepis	Dennstaedtiaceae	P	Lawrencia	Malvaceae	D
Hypoxis	Liliaceae	M	Laxmannia	Liliaceae	M
Ilex	Aquifoliaceae	D	Leiocarpa	Asteraceae	D
Imperata	Poaceae	M	Leionema	Rutaceae	D
Indigofera	Fabaceae	D	Lemna	Lemnaceae	M
<i>Inula</i>	Asteraceae	D	Leonotis	Lamiaceae	D
Ipomoea	Convolvulaceae	D	Leontodon	Asteraceae	D
Iris	Iridaceae	M	Lepidium	Brassicaceae	D
Isachne	Poaceae	M	Lepidosperma	Cyperaceae	M
Isoetes	Isoetaceae	P	Lepilaena	Zannichelliaceae	M
Isoetopsis	Asteraceae	D	Leptecophylla	Epacridaceae	D
Isolepis	Cyperaceae	M	Leptinella	Asteraceae	D
Isophysis	Iridaceae	M	Leptocarpus	Restionaceae	M
Isopogon	Proteaceae	D	Leptoceras	Orchidaceae	M
Isotoma	Campanulaceae	D	<i>Leptocyamus</i>	Fabaceae	D
Ixia	Iridaceae	M	Leptomeria	Santalaceae	D
<i>Ixiolaena</i>	Asteraceae	D	Leptorhynchus	Asteraceae	D
<i>Ixodia</i>	Asteraceae	D	Leptospermum	Myrtaceae	D

<i>Lepturus</i>	Poaceae	M	<i>Lyonsia</i>	Apocynaceae	D
Lepyrodia	Restionaceae	M	Lyperanthus	Orchidaceae	M
Leucanthemum	Asteraceae	D	Lysimachia	Primulaceae	D
Leucochrysum	Asteraceae	D	Lythrum	Lythraceae	D
Leucojum	Liliaceae	M	<i>Malaxis</i>	Orchidaceae	M
Leucophyta	Asteraceae	D	Malus	Rosaceae	D
Leucopogon	Epacridaceae	D	Malva	Malvaceae	D
Levenhookia	Stylidiaceae	D	<i>Marianthus</i>	Pittosporaceae	D
Leycesteria	Caprifoliaceae	D	Marrubium	Lamiaceae	D
Leymus	Poaceae	M	Marsilea	Marsileaceae	P
Libertia	Iridaceae	M	Matricaria	Asteraceae	D
Ligustrum	Oleaceae	D	Matthiola	Brassicaceae	D
Lilaeopsis	Apiaceae	D	Maytenus	Celastraceae	D
<i>Limnanthemum</i>	Menyanthaceae	D	Mazus	Scrophulariaceae	D
Limonium	Plumbaginaceae	D	<i>Mecodium</i>	Hymenophyllaceae	P
Limosella	Scrophulariaceae	D	Medicago	Fabaceae	D
Linaria	Scrophulariaceae	D	<i>Meionectes</i>	Haloragaceae	D
Lindsaea	Lindsaeaceae	P	Melaleuca	Myrtaceae	D
<i>Linguella</i>	Orchidaceae	M	<i>Melandrium</i>	Caryophyllaceae	D
Linum	Linaceae	D	Melicytus	Violaceae	D
Liparophyllum	Menyanthaceae	D	Melilotus	Fabaceae	D
Lissanthe	Epacridaceae	D	Melissa	Lamiaceae	D
Lithospermum	Boraginaceae	D	Mentha	Lamiaceae	D
<i>Litobrochia</i>	Pteridaceae	P	Mesembryanthemum	Aizoaceae	D
Lobelia	Campanulaceae	D	<i>Mesomelaena</i>	Cyperaceae	M
Lobularia	Brassicaceae	D	Micranthemum	Euphorbiaceae	D
Logfia	Asteraceae	D	Microcachrys	Podocarpaceae	G
Lolium	Poaceae	M	Microlaena	Poaceae	M
Lomandra	Xanthorrhoeaceae	M	Microseris	Asteraceae	D
<i>Lomaria</i>	Blechnaceae	P	Microsorium	Polypodiaceae	P
Lomatia	Proteaceae	D	<i>Microstrobos</i>	Podocarpaceae	G
Lonicera	Caprifoliaceae	D	Microtidium	Orchidaceae	M
<i>Lophochloa</i>	Poaceae	M	Microtis	Orchidaceae	M
<i>Lophopyrum</i>	Poaceae	M	Milligania	Liliaceae	M
<i>Lophozonia</i>	Fagaceae	D	Millotia	Asteraceae	D
Lotus	Fabaceae	D	<i>Mimosa</i>	Mimosaceae	D
Lunaria	Brassicaceae	D	<i>Mimulus</i>	Scrophulariaceae	D
Lupinus	Fabaceae	D	Minuartia	Caryophyllaceae	D
Luzula	Juncaceae	M	Mirabilis	Nyctaginaceae	D
<i>Lychnis</i>	Caryophyllaceae	D	Mirbelia	Fabaceae	D
Lycium	Solanaceae	D	Mitrasacme	Loganiaceae	D
Lycopodiella	Lycopodiaceae	P	<i>Mniarum</i>	Caryophyllaceae	D
Lycopodium	Lycopodiaceae	P	Modiola	Malvaceae	D
<i>Lycopsis</i>	Boraginaceae	D	Moenchia	Caryophyllaceae	D
Lycopus	Lamiaceae	D	Molineriella	Poaceae	M

<i>Monerma</i>	Poaceae	M	Ophioglossum	Ophioglossaceae	P
Monotoca	Epacridaceae	D	Oreobolus	Cyperaceae	M
Montia	Portulacaceae	D	Oreomyrrhis	Apiaceae	D
Montitega	Epacridaceae	D	<i>Oreoporanthera</i>	Euphorbiaceae	D
Moraea	Iridaceae	M	Orites	Proteaceae	D
Muehlenbeckia	Polygonaceae	D	Ornduffia	Menyanthaceae	D
Muscari	Liliaceae	M	Ornithogalum	Liliaceae	M
Myoporum	Myoporaceae	D	Ornithopus	Fabaceae	D
Myosotis	Boraginaceae	D	Orobanche	Orobanchaceae	D
Myosurus	Ranunculaceae	D	Orthoceras	Orchidaceae	M
<i>Myriogyne</i>	Asteraceae	D	<i>Oryzopsis</i>	Poaceae	M
Myriophyllum	Haloragaceae	D	Oschatzia	Apiaceae	D
<i>Myrsiphyllum</i>	Liliaceae	M	<i>Osteospermum</i>	Asteraceae	D
Nablonium	Asteraceae	D	Ourisia	Scrophulariaceae	D
<i>Nanozostera</i>	Zosteraceae	M	Oxalis	Oxalidaceae	D
Narcissus	Liliaceae	M	Oxylobium	Fabaceae	D
Nardus	Poaceae	M	Ozothamnus	Asteraceae	D
Nassella	Poaceae	M	Pachycladon	Brassicaceae	D
Nasturtium	Brassicaceae	D	<i>Panax</i>	Araliaceae	D
Navarretia	Polemoniaceae	D	Pandorea	Bignoniaceae	D
<i>Nemacianthus</i>	Orchidaceae	M	Panicum	Poaceae	M
<i>Nematoceras</i>	Orchidaceae	M	Papaver	Papaveraceae	D
Nematolepis	Rutaceae	D	Pappochroma	Asteraceae	D
<i>Neopaxia</i>	Portulacaceae	D	<i>Paquerina</i>	Asteraceae	D
Nepeta	Lamiaceae	D	Paracaleana	Orchidaceae	M
<i>Nephrodium</i>	Dryopteridaceae	P	<i>Parahebe</i>	Scrophulariaceae	D
Nertera	Rubiaceae	D	Parapholis	Poaceae	M
Nicotiana	Solanaceae	D	Paraserianthes	Mimosaceae	D
Nonea	Boraginaceae	D	Parentucellia	Scrophulariaceae	D
Notelaea	Oleaceae	D	Parietaria	Urticaceae	D
Nothofagus	Fagaceae	D	Paronychia	Caryophyllaceae	D
<i>Nothopanax</i>	Araliaceae	D	Parsonsia	Apocynaceae	D
Nothoscordum	Liliaceae	M	<i>Paspalidium</i>	Poaceae	M
<i>Notodanthonia</i>	Poaceae	M	Paspalum	Poaceae	M
Notogrammitis	Grammitidaceae	P	Passiflora	Passifloraceae	D
<i>Nymphoides</i>	Menyanthaceae	D	Pastinaca	Apiaceae	D
Odixia	Asteraceae	D	Patersonia	Iridaceae	M
Oenothera	Onagraceae	D	Pauridia	Liliaceae	M
Olearia	Asteraceae	D	Pelargonium	Geraniaceae	D
<i>Oligochaetochilus</i>	Orchidaceae	M	Pellaea	Adiantaceae	P
<i>Onobrychis</i>	Fabaceae	D	<i>Pennisetum</i>	Poaceae	M
<i>Onoclea</i>	Blechnaceae	P	Pentachondra	Epacridaceae	D
Ononis	Fabaceae	D	Pentaglottis	Boraginaceae	D
Onopordum	Asteraceae	D	Pentapogon	Poaceae	M
Opercularia	Rubiaceae	D	<i>Pernettya</i>	Ericaceae	D

Persicaria	Polygonaceae	D	Podalyria	Fabaceae	D
Persoonia	Proteaceae	D	Podocarpus	Podocarpaceae	G
<i>Petalochilus</i>	Orchidaceae	M	Podolepis	Asteraceae	D
Petasites	Asteraceae	D	<i>Podosperma</i>	Asteraceae	D
Petrorhagia	Caryophyllaceae	D	Podotheca	Asteraceae	D
Petroselinum	Apiaceae	D	Polycarpon	Caryophyllaceae	D
Phacelia	Hydrophyllaceae	D	Polygala	Polygalaceae	D
Phalaris	Poaceae	M	Polygonum	Polygonaceae	D
Phebalium	Rutaceae	D	Polyphlebium	Hymenophyllaceae	P
Pheladenia	Orchidaceae	M	<i>Polypodium</i>	Grammitidaceae	P
Pherosphaera	Podocarpaceae	G	<i>Polypodium</i>	Polypodiaceae	P
Philotheca	Rutaceae	D	Polypogon	Poaceae	M
Phlegmariusus	Lycopodiaceae	P	<i>Polypompholyx</i>	Lentibulariaceae	D
Phleum	Poaceae	M	Polyscias	Araliaceae	D
Phormium	Agavaceae	M	Polystichum	Dryopteridaceae	P
Phragmites	Poaceae	M	Pomaderris	Rhamnaceae	D
Phyllachne	Stylidiaceae	D	Populus	Salicaceae	D
Phyllangium	Loganiaceae	D	Poranthera	Euphorbiaceae	D
Phyllanthus	Euphorbiaceae	D	Portulaca	Portulacaceae	D
Phyllocladus	Podocarpaceae	G	Posidonia	Posidoniaceae	M
Phylloglossum	Lycopodiaceae	P	Potamogeton	Potamogetonaceae	M
Phyllota	Fabaceae	D	Potentilla	Rosaceae	D
<i>Phymatodes</i>	Polypodiaceae	P	<i>Poterium</i>	Rosaceae	D
<i>Phymatosorus</i>	Polypodiaceae	P	Prasophyllum	Orchidaceae	M
Physalis	Solanaceae	D	<i>Pratia</i>	Campanulaceae	D
Phytolacca	Phytolaccaceae	D	Prionotes	Epacridaceae	D
Picris	Asteraceae	D	Prostanthera	Lamiaceae	D
<i>Pilitis</i>	Epacridaceae	D	Prunella	Lamiaceae	D
Pilosella	Asteraceae	D	Prunus	Rosaceae	D
Pilularia	Marsileaceae	P	<i>Psamma</i>	Poaceae	M
Pimelea	Thymelaeaceae	D	Pseudanthus	Euphorbiaceae	D
Pinus	Pinaceae	G	<i>Pseudofumaria</i>	Fumariaceae	D
Piptatherum	Poaceae	M	<i>Pseudognaphalium</i>	Asteraceae	D
Pittosporum	Pittosporaceae	D	Pseudopanax	Araliaceae	D
<i>Plagianthus</i>	Malvaceae	D	<i>Psilotum</i>	Psilotaceae	P
Planocarpa	Epacridaceae	D	Psoralea	Fabaceae	D
Plantago	Plantaginaceae	D	Psychrophila	Ranunculaceae	D
Platylobium	Fabaceae	D	Pteridium	Dennstaedtiaceae	P
<i>Platyloma</i>	Adiantaceae	P	Pteris	Pteridaceae	P
<i>Pleurandra</i>	Dilleniaceae	D	Pterostylis	Orchidaceae	M
Pleurophyllum	Asteraceae	D	Pterygopappus	Asteraceae	D
Pleurosorus	Aspleniaceae	P	Ptilotus	Amaranthaceae	D
<i>Plumatichilos</i>	Orchidaceae	M	Puccinellia	Poaceae	M
Pneumatopteris	Thelypteridaceae	P	Pultenaea	Fabaceae	D
Poa	Poaceae	M	<i>Pygmea</i>	Scrophulariaceae	D

Pyrorchis	Orchidaceae	M	Scaevola	Goodeniaceae	D
<i>Racosperma</i>	Mimosaceae	D	Scandix	Apiaceae	D
Ranunculus	Ranunculaceae	D	<i>Schedonorus</i>	Poaceae	M
<i>Raoulia</i>	Asteraceae	D	Schenkia	Gentianaceae	D
Raphanus	Brassicaceae	D	Schizacme	Loganiaceae	D
Rapistrum	Brassicaceae	D	Schizaea	Schizaeaceae	P
Reseda	Resedaceae	D	<i>Schizostylis</i>	Iridaceae	M
<i>Restio</i>	Restionaceae	M	<i>Schoenodum</i>	Restionaceae	M
<i>Reynoutria</i>	Polygonaceae	D	<i>Schoenoides</i>	Cyperaceae	M
<i>Rhagadiolus</i>	Asteraceae	D	Schoenoplectus	Cyperaceae	M
Rhagodia	Chenopodiaceae	D	Schoenus	Cyperaceae	M
Rhamnus	Rhamnaceae	D	Scilla	Liliaceae	M
Rhodanthe	Asteraceae	D	<i>Scirpus</i>	Cyperaceae	M
Rhododendron	Ericaceae	D	Scleranthus	Caryophyllaceae	D
Rhytidosporum	Pittosporaceae	D	Sclerochloa	Poaceae	M
Ribes	Grossulariaceae	D	<i>Scleroleima</i>	Asteraceae	D
<i>Richea</i>	Asteraceae	D	<i>Sclerostegia</i>	Chenopodiaceae	D
Richea	Epacridaceae	D	Scorzonera	Asteraceae	D
Ricinocarpus	Euphorbiaceae	D	Scrophularia	Scrophulariaceae	D
<i>Roepera</i>	Zygophyllaceae	D	Scutellaria	Lamiaceae	D
Romulea	Iridaceae	M	Sebaea	Gentianaceae	D
Rorippa	Brassicaceae	D	Secale	Poaceae	M
Rosa	Rosaceae	D	Securigera	Fabaceae	D
Rostraria	Poaceae	M	Sedum	Crassulaceae	D
Rubus	Rosaceae	D	Selaginella	Selaginellaceae	P
Rumex	Polygonaceae	D	Selliera	Goodeniaceae	D
Rumohra	Dryopteridaceae	P	<i>Senebiera</i>	Brassicaceae	D
Ruppia	Ruppiaceae	M	Senecio	Asteraceae	D
<i>Rutidosis</i>	Asteraceae	D	Setaria	Poaceae	M
Rytidosperma	Poaceae	M	<i>Sheffieldia</i>	Primulaceae	D
Sagina	Caryophyllaceae	D	Sherardia	Rubiaceae	D
<i>Salicornia</i>	Chenopodiaceae	D	Sicyos	Cucurbitaceae	D
Salix	Salicaceae	D	<i>Sieglingia</i>	Poaceae	M
Salpichroa	Solanaceae	D	Sigesbeckia	Asteraceae	D
Salsola	Chenopodiaceae	D	Silene	Caryophyllaceae	D
Salvia	Lamiaceae	D	Siloxerus	Asteraceae	D
Sambucus	Caprifoliaceae	D	Silybum	Asteraceae	D
Samolus	Primulaceae	D	<i>Simpliglottis</i>	Orchidaceae	M
Sanguisorba	Rosaceae	D	Sinapis	Brassicaceae	D
Saponaria	Caryophyllaceae	D	Sisymbrium	Brassicaceae	D
Sarcochilus	Orchidaceae	M	Sisyrrinchium	Iridaceae	M
Sarcocornia	Chenopodiaceae	D	<i>Skirrhophorus</i>	Asteraceae	D
<i>Sarothamnus</i>	Fabaceae	D	Solanum	Solanaceae	D
Saxipoa	Poaceae	M	Soleirolia	Urticaceae	D
Scabiosa	Dipsacaceae	D	Solenogyne	Asteraceae	D



Soliva	Asteraceae	D	Syzygium	Myrtaceae	D
<i>Sollya</i>	Pittosporaceae	D	Tanacetum	Asteraceae	D
Sonchus	Asteraceae	D	Taraxacum	Asteraceae	D
Sorbus	Rosaceae	D	Tasmania	Winteraceae	D
Sorghum	Poaceae	M	<i>Taurantha</i>	Orchidaceae	M
Sowerbaea	Liliaceae	M	<i>Taxanthemea</i>	Plumbaginaceae	D
Sparaxis	Iridaceae	M	Tecticornia	Chenopodiaceae	D
Spartina	Poaceae	M	Teesdalia	Brassicaceae	D
Spartium	Fabaceae	D	Telopea	Proteaceae	D
<i>Speculantha</i>	Orchidaceae	M	Tetracarpaea	Escalloniaceae	D
Spergula	Caryophyllaceae	D	Tetragonia	Aizoaceae	D
Spergularia	Caryophyllaceae	D	Tetralia	Cyperaceae	M
<i>Sphaerocionium</i>	Hymenophyllaceae	P	Tetrarrhena	Poaceae	M
Sphaerolobium	Fabaceae	D	Tetratheca	Tremandraceae	D
<i>Spiculaea</i>	Orchidaceae	M	Teucrium	Lamiaceae	D
Spinifex	Poaceae	M	Thelionema	Liliaceae	M
Spiranthes	Orchidaceae	M	Thelymitra	Orchidaceae	M
<i>Spirodela</i>	Lemnaceae	M	Themeda	Poaceae	M
Sporadanthus	Restionaceae	M	Thesium	Santalaceae	D
Sporobolus	Poaceae	M	Thinopyrum	Poaceae	M
Sprengelia	Epacridaceae	D	Thismia	Burmanniaceae	M
Spyridium	Rhamnaceae	D	Thlaspi	Brassicaceae	D
Stachys	Lamiaceae	D	<i>Thonandia</i>	Poaceae	M
Stackhousia	Stackhousiaceae	D	Threlkeldia	Chenopodiaceae	D
<i>Statice</i>	Plumbaginaceae	D	Thryptomene	Myrtaceae	D
<i>Stegania</i>	Blechnaceae	P	Thynninorchis	Orchidaceae	M
<i>Stegostyla</i>	Orchidaceae	M	Thyridia	Scrophulariaceae	D
Stellaria	Caryophyllaceae	D	Thysanotus	Liliaceae	M
Stenanthemum	Rhamnaceae	D	<i>Tillaea</i>	Crassulaceae	D
<i>Stenanthera</i>	Epacridaceae	D	Tmesipteris	Psilotaceae	P
Stenopetalum	Brassicaceae	D	Todea	Osmundaceae	P
Stenotaphrum	Poaceae	M	Tolpis	Asteraceae	D
Sticherus	Gleicheniaceae	P	Torilis	Apiaceae	D
Stilbocarpa	Apiaceae	D	Townsonia	Orchidaceae	M
<i>Stipa</i>	Poaceae	M	Trachymene	Apiaceae	D
Stonesiella	Fabaceae	D	Tradescantia	Commelinaceae	M
Stuckenia	Potamogetonaceae	M	Tragopogon	Asteraceae	D
Stylidium	Stylidiaceae	D	<i>Trichinium</i>	Amaranthaceae	D
<i>Stypantra</i>	Liliaceae	M	<i>Trichomanes</i>	Hymenophyllaceae	P
Styphelia	Epacridaceae	D	<i>Trichonema</i>	Iridaceae	M
Suaeda	Chenopodiaceae	D	Tricoryne	Liliaceae	M
<i>Sullivania</i>	Orchidaceae	M	Tricostularia	Cyperaceae	M
Swainsona	Fabaceae	D	Trifolium	Fabaceae	D
Symphyotrichum	Asteraceae	D	Triglochin	Juncaginaceae	M
Symphytum	Boraginaceae	D	<i>Trigonella</i>	Fabaceae	D

<i>Triodia</i>	Poaceae	M	<i>Villarsia</i>	Menyanthaceae	D
Tripleurospermum	Asteraceae	D	Viminaria	Fabaceae	D
Triptilodiscus	Asteraceae	D	Vinca	Apocynaceae	D
Trisetum	Poaceae	M	Viola	Violaceae	D
Trithuria	Hydatellaceae	M	Vittadinia	Asteraceae	D
Triticum	Poaceae	M	Vulpia	Poaceae	M
Tritonia	Iridaceae	M	Wahlenbergia	Campanulaceae	D
Trochocarpa	Epacridaceae	D	Watsonia	Iridaceae	M
Tropaeolum	Tropaeolaceae	D	Westringia	Lamiaceae	D
Typha	Typhaceae	M	Wilsonia	Convolvulaceae	D
Ulex	Fabaceae	D	Winifredia	Restionaceae	M
Ulmus	Ulmaceae	D	Wolffia	Lemnaceae	M
Uncinia	Cyperaceae	M	Wurmbea	Liliaceae	M
<i>Uniola</i>	Poaceae	M	Xanthium	Asteraceae	D
<i>Urochilus</i>	Orchidaceae	M	Xanthorrhoea	Xanthorrhoeaceae	M
Urospermum	Asteraceae	D	Xanthosia	Apiaceae	D
Urtica	Urticaceae	D	Xerochrysum	Asteraceae	D
Utricularia	Lentibulariaceae	D	<i>Xerotes</i>	Xanthorrhoeaceae	M
<i>Vaccaria</i>	Caryophyllaceae	D	Xyris	Xyridaceae	M
Valerianella	Valerianaceae	D	<i>Zannichellia</i>	Zannichelliaceae	M
Vallisneria	Hydrocharitaceae	M	Zantedeschia	Araceae	M
<i>Valvaria</i>	Ranunculaceae	D	Zieria	Rutaceae	D
Velleia	Goodeniaceae	D	Zostera	Zosteraceae	M
Vellereophyton	Asteraceae	D	Zoysia	Poaceae	M
Verbascum	Scrophulariaceae	D	Zygophyllum	Zygophyllaceae	D
Verbena	Verbenaceae	D	<i>*Agropogon</i>	Poaceae	M
Veronica	Scrophulariaceae	D	<i>*Calassodia</i>	Orchidaceae	M
Viburnum	Caprifoliaceae	D	<i>*Glossadenia</i>	Orchidaceae	M
Vicia	Fabaceae	D	<i>*Taurodium</i>	Orchidaceae	M